

BSBI Gentian Handbook

Tim Rich & Andy McVeigh



Gentians of Britain and Ireland

Tim Rich & Andy McVeigh

BSBI Handbook no. 19

4. *Centaurium pulchellum* (Sw.) Druce Lesser Centaury

Annual 2.05–2.5(3) cm tall, erect, light greenish-green, glabrous. **Basal rosette** sometimes present or more usually withered at flowering (or sometimes shortly defined) with 2–3 pairs of leaves. **Basal leaves** 2–5 × 1.5–3 mm, broadly elliptic to ovate, acute, entire, 1-veined, broadly but shortly petiolate. **Stems** 6-angled, veined above, stipule-like branches mostly above. **Internodes** 0.5–2 × as long as their leaves. **Stem leaves** 13–20(–26) × as long as wide, 1 strong central vein and 2 weak lateral, acute or rarely obtuse, acute and scabrous stem but not coriaceous, increasing in size up the stem. Inflorescence simple or cymose, open and lax to dense. **Terminal pedicel** 1.5–10(–11) mm, lateral pedicels often longer. **Flowers** 6–8 mm across (8–10 mm in small plants or on lateral branches of big plants), 4–5 mm across. **Buds** whitish with calyx lobes uncontracted. **Calyx** 9–10(–10) mm, calyx lobes 3–4(–4) mm, rounded shaped, calyx lobes 12.5–16.7 mm, linear, acute, appressed, equal in size, membranous margins nearer base, acute on angles but not serrated. **Corolla** 12.5 nearly as long as corolla tube. **Corolla** 10.5–15(–16), 15.5 mm, corolla tube (6–10–12 mm cylindrical but slightly narrowed at apex, whitish, not noticeably veined, lobes 2–10(–15) mm × 3–2 mm, pink (rarely white), spreading, obtuse. **Corolla** 1.4 × 1.9(–2.2) × as long as calyx. **Anthers** twisted when deflexed. **Filaments** whitish, inserted at top of corolla tube. **Mean pollen size** 28.7–45.53 µm (range 26.1–30.6 µm). **Ovary** ellipsoid, style 2–3 mm, linear, stigma 2-lobed, corollate, yellowish-green. **Capsule** cylindrical, seeds 0.3–0.4 mm, brown, 2-ovoid, reticulate.

Chromosome numbers: 2n = 4, 34 (1 count Britain) and 36 (1 count British Isles) (BOB Cytology database). Other counts suggest it is consistently tetraploid 2n=36 with one diploid 2n=18 count from Israel (Zohary 1970, 1985).

Centaurium pulchellum is characterized by the annual habit, the few pairs of ovate stem leaves (mostly 2–4, sometimes up to 5) and the small, deep pink corolla (2–14 mm long with lobes 2–3 mm long (Figure 77)). The very small flowers 5–7 mm across are usually instantly recognizable.

It has sometimes been confused with *C. erythraea*, dwarf examples of which are not always easy to distinguish. In *C. pulchellum* the flowers are usually stalked (terminal pedicel 1–10 mm) and the basal rosette is usually withered at flowering. In *C. erythraea*, the flowers usually more or less sessile or shortly stalked (terminal pedicel to 1(–2) mm but usually more or less sessile) and the basal rosette usually still present even if remaining at flowering.

In coastal dune slacks, the very small corolla and ovate stem leaves will separate it from *C. litoreum* (which has corolla 11–14–16 mm and linear stem leaves 4–10 × as long as wide).

In drier heath *C. tetragynum* is having fewer pairs of stem leaves and smaller corolla (stem leaves 5–12 pairs, corolla 11.5–14 mm corolla fresh origin) (Fig. 77) for *C. pulchellum* and the angles of (2001) note that the dist. is impossible in Europe, but there is.

For hybrids with *C. litoreum*, see *C. litoreum*.



Figure 77. Flowers of *C. erythraea* (left: larger, paler pink) and *C. pulchellum* (right: smaller deep pink).



Figure 77. Distribution of *C. pulchellum* in Britain and Ireland.

It has been recorded in s.v. 1–23, 25, 27–38, 41, 44–46, 48, 49, 51–56, 58–63, 65–69, 72, 73, 82 and the Channel Islands, and in Ireland in H2, H5, H6, H12, H14, H21. Introduced in s.v. 70.

Widespread in Europe from the A introduced to North and South Am.

Habitats

In southern England, *C. pulchellum* occurs on a range of open grasslands and disturbed habitats. In all cases in open rich with minimal competition. In calcareous grassland it can be locally abundant on chalky clays, sometimes being the sides of animal tracks on the Downs, or colonizing bare chalk in old quarries, etc. It may be scattered on drier sites or grass in the open parts of rides in woodlands and plantations. On heathland it can be locally frequent on open dry acidic, sandy ground and on damp organic soils in sedge zones. On the coast, it occurs in dune slacks, especially beside trampled paths, and on the upper edges of salt-marshes where turf cutting may provide an important regeneration niche (Ray 1972). On sea cliffs it can occur in small seepage zones. In Ireland it now mainly occurs in dune slacks and associated paths.

It is quite tolerant of brackish but not saline conditions, and in the northern parts of its range becomes noticeably more restricted to coastal habitats.

Reproductive biology

Centaurium pulchellum has been little studied. It is relatively annual in the British and Ireland in being a summer annual (rather than winter annual which it may be in southern Europe), and is thus prone to vary from year to year depending on the spring weather. In hot dry summers, such as 1990, it can be very scarce, and then often only found in the damper habitats. In wet summers, such as 2007, it can be abundant in a wider range of habitats.

It probably has a long persistent seed bank. Seeds germinate in late spring, and they pale green rosettes can be found in April and May. Plants flower from late to October, but flowers may not open in dull weather. It is self-compatible and probably mostly self-pollinated. The mean pollen viability was




Figure 77. World distribution of *C. pulchellum* (after Mansell et al. 1965).



Figure 77. World distribution of *C. pulchellum* (after Mansell et al. 1965).



Figure 77. Habitats of *C. pulchellum*. A, Dune slacks, Anglesey; B, heathland rabbit, Portland; C, damp heathland, Devon.

79.4 ± 5.62% (range 6–96%) (West et al. 2014).

Conservation statuses

EU28 (2002) Conservation status Britain: Least Concern (JNCC, 2018); Ireland: Near Threatened (Wynne-Jackson et al. 2015).

In Ireland *C. pulchellum* is a protected species listed on the Flora Protection Order. In Britain, it is much more frequent though loss of heathland has resulted in some decline in the south. The distribution overall in Britain is currently more or less stable (Braithwaite et al. 2006), or with the warmer summers may be increasing.



Figure 77. Variation in *Centaurium pulchellum*. A, large branched plant c. 20 cm tall; B, densely branched plant from base c. 5 cm tall; C, small plants, the smallest c. 2 cm tall.





Figure 77. Habitats of *C. pulchellum*. A, Dune slacks, Anglesey; B, heathland rabbit, Portland; C, damp heathland, Devon.

Also just published! See leaflet!



Why are Gentianaceae difficult taxonomically?

- Some taxa are defined by quantitative differences between sets of partially overlapping morphological characters
 - partly related to their origin by alloploidy or autopolyploidy (Mansion *et al.* 2005).
- Some taxa are of relatively recent origin and closely related, such as the *Gentianella amarella* subspecies
 - genetically adapted to local environment
- Marked differences in growth form due to local environmental conditions
- Marked population fluctuations of annuals and biennials, making comparison between years difficult on the same site.
- After some work on population dynamics, we are less sure that some populations may have both annuals and biennials (c.f. Pritchard 1959)
- Hybridisation and introgression occur in some taxa
 - Most hybrids fertile, few are sterile
- Very difficult to cultivate plants

- Also long-standing problem of floras being compiled from herbaria, which give different character ranges to those in field due to selection of material and drying

Table 6. Comparison of measurements collected in the field and herbaria for *Gentianella germanica*.

Character	Field			Herbarium			<i>P</i> (<i>t</i> -test)
	<i>n</i>	mean ± s.e.	(1–)10–90(–100) percentile range	<i>n</i>	mean ± s.e.	(1–)10–90(–100) percentile range	
Internodes	321	10.4 ± 0.12	(5–)8–13(–16)	627	9.3 ± 0.08	(4–)7–12(–16)	<0.001
Leaf length [mm]	321	24.2 ± 0.40	(6–)14–32(–43)	578	16.7 ± 0.20	(6–)11–24(–42)	<0.001
Leaf width [mm]	321	12.2 ± 0.20	(3–)7–16.5(–22)	578	7.6 ± 0.10	(3–)5–11(–17)	<0.001
Leaf length / leaf width ratio	321	2.02 ± 0.02	(1–)1.6–2.40(–3.8)	578	2.3 ± 0.02	(1.3–)1.7–2.9(–4.3)	<0.001
Corolla length [mm]	321	28.6 ± 0.20	(12–)25–32(–37)	796	26.2 ± 0.10	(15–)22–30(–35)	<0.001
Calyx length [mm]	321	15.8 ± 0.20	(5–)12–20(–24)	796	11.6 ± 0.07	(5–)9–14(–18)	<0.001
Corolla length / calyx length ratio	321	1.86 ± 0.02	(1–2–)1.5–2.2(–3.2)	796	2.3 ± 0.01	(1.5–)1.9–2.7(–5.1)	<0.001

Data of A. McVeigh.

Parnell, J., Rich, T., McVeigh, A., Lim, A., Quigley, S., Morris, D. & Wong, Z. (2013). The effect of preservation methods on plant morphology. *Taxon* **62**: 1259–1265.

The case of *Centaurium tenuiflorum*

- first collected by C. C. Babington in 1837 in Guernsey (site now destroyed)
- subsequently found on Isle of Wight in 1879 by F. Townsend and Dorset in 1935 by R. Good
- In UK a very rare species



In Europe 3 taxa (G. Mansion work):

- diploid subsp. *acutiflorum*
- tetraploid subsp. *tenuiflorum*
- + widespread European taxon, informally '*C. ×tenuiflorum*' (allotetraploid derivative of *C. tenuiflorum* × *C. erythraea*)



N Spain



Brittany



Spain (Stace)

Morphological data Dorset/I of Wight vs European material:

	Mean Europe	Mean Dorset/IoW	T test Probability	significance
height cm	17.5	17.3	0.870917	NS
No. internodes	8.1	6.3	0.004188	**
Mid stem leaf length mm	9.8	11.4	0.06141	*
Mid stem leaf width mm	2.1	3.6	8.34E-06	***
Mid stem leaf L W ratio	5.1	3.4	0.000728	**
Terminal pedicel length mm	2.1	3.3	0.074825	NS
calyx length mm	8.0	8.9	0.000503	**
corolla tube mm	9.7	10.6	0.005761	**
corolla lobes mm	3.0	3.6	7.94E-06	***
total corolla mm	12.0	14.2	0.004047	**
corolla calyx ratio	1.5	1.6	0.46553	NS

- Dorset/I of Wight plants are different – generally bigger with broader leaves, and retain these in cultivation.

DNA (based on G. Mansion's work)

- ITS1 Dorset sequence is *C. tenuiflorum* (differs in 1 base pair)

Cytology

- one Ubsdell count of UK material = diploid (yet pollen sizes suggests tetraploid)

Hence described treated as new English endemic = *C. tenuiflorum* subsp. *anglicum**

- presumed to have arisen through isolation and local adaptation following reflooding of the English Channel
- **Post Handbook:** Now seen very similar material from N and W France

* Rich, T. C. G., McVeigh, A. & Stace, C.A.(2018). New taxa and new combinations in the British flora. *Edinburgh Journal of Botany* doi: 10.1017/S0960428618000288

The case of *Centaureum erythraea x littorale*

- Wheldon (1897) noted two forms of *C. littorale* on the Lancashire coast
 - normal widespread form
 - large form with the relatively long calyx: corolla tube ratio of *C. littorale* but the broader stem leaves and habit of *C. erythraea* (*Erythraea littoralis* var. *intermedia*).
- These regarded as of hybrid origin and *C. x intermedium* widely used for sterile tetraploid *C. erythraea x littorale*
- Ubsdell (1976a, b, 1979) showed the large form was a hexaploid forming cytologically stable, self-sustaining populations which were reproductively isolated from the parents, and suggested it should be a new allopolyploid species
 - Typification shows *Erythraea littoralis* var. *intermedia* belongs to these
- CTW and Stace acknowledge these hexaploids but did not separate them
 - Stace ed 4 “The latter could be treated as a distinct new sp. but the parents themselves are so close this is not feasible.”

Courtesy of Phil Smith, we went to see the Lancashire plants in 2018



C. intermedium

- Tall (to 40 cm), fertile, hexaploid
- Lancashire endemic



C. erythraea x littorale = C. x klattii

- Small (to 15 cm), infertile, tetraploid
- UK, Europe




- Admittedly hard to separate in the field without fertility data.

The case of *Gentianella anglica*

- Since 1993 TR has been saying *G. anglica* is a species, differing from *G. amarella* in:
 - small size 1–15 cm tall
 - few 1–3(–4) internodes
 - terminal internode 40–100% of stem height
 - flowering typically (March-)May-June(-early July)
- Tried several times to cultivate it without any success beyond first year



- a few populations had intermediates which were treated as hybrids (*G. x davidiana*)



Using AFLPs on British plants only, Wingfield et al. (2003) found *G. amarella*, *G. uliginosa* and *G. anglica* to be closely related

- In mixed populations of *G. anglica* and *G. amarella*, individuals of the two species were genetically more similar to each other than they were to individuals of the same species from other populations.
- But there were “morphological and phenological differences between the taxa regardless of the fact that apparently there is no, or very little, genetic difference”




Fig. 2. a) PCO plot showing the relationship between the five species included in the study, and b) UPGMA dendrogram showing the same relationship. The bootstrap values placed above the forks of the dendrogram show the percentage of times the group to the right of that fork occurred – only percentages greater than 75% are shown. A = *G. anglica*, *G. amarella*, *G. uliginosa* and hybrid populations; B = *G. campestris*; C = *G. germanica*)

Conclusion

- *G. anglica* has been recently derived from *G. amarella* as early-flowering form
- Relatively few characters separate them
- Completely interfertile
- Relegated to subspecies: *G. amarella* subsp. *anglica**

- Consequently the hybrid *G. x davidiana* is no longer recognised
 - Such intermediate plants included in the variable subsp. *amarella*

- This also fits better with subsp. *septentrionalis* (retained as subspecies)




The case of *Gentianella uliginosa*

- Pugsley first noted in 1923 Tenby plants was similar to European *G. uliginosa*
- characterised by:
 - **annual** with 0–2(–3) internodes (mean 1.3)
 - terminal internode on c. 1.7 × the internode length,
 - terminal pedicel forming up to 70% of total height
 - calyx teeth very unequal in width and usually out-curved
- A rare plant of South Wales and North Devon dune slacks (reported in error for v.c. 102)



- Molecular studies by Winfield *et al.* (2003) showed that British plants ascribed to the European *G. uliginosa* were genetically part of *G. amarella* (as above)
- Our studies with Gerard Oostermeijer and Sabrina de Carvalho found that British plants were genetically different from Swedish *G. uliginosa* (unpublished).
- Furthermore, morphologically, European *G. uliginosa* differs from South Wales ‘*G. uliginosa*’ by having more internodes (mean 3.2), terminal internode 0.9 × the average internode length and a short terminal pedicel forming 20% of total plant height.




- Conclude South Wales ‘*G. uliginosa*’ has evolved from subsp. *amarella* as an annual ecotype adapted to dune slacks around the Severn Estuary – a new taxon = *Gentianella amarella* subsp. *occidentalis**

• * Rich, T. C. G., McVeigh, A. & Stace, C.A.(2018). New taxa and new combinations in the British flora. *Edinburgh Journal of Botany* doi: 10.1017/S0960428618000288

G. amarella subsp. *hibernica*


Pritchard (1959) described all Irish plants as subsp. *hibernica*

- relatively longer corollas 19-22 mm
- more internodes 7-11
- a very contracted terminal internode



Foiness

- Irish field population samples show almost complete overlap with subsp. *amarella*, though clustering suggests some differentiation
 - Hence overall subsp. *hibernica* not maintained



PCA: black UK, green Ireland

Post Handbook

Blackstonia may be annual and biennial (page 21); detailed studies needed!

Centaureum intermedium also in v.c. 60 West Lancs (page 60)

Gentiana verna from Ingleborough (page 130) has been strongly queried!

Gentiana lutea naturalised on cricket field Horwich, v.c. 59 South Lancs, 1954 (MANCH)



Andy and Tim would like to thank

- Stacey Baldwin
- Julia Carey
- Sabrina de Carvalho
- Sep Greimler
- Andy Jones
- Liz Lavery
- Tommy Lennartsson
- Lola Lledó
- Gerard Oostermeijer
- Colin Pope
- Noel Pritchard
- Ivor and Jane Rees
- Phil Smith
- Clive Stace
- Paul Stanley
- Seren Thomas
- Willow West
- Mike Wilcox
- Phil Wilson



and everyone else who has helped with information etc since 1993