

First addendum to the Leafhoppers and Planthoppers of Germany

(Hemiptera: Auchenorrhyncha)

Herbert Nickel¹

Abstract: Recent additions to the fauna of leafhoppers and planthoppers of Germany are communicated. *Cixidia lapponica* (Zett.), *Empoasca decedens* Paoli, *Kyboasca maligna* (Walsh), *Ribautiana cruciata* (Rib.), *Synophropsis lauri* (Horv.) and *Paralimnus lugens* (Horv.) are new for the country. New distribution data which indicate a recent range expansion are given for *Chloriona sicula* Mats., *Zygina lunaris* (M. & R.), *Orientus ishidae* (Mats.), as well as for the previously overlooked and little known *Paralimnus rotundiceps* (Leth.). *Paralimnus zachvatkini* Em. is proposed as a younger synonym of *P. lugens* (Horv.).

Zusammenfassung: Es werden neuere Ergänzungen zur Zikadenfauna Deutschlands mitgeteilt. *Cixidia lapponica* (Zett.), *Empoasca decedens* Paoli, *Kyboasca maligna* (Walsh), *Ribautiana cruciata* (Rib.), *Synophropsis lauri* (Horv.) und *Paralimnus lugens* (Horv.) sind neu für das Land. Neue Verbreitungsdaten, die für eine jüngere Arealausweitung sprechen, werden für *Chloriona sicula* Mats., *Zygina lunaris* (M. & R.), *Orientus ishidae* (Mats.) sowie für den bisher übersehnen und wenig bekannten *Paralimnus rotundiceps* (Leth.) aufgeführt. *Paralimnus zachvatkini* Em. wird als jüngeres Synonym von *P. lugens* (Horv.) vorgeschlagen.

Key words: Fulgoromorpha, Cicadomorpha, checklist, Germany, invasive species, range expansion, *Paralimnus lugens*, *Paralimnus zachvatkini*

1. Introduction

Since the publication of the checklist to the German Auchenorrhyncha (Nickel & Remane 2002) and the following compilation of ecological and distributional data and their analysis (Nickel 2003) a number of new species has been recorded, and the information on some little known species has considerably increased. This paper presents some results of recent field work, notably new species records for the country, more distributional data of formerly little known or expanding species, new host plant records and a few nomenclatural changes. It also includes a number of photographic records published only on the web. All records leg. et det. H. Nickel, if not otherwise stated.

2. New species records for Germany

Cixidia lapponica (Zetterstedt, 1840) (Fig. 1a)

Bayern: Spiegelau, above Jägerfleck, 810 m, VII.2007, 1 ♂, in a window trap. St. Oswald-Riedlhütte, above Guglöd, VII. and VIII.2007, altogether 2 ♂♂, 3 ♀♀; in Malaise traps at two different sites, at 800 and 890 m. All three localities are situated within open spruce forest with a high proportion of standing and lying dead trees.

¹ J.F. Blumenbach Institute of Zoology and Anthropology, University of Goettingen, Berliner Straße 28, 37073 Goettingen, Germany, E-Mail: hnickel@gwdg.de

A remarkable finding, since all over central Europe there is only a single record of this species published more than 30 years ago from northern Slovakia: "Nízké Tatry, Vajakovského dolina, c. 700 m, 16.VII.1974, 1 ♂, 1 ♀, on decaying logs of *Abies alba*" (Dlabola 1976). From Finland nymphs are reported to have been found in pine trunks, along with *Cixidia confinis* (Zett.), feeding on fungi hyphae (Linnavuori 1951). Neither *Abies* nor *Pinus* was present at the German sites. Therefore, fungi living in *Picea abies* are the most likely hosts. The species' range extends from eastern Siberia to Norway, Sweden, Finland and Middle Russia (Anufriev & Emelyanov 1988, Anufriev & Kirillova 1998, Nast 1972, Söderman 2007), apparently with isolated mountain populations in central Europe. Due to the large quantities of dead trees in higher situations of the Bavarian Forest and in other mountain regions in Germany, the species may be more widespread in central Europe.

***Empoasca decadens* Paoli, 1932 (Fig. 1a)**

Baden-Württemberg: Eimeldingen near Weil am Rhein, 260 m, 14.VIII.2005, 1 ♂, on *Salix alba* in a sand pit.

According to Nast (1972, 1987) this species has been recorded in Iraq, Jordan, Israel, Egypt, Libya, Georgia, Cyprus, Slovakia, Italy and France. Furthermore it has been found in Anatolia (Lodos & Kalkandelen 1983), Greece (Drosopoulos *et al.* 1986), Slovenia (Holzinger & Seljak 2001), Montenegro (Velimirovic 1980), Spain (Alvarado *et al.* 1994) and Madeira (Freitas & Aguin Pombo 2004). It is known as a pest of a variety of woody plants, e.g. cotton in Anatolia (Atakan 2009), almond in Spain (Jacas *et al.* 1997) raspberry in north Italy (Grassi & Ri 2006), and peach in Slovenia (Seljak 1997). Since the turn of the century it has been found on the Swiss side of the river Rhein around Basel (Mühlethaler 2001). At least its occurrence in the northern parts of Switzerland (where H. Günt hart has been collecting for decades) and in adjacent parts of Germany is probably the result of a recent range expansion. In fact it might be much more widespread, since this area is not well covered by Auchenorrhyncha collectors. In Italy it lives polyphagously on a variety of shrubs and trees, particularly favouring narrow-leaved *Salix*. Records in April and May indicate overwintering in the adult stage (Arzone *et al.* 2008).

***Kyboasca maligna* (Walsh, 1862) (Photo 1, Fig. 1a)**

Nordrhein-Westfalen: Köln, Auweiler, Landwirtschaftskammer Nordrhein-Westfalen, Zentrum für ökologischen Land- und Gartenbau, 45 m, 6.VIII.2010, 16 ♀♀ in an apple orchard, both on standard and half-standard trees.

This is a recent invader, originally found in North America from British Columbia and Ontario southward to California and Georgia (Metcalf 1968). The first European record is from eastern France in 1997: La Wantzenau, near Strasbourg, only 3 km from the German border (della Giustina & Remane (2001). After that it was found in Belgium in 1998, 2 single ♂♂ (Baugnée 2003), then in Luxembourg in 2003 until 2006, 10 localities (Niedringhaus *et al.* 2010a, 2010b) and Czechia, central Moravia, in 2006, a single ♂ (Malenovský & Lauterer 2010). Despite the close proximity of almost all these records, it took a relatively long time until the species was found in Germany. According to Hamilton (1985) it primarily feeds on cultivated *Malus*, and occasionally on *Crataegus*. All French records were from apple (R. Remane, pers. comm.), in Belgium it was collected on *Rosa canina* and *Carpinus betulus*, the Luxembourg records were from apple orchards, urban parks and woodland margins, and apple was not present in each locality. The Czech record was a single ♂ in an apple orchard.




Fig 1: Localities: **a:** ▲ = *Synophropsis lauri* (Horv.), ■ = *Paralimnus lugens* (Horv.), ● = *Kyboasca maligna* (Walsh), ▼ = *Cixidia lapponica* (Zett.), ♦ = *Empoasca decedens* Paoli. **b:** *Ribautiana cruciata* (Rib.). **c:** *Chloriona sicula* Mats., ◊ = records until 2000, ♦ = after 2001. **d:** *Zygina lunaris* (M. & R.), ◊ = records until 2000, ♦ = after 2001.

No feeding damages were observed in any of these localities. The European records were all gathered between mid June and mid August, and ♂♂ were only taken until mid July, indicating a single generation and overwintering as egg. But this conclusion must be considered as preliminary due to the small amount of data. In Canada it is reported to occur from mid June until late October (Hamilton 1985), suggesting a possible second generation.

***Ribautiana cruciata* (Ribaut, 1931) (Photo 2, Fig. 1b)**

Rheinland-Pfalz: Koblenz, Deutsches Eck, 70 m, 21.VIII.2005, 5 ♂♂, 6 ♀♀, shores of the Rhein. Bacharach, 80 m, 10.VI.2006, 18 ♂♂, 7 ♀♀, shores of the Rhein. Offstein, 14.VIII.2008, 4 ♂♂, 4 ♀♀, in a hedge. **Nordrhein-Westfalen:** Hopsten, Heiliges Meer, 45 m, 19.VI.2007, 2 ♂♂, 1 ♀, and 31.VII.2008, 2 ♀♀, on roadside elms. Uffeln, 50 m, 20.VI.2007, 2 ♀♀, on roadside elms. **Hessen:** Darmstadt, garden of the Julius-Kühn-Institut, 170 m, 16.VII.2007, 4 ♂♂, 5 ♀♀. Vogelsberg region, 2008, no further details (R. Remane, pers. comm.). **Bayern:** Schwebheim, 210 m, VIII.2007, 2 ♂♂, 1 ♀, along a woodland margin. **Niedersachsen:** Gehrde, 30 m, 31.VII.2008, 6 ♀♀ in a hedge.

A mediterranean and western European elm-feeder, which was so far known to occur northward to England, Belgium, Switzerland and Slovenia (Nast 1987, Le Quesne & Payne 1981, Löcker 2003), and which has recently expanded its range eastward. It is unlikely that it was overlooked in Germany, since W. Wagner, R. Remane, R. Niedringhaus and the author frequently sampled elms all over the country. In Germany it was found on *Ulmus minor* and *U. × hollandica*. The new records indicate two generations per year and overwintering as egg.

***Synophropsis lauri* (Horváth, 1897) (Photo 4, Fig. 1a)**

Hamburg: Botanical Garden, 20 m, 13.IX.2008, 1 individual photographed on a beerglass and published on the web (Photo 4, downloaded from <http://insektenfotos.de>, Renate Ridley, Hamburg). In close vicinity there were two potted bay trees (*Laurus nobilis*). **Baden-Württemberg:** Öhringen, 240 m, 8.IX.2009, 1 ♂, at light in a living room, a large stand of *Hedera helix* nearby, but no *Laurus nobilis* (P. Dynort, pers. comm.).

This species is widespread in most Mediterranean countries (e.g. Nast 1987), feeding on a variety of evergreen trees and shrubs. After being recorded from Basel, Switzerland, in 2000 (Mühlethaler 2001) and England since 2007 (Bantock & Botting 2010) it is now likely to be more widespread in the western half of Germany, although populations may be largely synanthropic in public gardens and parks. *Laurus nobilis* and *Hedera helix* appear to be the most important host plants in central parts of Europe, where broad-leaved evergreen plants are relatively uncommon.

***Paralimnus lugens* (Horváth, 1897) (Photo 5, Fig. 1a)**

Deltocephalus phragmitis var. *lugens* Horváth, 1897

Paralimnus zachvatkini Emelyanov, 1964 syn. nov.

Niedersachsen: Adelebsen, Schwülme basin, 185 m, 13.VII.1995, 1 ♂, and 9.VII.2009, 9 ♂♂, 2 ♀♀, 8 nymphs, plus numerous individuals seen, all in a rather dry and dense stand of reed (*Phragmites australis*). The closely related *P. phragmitis* (Boh.) (Photo 6) was not found in this locality. Immediately after collecting, the first-mentioned specimen was misidentified and mislabelled as *P. phragmitis* (Boh.) by the author. Under this name it was offered to Biedermann & Niedringhaus (2004, 2009), who published a drawing of the dorsal view.



Photo 1: *Kyboasca maligna* (Walsh), ♀, Gostingen (Luxemburg state), 4.VIII.2010 (G. Kunz)



Photo 2: *Ribautiana cruciata* (Rib.), ♂, Darmstadt, 31.VIII. 2008 (G. Kunz)



Photo 3: *Synophropsis lauri* (Horv.), ♂, on a beer glass, Hamburg, Botanical Garden, 13.IX.2008 (R. Ridley)



Photo 4: *Zygina lunaris* (M. & R.), ♀, Lorch am Rhein, 6.VIII.2010 (G. Kunz)



Photo 5: *Paralimnus lugens* (Horv.), ♂, Adelebsen, 10.VII.2009 (G. Kunz)



Photo 6: *Paralimnus phragmitis* (Boh.), ♂, Rabenhof, Styria (Austria), 19.VIII.2007 (G. Kunz)



Photo 7: *Paralimnus rotundiceps* (Leth.), ♂, Gimte, 10.VII.2009 (G. Kunz)



Photo 8: *Orientus ishidae* (Mats.), ♂, Graz (Austria), 19.IX.2007 (G. Kunz)

Despite revisions of all available material by the author and colleagues (R. Remane, pers. comm.; R. Niedringhaus, pers. comm., S. Walter, pers. comm.) and targeted search in many similar habitats, this locality remained the first and only one confirmed from Germany so far. This is apparently a rare species in Europe. It has long been treated only as a variety of *P. phragmitis* (Boh.), until Koczor (2005), who found significant differences in coloration and genital morphology, raised it to species level. He listed all distributional records so far published, including Hungary, Transcaucasia (Horvàth's type series, comprising only 2 single ♀♀), Yemen (Zachvatkin 1935) and Germany (Hueber 1904), the two latter clearly in need of revision. Blöte (1927) in his Dutch leafhopper fauna and Haupt (1935) in his Auchenorrhyncha fauna of central Europe only mentioned the existence of var. *lugens* Horv., but did not give specific records. Later, *Paralimnus zachvatkini* Em., described from the delta of the river Volga (southern Russia, Astrakhan Province), was published from Hungary (Orosz 1999), Bulgaria (Emelyanov *et al.* 2002) and Greece (Drosopoulos *et al.* 1986). More recently it was found in south Finland (Söderman 2004, 2007), Lithuania (Söderman & Dapkus 2009) and Tuscany, Italy (Mazzoni 2005). It is also known from the southwest Russian provinces of Voronezh and Belgorod (A.F. Emelyanov, pers. comm.) and from eastern Zaysan, Kazakhstan (Mityaev 2002). Its synonymy with *P. lugens* (Horv.) was kindly confirmed by A.F. Emelyanov (St. Petersburg).

In Hungary adults were found from mid June until mid October (Koczor 2005), which indicates overwintering as egg and one, perhaps two generations per year.

3. Records of rare and little known species

Chloriona sicula Matsumura, 1910 (Fig. 1c)

Bayern: Volkach, Halbmilesee, 230 m, 14.VIII.2010, 10 ♂♂ (G. Kunz, pers. comm.).
Niedersachsen: Negenborn, Helstorfer Moor, 24.V.2008, 1 ♂; Resse, Schwarzes Moor, 45 m, 20.V. 2009, 1 ♂ (the two latter already published by Nickel & Gärtner 2009). Hannoversch Münden, Kiesgrube Ballertasche near Gimte, 110 m, 10.VII. 2009, 2 ♂♂. Friedland, 180 m, 19.VI. 2008, 7 ♂♂, 12.VIII.2010, 8 ♂♂. **Hessen:** Langen-Mörfelden, Langener Waldsee, 100 m, 10.VII.2008, 2 ♂♂. Darmstadt, Ruthsenbach-Grund, 190 m, 14.V.2008, 2 ♂♂. **Brandenburg:** Ludwigsfelde, Gröbener See, 35 m, 8.VIII.2010, 6 ♂♂. Recorded on *Phragmites australis* in a variety of habitats including sand pits, pond shores, peat bogs and saltmarshes.

This species is widespread from the Mediterranean region eastward to Kazakhstan (Nast 1972, 1987; Holzinger *et al.* 2003). In Germany it was until recently known only from 5 records along the southern Rhine between Waldshut and Speyer, the earliest one dating from 1971 (Remane & Fröhlich 1994, Nickel 2003). Since then it has expanded its range quickly. W. Wagner, R. Remane, H. Strübing, W. Fröhlich, R. Niedringhaus and the author of this paper have been collecting extensively on reed in central and northern parts of Germany, and even in at least two of the above-mentioned localities. So it is unlikely that this species was overlooked. Just recently it was also published from southern Moravia, Czechia (Malenovský & Lauterer 2010). Like its congeners, it overwinters as nymph. The presence of ♂♂ in August clearly indicates two generations a year, at least in favorable years.

Zygina lunaris (Mulsant et Rey, 1855) (Photo 4, Fig. 1d)

Rheinland-Pfalz: Kobern, Mosel, 70 m, 14.VIII.2004, 1 ♂, 4 ♀♀. Koblenz, Rhein, 70 m, 21.8.2005, 5 ♂♂, 9 ♀♀, on *Salix alba* and *S. babylonica*. Lorch, Rhein, 80 m, 01.IX.2008, 1 ♀, 1 nymph, on *S. fragilis*. Offstein (near Worms), 140 m, 01.IX.2008, 1 ♀, on *S. purpurea* in a garden. **Baden-Württemberg:** Höhfeld, Klosterhöhe, 340 m, 11.VI.2008, 1 ♂, on *S. purpurea* at a field margin. **Bayern:** Großwelzheim, Main, 105 m, 14.VIII.2009, 1 ♀, on *S. fragilis*. **Nordrhein-Westfalen:** Köln, Auweiler, 45 m, 5.VIII.2010, 2 ♂♂, 2 ♀♀ on ornamental *S. purpurea* in a garden.

This species is a recent invader from south-western Europe. Nast (1987) mentions it from Portugal, Spain and France. Mühlethaler (2001) recorded it in Basel, Switzerland, in 2000, and Niedringhaus *et al.* (2010a, 2010b) just published 5 localities in the south of Luxembourg dating from 2006. In Germany it was until recently only known from a few scattered localities along the southern upper Rhein near Kehl and Weil am Rhein (Lauterer & Malenovský 1995). The new records extend the known range into an area, where Kirschbaum (1868), Wagner (1939), Remane (1987) as well as students of the lower Rhein fauna (Mölleken & Topp 1997, Frommer 1996) would probably not have overlooked it.

Orientus ishidae (Matsumura, 1902) (Photo 8, Fig. 2a)

Baden-Württemberg: Wyhlen, Rheinufer, 260 m, 14.VIII.2005, 1 ♂, on *Salix babylonica*. Istein, Totengrien, 235 m, and villages of Mappach, 310 m, and Efringen-Kirchen, 250 m, 30.VII.2009, small numbers, respectively. Schwetzingen, 110 m, 24.VII.2009, 5 individuals, at a woodland margin. Öhringen, 240 m, first recorded 12.VII.2005, since then, until 2010, common in gardens, orchards and at light (P. Dynort, pers. comm.). Heilbronn, c. 160 m, 12.VII.2009, 1 nymph, on *Prunus laurocerasus*, reared until emergence; Jagsthausen, c. 200 m, 20.VII.2010. Horkheim, 165 m, 21.VII.2010; the two latter on apple in an orchard (the last 3 records K. Schrameyer, pers. comm.). Winnenden, 290 m, 21.VII.2009, 9 individuals, at light in a kitchen (H. Schlüter, pers. comm.). Rottenburg am Neckar, 350 m, 7.IX.2009, on *Hedera helix* in a garden, and 25.VII.2010, on *Urtica dioica* along a stream (the last two M. Albers, pers. comm.). **Rheinland-Pfalz:** Ludwigshafen, 95 m, 8.VIII. 2009, 1 ♀ (S. Schuch, pers. comm.). **Bayern:** Wechterswinkel, 265 m, 3.VIII.2008, 1 individual, on a house wall (E. Dietz, pers. comm.). **Hessen:** Darmstadt, garden of the Julius Kühn Institute, 170 m, 16.VII.2007, 1 nymph. Frankfurt, Langener Waldsee, 100 m, 10.VII.2008, 1 ♀, on *Salix alba*. Gelnhausen, 130 m, 30.VII.2010, 1 individual, on *Corylus avellana* (V. Wagner, pers. comm.). **Nordrhein-Westfalen:** Mülheim an der Ruhr, Heißen, 50 m, 17.IX.2006, 2 ♂♂, 1 ♀ (K. Winzer, pers. comm.), 22.X.2006, 1 individual (K. Winzer leg., R. Dietze det.). Dortmund, 95 m, 16.VII.2007, 1 nymph (S. Brüggenthies, pers. comm.). Neuss, Grefrath, 40 m, 27.VII.2008, 1 individual, in a garden; Köln, Stommeln, 55 m, 22.VIII. 2007, 1 individual; Köln, Feste Zons, 50 m, 28.VII.2008, 1 nymph (the last three M. Becker, pers. comm.). Köln, Wahner Heide, 6.VIII.2009, 1 individual, on a roadside embankment; Köln, Nippes, 10.VIII. 2009, 3 individuals, on stems of *Fagus sylvatica*; Köln, Rhiel, 23.VII.2010, 4 individuals, at light; Bergheim-Erf, 8.VIII.2009, 1 individual, on a house wall, and again 1 individual, 15.-29.VII. 2010, in a pitfall trap (the last 4 records: T. Hörren, pers. comm.). Köln, Auweiler, Landwirtschaftskammer Nordrhein-Westfalen, Zentrum für ökologischen Land- und Gartenbau, 45 m, 6.VIII.2010, at least 50 individuals, in apple orchards and hedges of *Carpinus betulus* and *Acer campestre*. Bochum, 22.VIII.2010, 4 individuals at light (A. Steiner, pers. comm.).




Fig 2: Localities: **a:** *Orientus ishidae* (Mats.), not all Berlin localities shown. **b:** *Paralimnus rotundiceps* (Leth.), ♦ = records until 2000, ♦ = after 2001.

Sachsen-Anhalt: Halle an der Saale, Heide-Süd, c. 100 m, since 27.VII. 2007, numerous nymphs and adults on hedges of *Carpinus betulus*; also recorded in other parts of the city and on other trees and shrubs (W. Witsack, in prep.). Halle an der Saale, inner city, 100 m, 9.VIII.2010, at least 6 individuals, found dead under a lamp inside a flat (J. Quitsch, pers. comm.). **Berlin:** first recorded in 2004 on 11 urban brownfield sites (Strauss 2007, N. Lange, pers. comm.). Johannisthal, 35 m, 17.IX.2009, at least 50 individuals, on *Acer campestre*, *Betula pendula*, *Salix x rubens*, *Populus nigra italicica*, on an abandoned airstrip. Grunewald, Im Jagen, 16.IX.2009, 1 individual. Biesenhorster Sand, 18.IX.2009, 2 ♀♀. Baumschulenweg, 25.IX.2008 and 13.VII.2010 (C. Brückner, pers. comm.).

In 1998 this new species to the European fauna was found near Milano, north Italy. Guglielmino (2005) was the first to raise questions about its specific identity, eventually leading to the description of *Orientus amurensis*, occurring in the Far East of Russia and eastern China. Her conclusion was further, that *O. ishidae* was originally restricted to Japan (and perhaps the Philippines and Taiwan), but introduced to North America and Europe. After its discovery in Italy, R. Mühlethaler found it to be common in urban areas of Basel, Switzerland, since 2000 (Günthart & Mühlethaler 2002). In 2002 it was recorded for the first time in Slovenia (Seljak 2004), in 2004 in Brno, Moravia, in 2006 in Prague, Bohemia (Malenovský & Lauterer 2010), in 2007 in Graz, Austria (Holzinger 2009).

The earliest German records date from 2002: Weil am Rhein (R. Mühlethaler) and Dresden (R. Remane), see Nickel & Remane (2003). Within only 8 years this invader has become widespread and locally common in Germany, with considerable population numbers at least in the Upper Rhine plain, urban areas of Berlin and the Lower Rhine and Ruhr region. It lives polyphagously on shrubs and low trees, notably *Salix*, *Carpinus*, *Betula*, *Corylus*, *Acer*, *Malus* and others, and it is also attracted to light, frequently flying into

houses. The majority of records is from urban sites, but the species was also found along rivers and woodland margins as well as in sand pits. Most findings are from July until September, the earliest from July 10th, the latest from October 22nd. This indicates a single generation per year and overwintering as egg.

***Paralimnus rotundiceps* (Lethierry, 1885) (Photo 7, Fig. 2b)**

Niedersachsen: Hannoversch Münden, Ballertasche near Gimte, 110 m, 4.VII.2006, 24.VI.2007, and again 10.VII.2009, a large population in a sand pit with sparsely growing stands of *Phragmites australis*. **Sachsen-Anhalt:** Halle an der Saale, Heide-Süd, c. 100 m, small numbers recorded from 2007 until 2010 (W. Witsack, pers. comm.). **Baden-Württemberg:** Weil am Rhein, 250 m, 30.VII.2009, 6 ♂♂, 2 ♀♀, in a sand pit. **Bayern:** Füssen, Lech floodplains, 790 m, VII.2003, 1 ♂, in a Malaise trap.

Until recently this rare species was known to occur in Germany only in two isolated localities in coal mining areas of Sachsen-Anhalt (Mücheln near Merseburg, see Funke & Witsack 2002) and in a riverine gravel bed in the Alps (Friedergries near Garmisch-Partenkirchen, see Nickel 2003). However, it appears to be more widespread though rare. It is confined to very sparse pioneer stands of reed, usually on non-flooded sites such as sand and gravel pits and banks of unregulated rivers. It is further reported from Norway, Sweden, Finland, Belgium, France, Switzerland, Austria, Italy, Slovenia, Hungary and Romania (Nast 1972, 1987, Holzinger & Seljak 2001).

4. Nomenclatural changes

Two generic changes have been proposed during recent years. According to Holzinger & Kunz (2006) *Mocuellus* Ribaut, 1946 is a younger synonym of *Henschia* Lethierry, 1892; therefore the valid name of the former *Mocuellus collinus* (Boh.) must now be *Henschia collina* (Boh.). Second, most species formerly placed into the genus *Recilia* Edwards, 1922, including *R. schmidgeni* (W.Wg.) and *R. horvathi* (Then) have been moved to *Maiestas* Distant, 1916 (Webb & Viraktamath 2009).

5. Discussion

Until recently the number of Auchenorrhyncha known from Germany included 620 species (Nickel & Remane 2002, Nickel 2003). The following checklist for the German federal states published by Nickel & Remane (2003) adopted the new synonymy of *Kelisia nervosa* Vilb. with *K. confusa* Lnv. proposed by Holzinger *et al.* (2003) and included the newly recorded *Orientus ishidae* (Mats.). Since then *Ribautiana cruciata* (Rib.) was the only addition (Nickel & Niedringhaus 2009). Together with the new species mentioned in the present paper, *Cixidia lapponica* (Zett.), *Empoasca decedens* Paoli, *Kyboasca maligna* (Walsh), *Synophropsis lauri* (Horv.) and *Paralimnus lugens* (Horv.), the German list now includes 626 species.

For three of the species mentioned here, notably *Cixidia lapponica* (Zett.), *Paralimnus lugens* (Horv.) and *P. rotundiceps* (Leth.), it is likely that they had been overlooked in Germany until now, due to their peculiar life habits and rather specific habitat requirements. But for the remaining species, *Chloriona sicula* Mats., *Ribautiana cruciata* (Rib.), *Synophropsis lauri* (Horv.), *Kyboasca maligna* (Walsh), *Empoasca decedens* Paoli, *Zygina lunaris* (M. & R.), and *Orientus ishidae* (Mats.), a more or less northeastward range expansion must be

assumed, which took place not before the last two or three decades, and which happened quite rapidly. In many cases, the earliest German records are from the southern upper Rhine plain, or less commonly, along the middle Rhine. A similar pattern of north-eastward expansion has also been observed in other species of Auchenorrhyncha such as *Muellerianella fairmairei* (Perr.), *Haematoloma dorsatum* (Ahr.), *Stictocephala bisonia* Kopp & Yonke, *Viridicerus ustulatus* (M. & R.), *Graphocephala fennahi* Young, *Liguropia juniperi* (Leth.), *Lindbergina aurovittata* (Dgl.), *Ribautiana debilis* (Dgl.), *Eupteryx decemnotata* (R.) and *Zyginidia scutellaris* (H.-S.) (Nickel 2003). Surprisingly this list includes invaders native to southern Europe but likewise species originating from other continents and first introduced to more southern parts of Europe. On the other hand, invaders originating from south-eastern and eastern Europe are rare. One example that is frequently grouped among these is the planthopper *Eurybregma nigrolineata* Scott (Remane & Fröhlich 1994), which has, however, been described from England in the 19th century.

Acknowledgements

I thank Gernot Kunz (Graz) for the photographs, Sasha Emelyanov (St. Petersburg) for the confirmation of the synonymy of *Paralimnus zachvatkini* Em., and Peter Dynort (Öhringen), Martin Albers (Rottenburg am Neckar), Claudia Brückner (Berlin), Thomas Hörren (Köln), Nora Lange (Berlin), Renate Ridley (Hamburg), Klaus Schrammeyer (Heilbronn), Sebastian Schuch (Göttingen), Axel Steiner (Breckerfeld), and Werner Witsack (Halle an der Saale) for the contribution of records. Michael Becker (Köln), Sigrun Brüggenthies (Dortmund), Eugen Dietz (Bad Neustadt), Julian Quitsch (Halle an der Saale), Horst Schlüter (Winnenden) and Volkmar Wagner (Gelnhausen) communicated details about their photographs published on the web. *Cixidia lapponica* (Zett.) was recorded within the project BIOKLIM, funded by Bayerisches Staatsministerium für Umwelt, Gesundheit und Verbraucherschutz.

Most of all I thank Reinhard Remane († April 27 2009) for almost 20 years of inspiring discussion and guidance, and for his encouragement to mistrust anything that is published.

6. References

- Alvarado, M., Berlanga, M., González Zamora, J.E., Serrano, A., de la Rosa, A., Villalgordo, E. (1994): Contribución al conocimiento del mosquito verde (*Empoasca decedens* Paoli) en melocotonero en el Valle del Guadalquivir. – Boletín de sanidad vegetal. – Plagas 20(3): 771-783.
- Anufriev, G.A., Emeljanov, A.F. (1988): Suborder Cicadinea (Auchenorrhyncha) - Cicads. – In: Ler, P.A. (ed): Keys to the identification of insects of the Soviet Far East. Vol. 2: Homoptera and Heteroptera. pp. 12-495. Nauka, Leningrad.
- Anufriev, G.A., Kirillova, B.I. (1998): The Cicadina of the Chuvash Republic. – Cheboksarij. 176 pp. [In Russian].
- Arzone, A., Alma, A., Mazzoglio, P.J. (2008): Collections made by prof. Carlo Vidano. – Memorie della Società Entomologica Italiana 86, 480 pp.
- Atakan, E. (2009): Damage assessment of the leafhopper complex [*Asymmetrasca decedens* (Paoli) and *Empoasca decipiens* Paoli] (Homoptera: Cicadellidae) in cotton. – Journal of Pest Science 82(3): 227-234.
- Bantock, T., Botting, J. (2010): Checklist of UK Auchenorrhyncha (Homoptera). – <http://www.britishbugs.org.uk> (accessed 20.III.2010)

- Baugnee, J.-Y. (2003): Sur la présence en Belgique du Cicadellidae *Kyboasca maligna* (Walsh, 1862) et du Psyllidae *Cacopsylla fulguralis* (Kuwayama, 1907) (Hemiptera Homoptera). – Bulletin de la Société Royale Belge d'Entomologie 139: 72-73.
- Biedermann, R., Niedringhaus, R. (2004): [English translation 2009] Die Zikaden Deutschlands. – Wissenschaftlich Akademischer Buchvertrieb Fründ, Scheeßel. 409 pp.
- Blöte, H.C. (1927): Fauna van Nederland. Afl. II. Homoptera (QXXI) A. Jassidae, Membracidae. – Leiden.
- Dlabola, J. (1976): Faunistic records from Czechoslovakia: Homoptera. Auchenorrhyncha. – Acta entomologica Bohemoslovaca 3: 348.
- Drosopoulos, S., Asche, M., Hoch, H. (1986): A preliminary list and some notes on the Cicadomorpha (Homoptera-Auchenorrhyncha) collected in Greece. – Proceedings of the 2nd International Congress Concerning the Rhynchota fauna of Balkan and adjacent Regions 1986: 8-13.
- Freitas, N., Aguin-Pombo, D. (2004): Is the leafhopper *Asymmetrasca decadens* (Paoli, 1932) invading Madeira Island? – Annales de la Société Entomologique de France 40 (1): 103-104.
- Frommer, W. (1996) Untersuchungen zur Zikadenfauna (Hemiptera: Homoptera, Auchenorrhyncha) ausgewählter Standorte in Köln. – Decheniana (Suppl.) 35: 163-174.
- Funke, T., Witsack, W. (2002): Zur Arthropodenfauna von Tagebaufolgelandschaften Sachsen-Anhalts. 2. Zikaden (Auchenorrhyncha, Hemiptera, Insecta) von Offenlandhabitaten. – Hercynia N.F. 35, 91-122.
- Giustina, W. della, Remane, R. (2001): Compléments à la faune de France des Auchenorrhyncha: espèces et données additionnelles; modifications à l'ouvrages de Nast (1987) (Homoptera). – Bull. Soc. Ent. Fr. 106 (3): 283-302.
- Grassi, A., Ri, M.D. (2006): *Asymmetrasca (Empoasca) decadens* Paoli (Hom., Typhlocybinae): a new pest of cultivated red raspberry in Trentino, Italy. – IOBC/WPRS Bulletin 29(9): 41-48.
- Günthart, H., Mühlthaler, R. (2002): Provisorische Checklist der Zikaden der Schweiz (Insecta: Hemiptera, Auchenorrhyncha). – Denisia 4: 329-338.
- Guglielmino, A. (2005): Observations on the genus *Orientus* (Rhynchota Cicadomorpha Cicadellidae) and description of a new species: *O. amurensis* n. sp. from Russia (Amur Region and Maritime Territory) and China (Liaoning Province). – Marburger entomologische Publikationen 3(3): 99-110.
- Hamilton, K.G.A. (1985): Leafhoppers of ornamental and fruit trees in Canada. – Agriculture Canada Publication 1779E/F: 1-71.
- Haupt, H. (1935): Unterordnung: Gleichflügler, Homoptera. – In: Brohmer, P., Ehrmann, P., Ulmer, G. (eds.): Die Tierwelt Mitteleuropas IV (X): 115-262.
- Holzinger, W. E. (2009): Rote Listen gefährdeter Zikaden (Auchenorrhyncha) Österreichs. – In: Wallner R., Zulka K.P. (eds.): Rote Listen gefährdeter Tiere Österreichs 14(3). Böhlau, Wien, pp. 41-317.
- Holzinger, W. E., Kunz, G. (2006): New records of leafhoppers and planthoppers from Austria (Hemiptera: Auchenorrhyncha). – Acta entomologica Slovenica 14(2): 163-174.
- Holzinger, W.E., Seljak, G. (2001): New records of planthoppers and leafhoppers from Slovenia, with a checklist of hitherto recorded species (Hemiptera: Auchenorrhyncha). – Acta Entomologica Slovenica 9(1): 39-66.
- Hüeber, T. (1904): Catalogus insectorum faunae germanicae: Hemiptera homoptera. Systematisches Verzeichnis der Zikaden Deutschlands (und der nächst angrenzenden Landesteile). – Jahreshefte des Vereins für vaterländische Naturkunde in Württemberg 60: 253-277.
- Jacas, J.A., Hermoso Mendoza, A., Cambra, M., Balduque, R. (1997): *Asymmetrasca decadens*: a new pest of almond in Spain. – EPPO Bulletin 27(4): 523-524.
- Kirschbaum, C.L. (1868): Die Cicadinen der Gegend von Wiesbaden und Frankfurt a.M. nebst einer Anzahl neuer oder schwer zu unterscheidender Arten aus anderen Gegenden Europas. – Jahrbücher des nassauischen Vereins für Naturkunde 21/22: 1-202.

- Koczor, S. (2005): Revision of the taxonomic status of *Paralimnus phragmitis* var. *lugens* (Horváth, 1897) (Auchenorrhyncha: Cicadellidae). – *Folia Entomologica Hungarica* 66: 51-56.
- Lauterer, P., Malenovský, I. (1995): Eine neue Art der Kleinzikaden (Homoptera Cicadellidae Typhlocybinae) auf dem Gebiet der Bundesrepublik Deutschland. – *Marburger entomologische Publikationen* 2(9): 76-77.
- Le Quesne, W., Payne, K.R. (1981): Cicadellidae (Typhlocybinae) with a checklist of the British Auchenorrhyncha (Hemiptera, Homoptera). – *Handbooks for the Identification of British Insects* II (2 c), 95 pp.
- Linnauvori, R. (1951): Hemipterological observations. – *Annales Entomologici Fennici* 17: 51-65.
- Löcker, H. (2003): Arborikole Zikaden-Gilden in Slowenien (Hemiptera, Auchenorrhyncha). – *Beiträge zur Zikadenkunde* 6: 7-38.
- Lodos, N., Kalkandelen, A. (1983): Preliminary list of Auchenorrhyncha with notes on distribution and importance of species in Turkey, XII. Family Cicadellidae: Typhlocybinae: Empoascini. *Türk Bit. Kor. Derg.* 7(3): 153 -165.
- Mazzoni, V. (2005): Contribution to the knowledge of the Auchenorrhyncha (Hemiptera Fulgoromorpha and Cicadomorpha) of Tuscany (Italy). – *REDIA* 88: 85-102.
- Metcalf, Z.P. (1968): General Catalogue of the Homoptera. Cicadelloidea, Part 17, Cicadellidae. United States Department of Agriculture, Agricultural Research Service, Washington, D.C. 6(17), viii + 1513 pp.
- Mityaev, I. (2002): Fauna, ecology and zoogeography of leafhoppers (Homoptera, Cicadinea) of Kazakhstan. – *Tethys Entomological Research* 5: 1-168. [In Russian]
- Mölleken, H., Topp, W. (1997): Die Insektenfauna auf Silberweiden (*Salix alba* L.): Einfluß des Geschlechts und der Pflegemaßnahmen. – *Zeitschrift für Ökologie und Naturschutz* 6: 193-206.
- Mühlethaler, R. (2001): Untersuchungen zur Zikadenfauna der Lebensraumtypen von Basel. – *Beiträge zur Zikadenkunde* 4: 11-32.
- Nast, J. (1972): Palaearctic Auchenorrhyncha (Homoptera). An annotated check list. – Polish Scientific Publ. Warszawa. 550 pp.
- Nast, J. (1987): The Auchenorrhyncha (Homoptera) of Europe. – *Annales zoologici Warszawa* 40: 535-662.
- Nickel, H. (2003): The leafhoppers and planthoppers of Germany (Hemiptera, Auchenorrhyncha): patterns and strategies in a highly diverse group of phytophagous insects. – Pensoft, Sofia und Moskau. 460 pp.
- Nickel, H., Gärtner, E. (2009): Tyrphobionte und tyrphophile Zikaden (Homoptera, Auchenorrhyncha) in der Hannoverschen Moorgeest – Biotopspezifische Insekten als Zeigerarten für den Zustand von Hochmooren. – *Telma* 39: 45-74.
- Nickel, H., Niedringhaus, R. (2009): Die Zikaden des Naturschutzgebietes „Heiliges Meer“, mit Anmerkungen zur Fauna Westfalens und Nordwest-Deutschlands (Homoptera, Fulgoromorpha et Cicadomorpha). – Abhandlungen aus dem Westfälischen Museum für Naturkunde 71 (4): 213-271.
- Nickel, H., Remane, R. (2002): Artenliste der Zikaden Deutschlands, mit Angaben zu Nährpflanzen, Nahrungsbreite, Lebenszyklen, Areal und Gefährdung (Homoptera, Fulgoromorpha et Cicadomorpha). – *Beiträge zur Zikadenkunde* 5: 27-64.
- Nickel, H., Remane, R. (2003): Verzeichnis der Zikaden (Auchenorrhyncha) der Bundesländer Deutschlands. – *Entomologische Nachrichten und Berichte, Suppl.* 8: 130-154.
- Niedringhaus, R., Biedermann, R., Nickel, H. (2010a): Verbreitungsatlas der Zikaden des Großherzogtums Luxemburgs. Textband. – Ferrantia - Travaux scientifiques du Musée naturelle Luxembourg 60, 105 pp.
- Niedringhaus, R., Biedermann, R., Nickel, H. (2010b): Verbreitungsatlas der Zikaden des Großherzogtums Luxemburgs. Atlasband. – Ferrantia - Travaux scientifiques du Musée naturelle Luxembourg 61, 395 pp.

- Orosz, A. (1999): New Auchenorrhyncha species from Hungary (Homoptera). – *Folia entomologica hungarica* 60: 153-164.
- Remane, R. (1987): Zum Artenbestand der Zikaden (Homoptera: Auchenorrhyncha) auf dem Mainzer Sand. – *Mainzer naturwissenschaftliches Archiv* 25: 273-349.
- Remane, R., Fröhlich, W. (1994): Beiträge zur Chorologie einiger Zikaden-Arten (Homoptera Auchenorrhyncha) in der Westpaläarktis. – *Marburger entomologische Publikationen* 2(8): 131-188.
- Seljak G., (1997): *Empoasca decedens* Paoli - a new pest on peach trees in Slovenia. – *Zbornik predavanj in referatov s 3. Slovenskega posvetovanja o varstvu rastlin*, Portoroz, 4.-5. Marec 1997, pp. 339-344. [in Slovenian]
- Seljak G., (2004): Contribution to the knowledge of planthoppers and leafhoppers of Slovenia (Hemiptera: Auchenorrhyncha). – *Acta Entomologica Slovenica* 12(2): 189-216.
- Söderman, G. (2004): Auchenorrhyncha captured at light in Finland (Hemiptera, Fulgoromorpha et Cicadomorpha). – *Beiträge zur Zikadenkunde* 7: 17-28.
- Söderman, G. (2007): Taxonomy, distribution, biology and conservation of Finnish Auchenorrhyncha. – *The Finnish Environment* 7, 101 pp.
- Söderman, G., Dapkus, D. (2009): New records of Hemiptera from the Čepkeliai strict Nature Reserve (Southern Lithuania) in 2007. – *Naujos ir retos Lietuvos vabzdziu rusys* 21: 12-14.
- Strauss, G. (2007): Insects in urban brownfields. Analyses of species occurrences, community composition, and trait frequencies along a successional gradient. – Doctoral thesis, University of Oldenburg. 145 pp.
- Velimirovic, V. (1980): Cikada – *Empoasca decedens* Paoli stetocinana Citrus kulturama u predjelu crnogorskog Primorja. – *Zastita bilja* 31(3): 273-276.
- Wagner, W. (1939): Die Zikaden des Mainzer Beckens. – *Jahrbücher des nassauischen Vereins für Naturkunde* 86: 77-212.
- Webb, M.D., Viraktamath, C.A. (2009): Annotated check-list, generic key and new species of Old World Deltocephalini leafhoppers with nomenclatorial changes in the Deltocephalus group and other Deltocephalinae (Hemiptera, Auchenorrhyncha, Cicadellidae). – *Zootaxa* 2163: 1–64.