

Plant Propagation Protocol for *Alnus rubra*

ESRM 412 – Native Plant Production

Protocol URL: <https://courses.washington.edu/esrm412/protocols/ALRU2.pdf>




© Smithsonian Institution
10



© Susan McDougal
10

TAXONOMY	
Plant Family	
Scientific Name	Betulaceae
Common Name	Birch family
Species Scientific Name	
Scientific Name	<i>Alnus rubra</i> Bong.
Varieties	
Sub-species	
Cultivar	
Common Synonym(s)	<i>Alnus oregona</i> Nutt. ¹⁵
Common Name(s)	Red alder, Oregon alder
Species Code (as per USDA Plants database)	ALRU2
GENERAL INFORMATION	



Coastal North America and a few isolated populations in Idaho and Montana ¹⁰

Ecological distribution
 Will grow along rivers or roads, areas that regularly experience stressful conditions or that are recovering from a disturbance ^{6, 12}.
 Upland stands of *Alnus rubra* in coniferous forests with *Rubus spectabilis* Pursh or *Sambucus racemosa* L. in the understory
 In riparian environments where conifers also dominate
Alnus rubra in mixed stands with codominant species *Acer macrophyllum* Pursh. and *Populus trichocarpa* Brayshaw.
 A swamp environment where *Thuja plicata* D.Don is dominant ⁵.
 Found often in low nutrient soils because its ability to fix nitrogen ⁷.

Climate and elevation range
 Sea level to approximately 700 meters in elevation ^{2, 5}
 Within 160 kilometers from the coast
 Cold, wet winters and hot, dry summers, mild coastal climates ⁵

Local habitat and abundance
 Can be found in a pure stand of all *Alnus rubra*, co-dominating with other deciduous woody shrubs or trees such as *Acer macrophyllum* and *Populus trichocarpa*, co-dominating with *Thuja plicata*, and have an understory of *Rubus spectabilis* and *Sambucus racemosa* ⁵.

Plant strategy type / successional stage
 Early seral species, will easily establish themselves after disturbance such as forest fires, landslides, or logging ^{5, 9}.
 Allow for shade-tolerant species to grow and establish themselves by using *Alnus rubra* canopy ⁹.
Alnus rubra are shade intolerant, and will therefore self-thin to maintain a uniform height ⁵.

Plant characteristics
 Up to 20-30 meters tall ^{2, 5}.
 Hardy, relatively short-lived tree species ^{5, 12}.
 Dark grey to white speckled bark
 Prominently veined and serrated yellow to bright green leaves
 Maximum diameter at breast height 80 cm ^{5, 11}.

PROPAGATION DETAILS

Ecotype

Propagation Goal	Plants
Propagation Method	Seed (Cuttings of <i>Alnus rubra</i> have a difficult time rooting ⁵)
Product Type	Bareroot
Stock Type	Seedlings
Time to Grow	One growing season to have a seedling ready to be out planted.
Target Specifications	Roots should fully occupy the container but seedlings are not rootbound.
Propagule Collection Instructions Processing/Propagule Characteristics	<p>Seeds should be ready in early autumn, anywhere from early September in some regions to mid October in other regions ¹³.</p> <p>Collect catkins before disintegration while they are still green. Lay flat in a controlled humid environment, such as in a greenhouse, to encourage disintegration. Dry catkin in a kiln at 32°C for 24 hours no later than 6 weeks after collecting. Strip seeds from catkin using hands ⁸.</p> <p>Another way to collect the seeds is to check the strobile of the cone-like structure. If the strobile is easy to twist, the seeds should be ready. The seeds that are easiest to take and most often highest percent viable are on the top third of the tree ¹³.</p> <p>After collection, keep seeds in a highly ventilated and stable environment, such as a refrigerator, to maintain seed integrity before cleaning ⁵.</p> <p>Alternatively, store seeds in an airtight container for one to two years while seeds are viable ⁴.</p>
Pre-Planting Propagule Treatments	<p>Keep seeds in cold stratification for three to five weeks before sowing to overcome seed dormancy and have the highest percent viability ^{8,16}.</p> <p>Germination is higher in light environments versus dark environments ³.</p>
Growing Area Preparation / Annual Practices for Perennial Crops	<p>400 seeds per square meter ⁸.</p> <p>The media may be fertilized with low-nitrogen and high phosphorus and should absorb water well with low drainage ¹³.</p>
Establishment Phase Details	<p>Seeds should not be planted more than 1 cm below the soil, due to their small size ¹³. Soil may be sterilized, but if so, it is suggested to inoculate the soil with cultured <i>Frankia</i> to increase the nodulation of <i>Alnus rubra</i> roots ¹.</p> <p>Seedlings may be transplanted from late winter to mid spring, where the same media may be used from the bare root bed ¹⁴.</p>
Length of Establishment Phase	1 month ¹⁴
Active Growth Phase	<p>Water during times of minimum wind.</p> <p>Hand weed the seedlings.</p> <p>Pests do not usually occur during this growing phase, but in large populations of alders, it is common to see the harmless tent caterpillars ^{12, 14}.</p>
Length of Active Growth Phase	3 months
Hardening Phase	<p>Decrease frequency of watering to induce hardening in late summer.</p> <p>Roots will spiral if potted in a container. To avoid this, transplant plant into a larger container before winter.</p>

Length of Hardening Phase	3 months
Harvesting, Storage and Shipping	Seedlings should remain in their containers while shipping or storing.
Length of Storage	Keep stored in a greenhouse. Plant before or soon after roots have spiraled ¹⁴ .
Guidelines for Outplanting / Performance on Typical Sites	
Other Comments	

INFORMATION SOURCES

References	<p>¹ A. J. MOFFAT, Nursery Sterilization and Inoculation Regimes for Alder Production, <i>Forestry: An International Journal of Forest Research</i>, Volume 67, Issue 4, 1994, Pages 313–327, https://doi.org/10.1093/forestry/67.4.313</p> <p>² Bong. “Alnus Rubra - Bong.” <i>Pfaf Plant Search</i>, 2012, Plants for a Future, pfaf.org/user/Plant.aspx?LatinName=Alnus%2Brubra.</p> <p>³ Bormann, Bernard T. <i>Ecological Implications of Phytochrome-Mediated Seed Germination in Red Alder</i>. 4th ed., vol. 29, Forest Sciences, by the Society of American Foresters, 1983, pp. 734–738, <i>Ecological Implications of Phytochrome-Mediated Seed Germination in Red Alder</i>.</p> <p>⁴ Dirr, Michael, and Charles W. Heuser. <i>The Reference Manual of Woody Plant Propagation</i>. Timber Press, Inc., 2006.</p> <p>⁵ “Index of Species Information.” <i>Fire Effects Information System (FEIS)</i>, U. S. Department of Agriculture, www.fs.fed.us/database/feis/plants/tree/alnrub/all.html.</p> <p>⁶ Jensen E, W. Randall, R. Keniston, and D. Bever. <i>Manual of Oregon Trees and Shrubs</i>. Eighth edition. Oregon State University, Corvallis, OR. 2002</p> <p>⁷ “Landscape Plants.” <i>Alnus Rubra Landscape Plants Oregon State University</i>, College of Agriculture Sciences, Department of Horticulture, landscapeplants.oregonstate.edu/plants/alnus-rubra.</p> <p>⁸ MacDonald, Bruce. <i>Practical Woody Plant Propagation for Nursery Growers</i>. Timber Press, 1986.</p> <p>⁹ Maia, Eric. “Native Plant Guide.” <i>Red Alder: Alnus Rubra - Native Plant Guide</i>, Information and Services for King County, Washington, 2013, green2.kingcounty.gov/gonative/Plant.aspx?Act=view&PlantID=11.</p>
------------	---

	<p>¹⁰ “Plant Profile for <i>Alnus Rubra</i> Bong. Red Alder.” <i>Natural Resources Conservation Service</i>, United States Department of Agriculture, plants.usda.gov/core/profile?symbol=alru2.</p> <p>¹¹ “Red Alder (<i>Alnus Rubra</i>).” <i>Red Alder (Alnus Rubra) Oregon Wood Innovation Center</i>, Oregon State University, owic.oregonstate.edu/red-alder-alnus-rubra.</p> <p>¹² Robson, Kathleen A., et al. <i>Encyclopedia of Northwest Native Plants for Gardens and Landscapes</i>. Timber Press, 2008.</p> <p>¹³ Rose, Robin, et al. <i>Propagation of Pacific Northwest Native Plants</i>. Oregon State University Press, 1998.</p> <p>¹⁴ Steinfeld, David. “Native Plant Network.” <i>Reforestation, Nurseries and Genetics Resources</i>, Native Plant Network, 2003, npn.rngr.net/renderNPNProtocolDetails?selectedProtocolIds=betulaceae-alnus-2376.</p> <p>¹⁵ “The Plant List - A Working List for All Plant Species.” <i>Alnus Rubra</i> Bong. - <i>The Plant List</i>, The Plant List, 2010, www.theplantlist.org/tpl/record/kew-6584.</p> <p>¹⁶ <i>The Woody Plant Seed Manual</i>. vol. 727, United States Forest Service, 2008, p. 101, <i>The Woody Plant Seed Manual</i>.</p>
Other Sources Consulted	<p>Grant, John A., and Carol L. Grant. <i>Trees and Shrubs for Pacific Northwest Gardens</i>. Pacific Books, 1974.</p> <p>Potash, Laura L, and Carol A Aubry. <i>Olympic National Forest Native Plant Notebook</i>. Unpublished Manuscript on File at U.S. Department of Agriculture, Forest Service, Olympic National Forest, Olympia, WA., 2007, pp. 40-+ appendices, <i>Olympic National Forest Native Plant Notebook</i>.</p>
Protocol Author	Chloe May
Date Protocol Created or Updated	05/29/2019

Plant Propagation Protocol
 ESRM 412 – Native Plant Production
 JD Bakker
 Spring 2007

Family Names	
Family Scientific Name:	<i>Betulaceae</i>
Family Common Name:	Alder, Birch
Scientific Names	
Genus:	<i>Alnus</i>
Species:	<i>rubra</i>
Species Authority:	Bong.
Variety:	
Sub-species:	
Cultivar:	
Authority for Variety/Sub-species:	
Common Synonym	
Genus:	<i>Alnus</i>
Species:	<i>Oregona</i>
Species Authority:	Nutt. (ITIS, 2007)
Variety:	<i>Pinnatisecta</i>
Sub-species:	
Cultivar:	
Authority for Variety/Sub-species:	Starker (ITIS, 2007)
General Information	
Common Name:	red alder, Oregon alder, Pacific coast alder, alder (AgroForestry Tree Database, 2007)
Species Code (as per USDA Plants database):	ALRU2
Ecotype:	
Date Entered or Updated (MM/DD/YY):	04/10/2007
General Distribution (elevation range, ecosystems, etc):	Occurs along streams and in moist bottoms, and on moist mountain slopes. Ranges from southeastern Alaska southward through western British Columbia, and on the west side of Cascade mountains as far south as southern California. Elevational range: sea level to 3,500 feet. (Jensen et al, 2002)
Propagation Details	
Propagation Goal:	Plants
Propagation Method :	Seed
Product Type :	Bareroot (field grown)
Stock Type:	1 + 0 seedlings
Time to Grow:	A plantable bareroot seedling can be produced at the nursery in one growing season. (Potash and Aubry, 1997)
Target Specifications:	A plantable bareroot seeding which can be planted in the field, or lifted and transplanted into containers or another bed for growing on.
Propagule Collection:	<i>A. rubra</i> reaches sexual maturity after 10 years (Young, 1992). The tree is generally monoecious, with separate male and female catkins developing on previous years' growth. Male catkins are thin and 10-15 cm long while female catkins are thicker and shorter, 0.8-1.2 cm long. The seeds are tiny winged nutlets found inside woody cone-like fruits

	<p>called strobiles. The strobiles are 1.3-2.5 cm long, cylindrical shaped, persistent, and turn from green to yellow to brown as they ripen. (Rose et al, 1998) Seed dispersal begins in late September in the middle of the species range, somewhat earlier in Alaska, and several weeks later in California. <i>A. rubra</i> is a consistent and prolific producer of seed. (Agroforestry Tree Database, 2007) Handpicking is particularly convenient for <i>Alnus</i> sp. (Macdonald, 1986) Seed maturity can be checked by twisting a strobile along the long axis. If it is mature, the strobile will twist easily and scales part slightly. (Rose et al, 1998) Another way to tell when cones are ripe is that the stem below the cluster of cones thins noticeably and you are able to break off clusters at this point. (Potash and Aubry, 1997) To determine whether or not the cones on a certain alder tree are worth collecting take a knife and split a few cones lengthwise. The light colored seeds are located between the cone scales and near their base. A good seed crop is indicated by a count of five or more seeds. (Hibbs and Ager, 1989) Collect from standing or recently felled trees when the bracts start to separate on the earliest strobiles. Collect strobiles by stripping from the branches and blow out any leaf litter with a fan. Strobiles found on the upper third of the tree usually contain the most viable seed. (Rose et al, 1998) Collect seed from a source as geographically and climactically close as possible to the final planting site of seedlings, for greatest seedling growth rates and viability. (Hamann et.al, 2000) While one study found high elevation stands to have significantly higher viable seed production per catkin than low elevation stands, (Markham, 2002) another found this correlation to be negligible. The second study found the best growth performance associated with seed sources from warmer areas with low moisture demands. In general, this combination of climate conditions exists in low-elevation coastal areas; in this study, all seed collection locations fitting this description were in northwestern Washington state. (Hibbs et al, 1994)</p>
<p>Propagule Processing/Propagule Characteristics (including seed density (# per pound), seed longevity, etc):</p>	<p>Once picked, dry immediately with good air circulation (on screens or in mesh bags at 16-27° C) to prevent molding. Most strobiles will open after being in above storage treatment for several weeks. They can be opened in a shorter time by kiln drying at 32° C for 24 hours within 4-6 wks of collecting. (Macdonald, 1986, Rose et al, 1998) Most seeds fall out of the strobiles during the drying process. The remainder, if needed, may be extracted by tumbling and shaking. (Young, 1992) Once cleaned, seed can be stored for a short time by refrigeration. For long term storage, dry to less than 10% moisture content and store in moisture proof containers at -12 to -13° C for up to five years. (Rose, et al, 1998) Moderate seed crops are produced annually and large crops occur every 3-5 years. The seeds are very light, numbering 800,000-3,000,000/kg (Young, 1992) Higher number of seeds per gram may indicate lower quality. (Young, 1992)</p>

Pre-Planting Propagule Treatments (cleaning, dormancy treatments, etc):	<i>Alnus rubra</i> seeds have no or little dormancy. (Burns and Honkala, 1990) Seeds germinate easily. Generally, no stratification is required for red alder seed, (Rose et al, 1998) although seed not planted in the same fall it was collected may be cold-stratified for 30 days to break seed dormancy. (Stevens, 1993)
Growing Area Preparation/Annual Practices for Perennial Crops	Will germinate in fall or spring on moist mineral soil in full sunlight. (Baskin, 2002) Sowing smaller seed such as alder is made easier by bulking up the seed with fine peat moss or fine sand to a known volume. (Macdonald, 1986) Sow at less than one pound per acre. (Potash and Aubry, 1997) Seeds germinate at 24/16° C. Due to their size, seeds should not be covered, or they will not germinate. (Rose et al, 1992) If seeds are to be covered depth should be no greater than 1 cm. Seedling growth is stimulated by soil sterilization before sowing seeds. This decreases root nodulation, however, so it is recommended to follow sterilization with inoculation with a cultured strain of the nitrogen fixing bacteria <i>Frankia</i> , which greatly increases seedling root nodulation. (Moffat, 1994, Martin and Myrold, 2003) Pre-germination inoculation with cultured strains of <i>Frankia</i> has been shown to increase seedling growth more than treatment with standard nursery fertilizers. (Wheeler, et al 1991)
Establishment Phase:	It may be necessary to give 3-4 applications of water each day during dry weather after sowing seed until the radicle has developed sufficiently to function and maintain moisture status. This is critical during the first few days for small seeds such as <i>Alnus</i> . (Macdonald, 1986) Results of one study indicate that red alder would benefit in total plant growth from increased ambient CO ₂ during seed stratification, germination, and establishment. The same study also indicates this increased ambient CO ₂ would increase the seedlings drought tolerance. (Hibbs et. al, 1995)
Length of Establishment Phase:	Few weeks
Active Growth Phase:	Seedlings are irrigated in the early morning when the wind is at a minimum. Irrigation scheduling is based on the moisture content of the seed bed. Generally, the bed is not allowed to dry down much below field capacity. During periods of maximum summer temperatures, seedlings are irrigated every 2 to 3 days for up to 4.5 hours an irrigation. Seedlings are not cooled in the afternoon with overhead irrigation. Seedlings are monitored for pests but generally insects and diseases do not occur in this species and stocktype. Seedlings must be hand-weeded every 6 weeks. (Steinfeld, 2003)
Length of Active Growth Phase:	3 months
Hardening Phase:	Hardening begins in the late summer with a reduction in the frequency of irrigations. Generally seedlings are hardy to any nighttime low temperatures that are encountered in the fall and winter months. For winter freezes where temperatures reach into the low teens, seedbeds can be covered with poly hoop houses until the event has passed.
Length of Hardening Phase:	3 months
Harvesting, Storage and Shipping:	1 + 0 bare root seedlings are lifted in the fall and can be

	transported immediately to a planting site, to another nursery, potted up or put in another bare root bed to grow on another year.
Length of Storage:	NA
Guidelines for Outplanting / Performance on Typical Sites:	Seedlings are typically planted in the early spring when snow has left the site and soils are no longer frozen or in early fall. Seedlings are placed in holes so that the top of the plug is several inches below the surface of the soil. Soil is placed back over the planting hole so that no media is exposed. The resulting depression around the seedling will store rain or irrigation water. Competitive grasses and forbs are often removed from around the seedling and the site then mulched. Seedlings are protected from animal damage with deer repellent or netting. Seedlings that are planted in the early fall are usually watered after planting. (Steinfeld, 2003) Underplanting in thinned stands of Douglas fir, which is done for various reasons, has good success, and plants may not need to be protected from deer or elk browse. (Maas-Hebner et al, 2005) One study suggests that rectangular planting of red alder at dense spacing enhance stand differentiation, accelerate competition-related mortality, and thus lead to improved growth of surviving trees. (Debell, 2002)
Other Comments:	
References:	<p>AgroForestry Tree Database. URL: http://www.worldagroforestrycentre.org. (accessed 8 April 2007).</p> <p>Baskin, Carol J.; Baskin, Jerry M. 2002. Propagation protocol for production of container <i>Alnus rubra</i> plants; University of Kentucky, Lexington, Kentucky. In: Native Plant Network. URL: http://www.nativeplantnetwork.org (accessed 8 April 2007).</p> <p>Burns, R. and B. Honkala 1990. Silvics of North America, Volume 2, Hardwoods. Agricultural Handbook 654. U.S. Department of Agriculture, Forest Service, Washington, D. C. 877 p.</p> <p>DeBell DS, Harrington CA. "Density and rectangularity of planting influence 20-year growth and development of red alder." Canadian Journal of Forest Research 32 (7): 1244-1253 Jul 2002 .</p> <p>Hamann A, Koshy MP, Namkoong G, Ying CC. "Genotype x environment interactions in <i>Alnus rubra</i>: developing seed zones and seed-transfer guidelines with spatial statistics and GIS." Forest Ecology and Management 136 (1-3): 107-119 Oct 1 2000.</p> <p>Hibbs DE, Bouvarel L, Ducros ET. "Performance of Red Alder seed sources in France." Canadian Journal of Forest Research 24 (5): 1008-1014 May 1994</p>

Hibbs, D.E. and A.A. Ager. 1989. Red Alder: Guidelines for Seed Collection, Handling and Storage. Forestry Research Laboratory, Oregon State University. Corvallis, OR. Special Publication 18, 6p.

Integrated Taxonomic Information System. URL: <http://www.itis.gov> (accessed 8 April 2007).

Jensen E, W. Randall, R. Keniston, and D. Bever. Manual of Oregon Trees and Shrubs. Eighth edition. Oregon State University, Corvallis, OR. 2002

Potash, L., and C.A. Aubry. 1997. *Native plant notebook: Mt. Baker-Snoqualmie National Forest*. North Cascades Institute, Sedro Woolley, WA.

Maas-Hebner KG, Emmingham WH, Larson DJ, Chan SS. 2005. "Establishment and growth of native hardwood and conifer seedlings underplanted in thinned Douglas-fir stands." *Forest Ecology and Management* 208 (1-3): 331-345.

Macdonald, Bruce. Practical Woody Plant Propagation for Nursery Growers. Timber Press, Portland, OR 1986

Markham, JH. "A hierarchical analysis of seed production by *Alnus rubra*." *American Midland Naturalist* 148 (2): 246-252 Oct 2002.

Martin KJ, Tanaka Y, Myrold DD. "Dual inoculation increases plant growth with *Frankia* on red alder (*Alnus rubra* Bong.) in fumigated nursery beds." *Symbiosis* 34 (3): 253-260 2003.

Moffat, AJ. "Nursery Sterilization and Inoculation regimes for alder production" *Forestry* 67 (4): 313-327 1994

Rose, Robin, Caryn Chachulski, Dinae Haase. Propagation of Pacific Northwest Plants. Oregon State University Press. Corvallis, OR. 1998

Steinfeld, David. 2003. Propagation protocol for production of container *Alnus rubra* plants; J. Herbert Stone Nursery, Central Point, Oregon. In: Native Plant Network. URL: <http://www.nativeplantnetwork.org> (accessed 8 April 2007).

Stevens ML, and Vanbianchi R. "Restoring Wetlands in Washington" Publication #93-17. Washington State Department of Ecology, Olympia, Washington. 1993.

Wheeler CT, Holingsworth MK, Hooker JE, McNeill JD, Mason WL, Moffat AJ, Sheppard LJ. "The effect of inoculation with either cultured *Frankia* or crushed nodules on nodulation and growth of *Alnus rubra* and *Alnus*

	<p><i>glutinosa</i> seedlings in forest nurseries.” Forest Ecology and Management 43 (1-2): 153-166 SEP 1991</p> <p>Young, James, and Cheryl Young. Seeds of Woody Plants in North America. Dioscorides Press, Portland, OR. 1992</p>
<p>Propagator (Author) That Developed This Protocol</p>	
<p>First Name:</p>	<p>Katie</p>
<p>Last Name:</p>	<p>Pencke</p>

Plant Data Sheet

Plant Data Sheet

Red alder, *Alnus rubra*

Range

Pacific Northwest coastal areas into SE Alaska (Silvics of N.A. 1990)

Climate, Elevation

Low winter temperatures and lack of precipitation limit red alder. Locally, elevations below 450 m (Silvics of N.A. 1990)

Local occurrence (where, how common)

Forest understory, disturbed sites, riparian, wetlands

Habitat preferences

Moist soils, sunny sites, disturbances (Silvics of N.A. 1990)

Plant strategy type/successional stage (stress-tolerator, competitor, weedy/colonizer, seral, late successional)

Weedy, colonizer and seral. Nitrogen-fixer

Associated species

Big-leaf maple, Douglas fir, willow, western red cedar, western hemlock, grand fir, black cottonwood (Silvics of N.A. 1990)

May be collected as: (seed, layered, divisions, etc.)

Seed, stump-sprouts, greenwood cuttings, mound layering (Silvics of N.A. 1990)

Collection restrictions or guidelines

Seeds shed in September (Alaska) through December (California) (Silvics of N.A. 1990)

Seed germination (needs dormancy breaking?)

Will germinate in fall or spring on moist mineral soil in full sunlight. No or little dormancy. (Silvics of N.A. 1990)

Seed life (can be stored, short shelf-life, long shelf-life)

Short; very small seeds. Store no more than two seasons.

Recommended seed storage conditions

Store in paper sacks in cool, dry environment

Propagation recommendations (plant seeds, vegetative parts, cuttings, etc.)

Collect cones, separate seeds, plant seeds. Pull-ups also convenient and plentiful. (Rose et al. 1996)

Soil or medium requirements (inoculum necessary?)

Will grow in non-soil medium, but requires inoculum for nitrogen-fixing actinobacteria *Frankia* in nodules.

Installation form (form, potential for successful outcomes, cost)

Seeds, container-plants grown from seeds, green cuttings, bare root (Silvics of N.A. 1990)

Recommended planting density

6' centers (Stevens and Vanbiatchi 1994)

Care requirements after installed (water weekly, water once, never water, etc.)

Requires moist soil (Stevens and Vanbiatchi 1994)

Plant Data Sheet

Normal rate of growth or spread; Lifespan
Fast growth: 30' at 5 yrs., 50' at 10 yrs. 80 ft. at 20 yrs. Mature at 60-70 yrs. Maximum age 100 yrs. (Silvics of N.A. 1990)

Sources cited

Burns, R. and B. Honkala 1990. Silvics of North America, Volume 2, Hardwoods. Agricultural Handbook 654. U.S. Department of Agriculture, Forest Service, Washington, D. C. 877 p.

Rose, R., C. Chachulski and D. Haase. 1996. Propagation of Pacific Northwest Native Plants: A Manual, Volume Two, First Edition. Nursery Technology Cooperative, Oregon State University, Corvallis, Oregon, 73 p.

Stevens, M. and R. Vanbiatchi. 1993. Restoring Wetlands in Washington: A Guidebook for Wetland Restoration, Planning and Implementation. Washington State Department of Ecology Publication 93-17, 110 p and Appendices.

Data compiled by: Kern Ewing, 14 Mar 2003