

2015-11

# The macroecology of phylogenetically structured hummingbird-plant networks

Martin Gonzalez, AM

<http://hdl.handle.net/10026.1/4865>

---

10.1111/geb.12355

Global Ecology & Biogeography

Wiley

---

*All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.*

1 **Article title:** The macroecology of phylogenetically structured hummingbird-plant networks

2 **Running title:** Macroecology of hummingbird-plant networks

3  
4  
5  
6  
7  
8 **Authors and affiliations:** Ana M. Martín González<sup>1,2</sup>, Bo Dalsgaard<sup>2</sup>, David Nogués-Bravo<sup>2</sup>,  
9 Catherine H. Graham<sup>3</sup>, Matthias Schleuning<sup>4</sup>, Pietro K. Maruyama<sup>2,5,6</sup>, Stefan

10 Abrahamczyk<sup>7,8</sup>, Ruben Alarcón<sup>9</sup>, Andréa C. Araujo<sup>10</sup>, Francielle P. Araújo<sup>5</sup>, Severino Mendes  
11 de Azevedo Jr<sup>11</sup>, Andrea C. Baquero<sup>2</sup>, Peter A. Cotton<sup>12</sup>, Tanja Toftemark Ingversen<sup>13</sup>, Glauco

12 Kohler<sup>14</sup>, Carlos Lara<sup>15</sup>, Flor Maria Guedes Las-Casas<sup>16</sup>, Adriana O. Machado<sup>6</sup>, Caio Graco  
13 Machado<sup>17</sup>, María Alejandra Maglianesi<sup>4,18</sup>, Jimmy A. McGuire<sup>19</sup>, Alan Cerqueira Moura<sup>17</sup>,

14 Genilda M. Oliveira<sup>20</sup>, Paulo E. Oliveira<sup>6</sup>, Juan Francisco Ornelas<sup>21</sup>, Licléia da Cruz

15 Rodrigues<sup>22</sup>, Liliana Rosero-Lasprilla<sup>23</sup>, Ana M. Rui<sup>24</sup>, Marlies Sazima<sup>5</sup>, Allan Timmermann<sup>25</sup>,

16 Isabela Galarda Varasin<sup>26</sup>, Jeferson Vizentin-Bugoni<sup>2,5</sup>, Zhiheng Wang<sup>2, 27</sup>, Stella Watts<sup>28</sup>,  
17 Carsten Rahbek<sup>2,29</sup>, Neo D. Martinez<sup>1,30</sup>.

18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33 **1** Pacific Ecoinformatics and Computational Ecology Lab, 1604 McGee Ave, 94703 Berkeley,  
34 California, USA.

35  
36  
37 **2** Center for Macroecology, Evolution and Climate, Natural History Museum of Denmark,  
38 University of Copenhagen, DK-2100 Copenhagen Ø, Denmark.

39  
40  
41 **3** Department of Ecology and Evolution, Stony Brook University, 650 Life Sciences Building,  
42 Stony Brook, NY 11784-5245, USA.

43  
44  
45  
46 **4** Biodiversity and Climate Research Centre (BiK-F) and Senckenberg Gesellschaft für  
47 Naturforschung, Senckenberganlage 25, 60325 Frankfurt (Main), Germany.

48  
49  
50 **5** Instituto de Biologia, Universidade Estadual de Campinas, Campinas, São Paulo, Brazil.

51  
52  
53 **6** Instituto de Biologia, Universidade Federal de Uberlândia -UFU, Uberlândia, Minas Gerais,  
54 Brazil.

55  
56  
57  
58  
59  
60

- 1  
2 26 **7** Nees Institute of Plant Biodiversity, Meckenheimer Allee 170, 53115 Bonn, Germany.  
3  
4 **8** Institute of Systematic Botany, Zollikerstr.107 Zurich, Switzerland.  
5  
6 28 **9** Biology Program, California State University Channel Islands, 93012 Camarillo, California,  
7  
8 USA.  
9  
10  
11 30 **10** Centro de Ciências Biológicas e da Saúde, Universidade Federal de Mato Grosso do Sul,  
12  
13 79070-900, Campo Grande, Mato Grosso do Sul, Brazil.  
14  
15 32 **11** Department of Biology, Rural Federal University of Pernambuco, Recife, Pernambuco,  
16  
17 Brazil.  
18  
19 34 **12** Marine Biology & Ecology Research Centre, Plymouth University, Plymouth PL4 8AA, UK.  
20  
21 **13** Stationsvej 33, DK-6580 Vamdrup, Denmark.  
22  
23  
24 36 **14** Instituto Nacional de Pesquisas da Amazônia, Av. André Araújo 2936, Petrópolis, CEP  
25  
26 69080-971, Manaus, Amazonas, Brazil.  
27  
28 38 **15** Centro de Investigación en Ciencias Biológicas, Universidad Autónoma de Tlaxcala. Km  
29  
30 10.5 Autopista Tlaxcala-San Martín Texmelucan, San Felipe Ixtacuixtla, 90120, Tlaxcala,  
31  
32 México.  
33  
34 **16** Department of Zoology, Federal University of Pernambuco, Recife, Pernambuco, Brazil.  
35  
36  
37 42 **17** Laboratório de Ornitologia, Departamento de Ciências Biológicas, Universidade Estadual  
38  
39 de Feira de Santana, Feira de Santana, 44036-900, Bahia, Brazil.  
40  
41  
42 44 **18** Vicerrectoría de Investigación. Universidad Estatal a Distancia (UNED), San José, Costa  
43  
44 Rica.  
45  
46 46 **19** Museum of Vertebrate Zoology and Department of Integrative Biology, University of  
47  
48 California, Berkeley, California 94720.  
49  
50  
51 48 **20** Instituto Federal do Triângulo Mineiro, Campus Uberlândia, Uberlândia, Minas Gerais,  
52  
53 Brazil.  
54  
55 50 **21** Departamento de Biología Evolutiva, Instituto de Ecología AC, Carretera antigua a  
56  
57  
58  
59  
60

1 Coatepec 351, El Haya, Xalapa, Veracruz 91070, Mexico.

2  
3  
4 52 **22** Laboratory of Ornithology, Department of Zoology, ICB, Minas Gerais Federal University,  
5  
6 Post office box 486, 31270-901, Belo Horizonte, Minas Gerais, Brazil.

7  
8 54 **23** Escuela de Ciencias Biológicas, Grupo de Investigación Biología para la Conservación,  
9  
10 Universidad Pedagógica y Tecnológica de Colombia, Colombia.

11  
12 56 **24** Departamento de Ecologia, Zoologia e Genética, Instituto de Biologia, Universidade  
13  
14 Federal de Pelotas, Capão do Leão, Rio Grande do Sul, Brazil.

15  
16  
17 58 **25** Section for Ecoinformatics and Biodiversity, Department of Bioscience, Aarhus University,  
18  
19 Ny Munkegade 114, DK-8000 Aarhus C, Denmark.

20  
21 60 **26** Laboratório de Ecologia Vegetal, Departamento de Botânica, Universidade Federal do  
22  
23 Paraná, 81531-980 Curitiba, Paraná, Brazil.

24  
25  
26 62 **27** Department of Ecology and Key Laboratory for Earth Surface Processes of the Ministry of  
27  
28 Education, College of Urban and Environmental Sciences, Peking University, Beijing  
29  
30 100871, China.

31  
32  
33 **28** Landscape and Biodiversity Research Group, Department of Geographical and  
34  
35 Environmental Sciences, University of Northampton, Avenue Campus, St George's Avenue,  
36  
37 Northampton NN2 6JD, UK.

38  
39 68 **29** Imperial College, London, Silwood Park Campus, Ascot, Berkshire SL5 7PY, UK.

40  
41 **30** Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, AZ 85721  
42  
43 USA.  
44  
45

46  
47  
48 **Email addresses:**

49 Ana M. Martín González <ana.maria.martingonzalez@gmail.com>

50  
51  
52 74 Bo Dalsgaard <bo.dalsgaard@snm.ku.dk>

53 David Nogués-Bravo <dnogues@snm.ku.dk>

- 1  
2 76 Catherine Graham <musaphaga@yahoo.com>  
3  
4 Matthias Schleuning <Matthias.Schleuning@senckenberg.de>  
5  
6 78 Stefan Abrahamczyk <stefan.abrahamczyk@uni-bonn.de>  
7  
8 Ruben Alarcon <ruben.alarcon@csuci.edu>  
9  
10 80 Andréa Araujo <andrea.c.araujo@uol.com.br>  
11  
12 Francielle P. Araújo <franciaralp@yahoo.com.br>  
13  
14  
15 82 Severino Mendes de Azevedo Jr <smaj@db.ufrpe.br>  
16  
17 Andrea Baquero <baquero.andrea@gmail.com>  
18  
19 84 Pete Cotton <P.Cotton@plymouth.ac.uk>  
20  
21 Tanja Ingversen <Tanja.ingversen@gmail.com>  
22  
23  
24 86 Glauco Kohler <certhiaxis@gmail.com>  
25  
26 Carlos Lara <laracar@posgradouatx.com.mx>  
27  
28 88 Flor Maria Guedes Las-Casas <flormarialc@hotmail.com>  
29  
30 Adriana O. Machado <adrianaoliveiramachado@yahoo.com.br>  
31  
32  
33 90 Caio Graco Machado <gracom@uol.com.br>  
34  
35 Alejandra Maglianesi <Alejandra.Maglianesi@senckenberg.de>  
36  
37 92 Pietro Kiyoshi Maruyama <pietrokiyoshi@gmail.com>  
38  
39 Jimmy A. McGuire <m McGuirej@calmail.berkeley.edu>  
40  
41 94 Alan Cerqueira Moura <daniellcerq@yahoo.com.br>  
42  
43 Genilda M. Oliveira <genioliv@yahoo.com.br>  
44  
45  
46 96 Paulo Eugenio Oliveira <poliveiragm@gmail.com>  
47  
48 Juan Francisco Ornelas <francisco.ornelas@inecol.edu.mx>  
49  
50  
51 98 Licléia Rodrigues <licleia@yahoo.com.br>  
52  
53 Liliana Rosero <lilianaroslal@yahoo.com.mx>  
54  
55 100 Ana M. Rui <ana.rui@ufpel.tche.br>  
56  
57  
58  
59  
60

1 Marlies Sazima <msazima@unicamp.br>

2  
3  
4 102 Allan Timmermann <allantimmermann@yahoo.dk>

5  
6 Isabela Galarda Varassin <isagalarda@gmail.com>

7  
8 104 Jeferson Vinzentin-Bugoni <jbugoni@yahoo.com.br>

9  
10 Zhiheng Wang <zhiheng.wang@pku.edu.cn>

11  
12 106 Stella Watts <hummingbird\_pe@yahoo.com>

13  
14 Carsten Rahbek <crahbek@snm.ku.dk>

15  
16  
17 108 Neo Martinez <neo@peacelab.net>

18  
19  
20  
21  
22 110 **Corresponding author:** Ana M. Martín González [ana.maria.martingonzalez@gmail.com](mailto:ana.maria.martingonzalez@gmail.com)

23  
24  
25  
26 112 **Type of article:** Research paper

27  
28 **Number of words in abstract:** 298

29  
30 114 **Number of words in main text:** 5266

31  
32 **Number of references:** 52

33  
34 116 **Number of tables in main text:** 1

35  
36 **Number of figures in main text:** 3

37  
38 118 **Number of supplementary appendices:** 5

39  
40 **Number of supplementary tables:** 3

41  
42 120 **Number of supplementary figures:** 1

43  
44 **Keywords:** community ecology, current climate, historical climate, hummingbird

45  
46  
47  
48 122 biogeography, specialization, macroecology, modularity, phylogenetic signal, pollination,  
49 quantitative networks.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

124 **ABSTRACT**

**Aim** To investigate the association between species richness, species' phylogenetic signal, insularity and historical and current climate with hummingbird-plant network structure.

**Location** 54 communities along a c. 10,000 kilometer latitudinal gradient across the Americas (39°N - 32°S), ranging from sea level to c. 3700 m asl, located on the mainland and on islands, and covering a wide range of climate regimes.

**Methods** We measured null-modeled corrected complementary specialization and bipartite modularity (compartmentalization) in networks of quantitative interactions between hummingbird and plant species. Using an ordinary least squares multi-model approach, we examined the influence of species richness, phylogenetic signal, insularity, and current and historical climate conditions on network structure.

**Results** Phylogenetically-related species, especially plants, showed a tendency to interact with a similar array of partners. The spatial variation in network structure exhibited a constant association with species' phylogeny ( $R^2=0.18-0.19$ ). Species richness and environmental factors showed the strongest associations with network structure ( $R^2=0.20-0.44$ ;  $R^2=0.32-0.45$ , respectively). Specifically, higher levels of complementary specialization and modularity were associated to species-rich communities and communities in which closely-related hummingbirds visited distinct sets of flowering species. On the mainland, warmer temperatures and higher historical temperature stability associated to higher levels of complementary specialization.

**Main conclusions** Previous macroecological studies of interaction networks have highlighted the importance of environment and species richness in determining network structure. Here, for the first time, we report an association between species phylogenetic signal and network structure at macroecological scale. Specifically, null model corrected complementary specialization and modularity exhibited a positive association with species

1 richness and a negative association with hummingbird phylogenetic signal, indicating that  
2  
3  
4 150 both high richness and high inter-specific competition among closely-related hummingbirds  
5  
6 exhibit important relationships with specialization in hummingbird-plant networks. Our results  
7  
8 152 document how species richness, phylogenetic signal and climate associate with network  
9  
10 structure in complex ways at macroecological scale.  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review



## 154 INTRODUCTION

1  
2  
3  
4 An urgent challenge facing contemporary ecologists is to understand how ecological,  
5  
6 156 evolutionary and environmental mechanisms affect the structure and function of ecological  
7  
8 communities. Better understanding of how these processes form and maintain communities  
9  
10  
11 158 should help us to construct more robust theories and models of community dynamics that  
12  
13 better predict how species and communities may respond to disturbance (Vázquez *et al.*,  
14  
15 160 2009a; Woodward *et al.*, 2010). For this purpose, the use of network approaches to study  
16  
17 complex communities of interacting species, i.e. networks representing species as nodes and  
18  
19  
20 162 interactions as links between species, has significantly advanced our understanding of  
21  
22 ecological systems (Woodward *et al.*, 2010).

23  
24 164 For instance, research on mutualistic networks such as plant-pollinator or plant-  
25  
26 frugivore interactions, has revealed that these communities typically exhibit a number of  
27  
28 166 architectural attributes, such as complementary specialization and modularity, which differ  
29  
30 significantly from random interactions among species (Olesen *et al.*, 2007; Blüthgen 2010;  
31  
32 168 Dalsgaard *et al.*, 2011, 2013; Schleuning *et al.*, 2012, 2014; Trøjelsgaard & Olesen, 2013;  
33  
34 Dormann & Strauss, 2014; Sebastián González *et al.*, 2015). Complementary specialization  
35  
36 170 measures the exclusiveness in species' interactions. Specifically, it is a measure of the  
37  
38 deviation from a neutral scenario in which species interact solely according to their  
39  
40 172 availability, measuring species' availability either as its abundance or its interaction frequency  
41  
42 (Blüthgen *et al.*, 2006). Modularity quantifies whether species interact more frequently with  
43  
44 174 subsets of available species within a community, forming modules of densely interacting  
45  
46 plants and hummingbirds, with loose connections between them (Olesen *et al.*, 2007). Such  
47  
48 176 modules have been suggested to reflect co-evolutionary units (Olesen *et al.*, 2007), within-  
49  
50 network functional specialization (Maruyama *et al.*, 2014), and phenological units of species  
51  
52 in environments with strong climatic seasonalities (Martín González *et al.*, 2012; Schleuning  
53  
54  
55  
56  
57  
58  
59  
60

1 *et al.*, 2014).

2  
3  
4 180 These architectural attributes have a profound impact on the dynamics of ecological  
5  
6 communities. For example, higher levels of complementary specialization have been linked to  
7  
8 182 higher dependencies between species and hence, to a higher risk of secondary extinctions  
9  
10 (Blüthgen, 2010), whereas high levels of modularity may at first provide higher network  
11  
12 184 resilience to perturbations, as these are not typically spread across modules, but ultimately  
13  
14 may result into unconnected modules, community fragmentation and rapid species loss  
15  
16 186 (Thébault & Fontaine, 2010; Stouffer & Bascompte, 2011). Complementary specialization and  
17  
18 modularity are typically related, as some degree of modularity necessarily implies a certain  
19  
20 188 level of complementary specialization on a subset of species from the available species pool  
21  
22 (Dormann & Strauss, 2014). Nevertheless, these metrics measure different aspects of  
23  
24 190 interaction specialization and, hence, may show disparate associations with different  
25  
26 ecological, evolutionary and environmental factors.  
27  
28  
29

30 192 Thus, studies at macroecological scales, where differences among network structure  
31  
32 can be associated to varying ecological, evolutionary and environmental factors, may provide  
33  
34 194 valuable insights into the structure and dynamics of ecological networks (Dalsgaard *et al.*,  
35  
36 2011, 2013; Schleuning *et al.*, 2012, 2014; Trøjelsgaard & Olesen, 2013, Sebastián González  
37  
38 *et al.*, 2015). For instance, species richness may promote higher degrees of complementary  
39  
40 196 specialization and modularity by increasing interspecific competition and providing more  
41  
42 species to interact with, *i.e.* allowing a finer niche partitioning (e.g. Dalsgaard *et al.*, 2011;  
43  
44 198 Junker *et al.*, 2013). Additionally, evolution may structure biotic interactions via niche  
45  
46 conservatism, as closely-related species may display more similar phenotypes, spatial  
47  
48 200 distributions, and ecological interactions than distantly-related species (Webb *et al.*, 2002;  
49  
50 Ives & Godfray, 2006; Rezende *et al.*, 2007; Vázquez *et al.*, 2009a). Alternatively, closely-  
51  
52 202 related species experiencing strong interspecific competition may undergo niche partitioning,  
53  
54  
55  
56  
57  
58  
59  
60

1  
2 204 competitive exclusion, or resource-use complementarity (Webb *et al.*, 2002; Rezende *et al.*,  
3  
4 2009; Krasnov *et al.*, 2012). Environmental setting may also affect biotic interactions.  
5  
6 206 Historical climatic stability may promote specialization and modularity by providing species  
7  
8 more opportunity to co-evolve over longer periods of time compared to species from  
9  
10 208 climatically more unstable areas. For example, large late Quaternary glacial-interglacial  
11  
12 climate change is coupled with a decreased modularity in pollination networks (Dalsgaard *et*  
13  
14 *al.*, 2013), and decreased specialization in hummingbird-plant networks (Dalsgaard *et al.*,  
15  
16 210 2011). Likewise, current environmental conditions may influence the identity and strength of  
17  
18 species interactions by altering species' spatial distribution, phenophases and foraging  
19  
20 212 capability (Cruden, 1972; Martín González *et al.*, 2009; Dalsgaard *et al.*, 2011, 2013;  
21  
22 Schleuning *et al.*, 2012, 2014). For example, bird pollination has been reported to become  
23  
24 214 increasingly important and specialized in areas of high precipitation and low temperatures,  
25  
26 possibly because insect pollinator diversity and activity is lower in cold and wet environments  
27  
28 216 due to thermoregulatory constraints (Cruden, 1972; Dalsgaard *et al.*, 2009, 2011; Martín  
29  
30 González *et al.*, 2009). Finally, insularity may influence pollination network structure, with  
31  
32 significantly lower levels of specialization and modularity on islands compared to the  
33  
34 218 mainland (Olesen *et al.*, 2002; Dalsgaard *et al.*, 2013).  
35  
36  
37  
38  
39

40  
41  
42 222 Despite the recently increasing interest in determining the influence of species  
43  
44 richness, phylogenetic signal and past and current environmental conditions on network  
45  
46 224 structure (Dalsgaard *et al.*, 2011, 2013; Schleuning *et al.*, 2012, 2014; Sebastián González *et*  
47  
48 *al.*, 2015), very little is known about the relative importance of these factors. Only Schleuning  
49  
50 226 *et al.* (2014) studied their combined influence on the modular patterns of mutualistic  
51  
52 networks. Specifically, they used a dataset of 18 frugivore bird-plant networks, and showed  
53  
54 that modularity was high in areas having low temperatures and high temperature seasonality,  
55  
56 228 whereas species' phylogenetic history and historical climate exhibited no detectable  
57  
58  
59  
60

1 association to modularity. Here, we likewise examine how species richness, phylogenetic  
2  
3  
4 230 signal and environmental conditions associate to the structure of hummingbird-plant  
5  
6 networks, another type of mutualistic assemblage. To do so, we compiled a data-set  
7  
8 232 consisting of 54 high-resolution quantitative hummingbird-plant interaction networks, and  
9  
10 combined them with state-of-the-art species phylogenies and simulations of paleo and  
11  
12 234 current climate. In accordance with previous studies on pollination networks, we show that  
13  
14 species richness and climatic conditions influence complementary specialization and  
15  
16 236 modularity. Additionally, for the first time, we report an association between species  
17  
18 phylogenetic signal and network structure at macroecological scale. Specifically, communities  
19  
20 238 where closely-related hummingbirds interact with distinct sets of plant species exhibit higher  
21  
22 levels of complementary specialization and modularity.  
23  
24  
25  
26  
27

240

## 28 MATERIAL & METHODS

### 30 242 *Hummingbird-plant interaction networks*

32 We compiled a large data-set of 54 high-resolution and geographically widely distributed  
33  
34 244 hummingbird-plant interaction networks from published and unpublished sources (Fig. 1; see  
35  
36 Table S1 for full references, and Dalsgaard *et al.*, 2011 for a previous version of this dataset).  
37  
38 246 This dataset does not include networks which have not sampled the entire floral community,  
39  
40 i.e. we discarded studies which had focused only on ornithophilous species or on specific  
41  
42 floral or hummingbird groups. We also discarded illegitimate interactions, i.e. our database  
43  
44 248 focuses on interactions with potential for pollination. These networks describe the interaction  
45  
46 frequency between plants and hummingbirds, recorded as number of observed visits. In total,  
47  
48 250 our dataset is composed of 141 species of hummingbirds belonging to all major hummingbird  
49  
50 lineages (about 42% of all described species; McGuire *et al.*, 2014) and 824 plant species  
51  
52 252 belonging to 79 different plant families (Appendix S1).  
53  
54  
55  
56  
57  
58  
59  
60

1  
2 254 **Mutualistic** hummingbird-plant interaction networks constitute a particularly suitable  
3  
4 model system to explore large-scale patterns in network structure as there is ample  
5  
6 256 knowledge of the historical biogeography of hummingbirds, these assemblages are widely  
7  
8 distributed in a variety of ecosystems **across the Americas**, and they offer examples covering  
9  
10 258 the entire interaction specialization-generalization spectrum (Bleiweiss, 1998; Dalsgaard *et*  
11  
12 *al.*, 2011). Furthermore, our data consists of a single and monophyletic pollinator group  
13  
14 (Brown & Bowers, 1985; McGuire *et al.*, 2014), allowing us to explore the **association**  
15 260  
16 **between** phylogeny on overall network structure more precisely.  
17  
18

19 262 Finally, the spatial distribution of these networks extend from sea level to **c.** 3700 m  
20  
21 a.s.l. and spans a **c.** 10,000 kilometer gradient from 39°N to 32°S, from **tropical and**  
22  
23 264 subtropical ecosystems in South America to temperate communities in North America. The  
24  
25 spatial distribution includes both mainland and insular communities (Fig. 1), and consists of  
26  
27 266 communities from various environmental settings.  
28  
29  
30  
31

### 32 268 *Network composition and network metrics*

33  
34 Network interactions were quantified in terms of the number of interaction events between  
35  
36 270 hummingbird and plant species at particular locations. For each network we calculated two  
37  
38 complementary metrics that describe the degree of specialization or specificity of  
39  
40 272 hummingbird-plant interactions: **complementary specialization** (Blüthgen *et al.*, 2006) and  
41  
42 quantitative bipartite modularity (Dormann & Strauss, 2014). These metrics were correlated in  
43  
44 274 our dataset  $R^2=0.78$  (Table S2) **but** were chosen to more fully characterize **how** evolutionary  
45  
46 and environmental **factors associate to** specialized network structures. For the calculations  
47  
48 276 we used the R package *bipartite* 1.20 (Dormann *et al.*, 2009).  
49  
50  
51

52  
53 (1) **Complementary specialization ( $H_2'$ )** describes how species restrict their interactions  
54  
55 278 **relative to random expectations based on species' abundances/interaction frequency.**  
56  
57  
58  
59  
60

Complementary specialization is calculated as  $H_2 = - \sum_{i=1}^r \sum_{j=1}^c (p_{ij} \ln p_{ij})$ , where  $p_{ij}$  reflects the proportional number of interactions of each species relative to their availability, i.e., for their respective marginal total (in our case total interaction frequency) for  $r$  plant and  $c$  animal species. We illustrate this in Figure 1; for instance, from the example community A from Fig. 1,  $p_{1A}=0.99$  (which results from an interaction frequency of 155 between plant 1 and hummingbird A, over a marginal total of  $155+1=156$  for plant 1) and  $p_{A1}=1$  (155 over 155). Hence the interaction A-1 exhibits a high complementary specialization, as hummingbird A visits exclusively plant 1 and at a high frequency, while plant 1 is only visited, and at a very low frequency, by another hummingbird species. If a community is composed majoritarily by interactions with such high complementarity, the community will exhibit high values of complementary specialization (Blüthgen *et al.*, 2006). On the other hand, species C and 4 also interact 155 times, but in this case hummingbird C interacts with many other plant species and has a higher marginal total, lowering the complementary specialization of this pair ( $p_{C4}=0.36$ ,  $p_{4C}=1$ ).

We standardized complementary specialization as  $H_2' = \frac{H_{2\max} - H_2}{H_{2\max} - H_{2\min}}$ , so that  $H_2'$  ranges from minimum ( $H_2'=0$ ) to maximum ( $H_2'=1$ ) link selectiveness where species establish distinct and highly specific interactions far different than expected interactions (Blüthgen *et al.*, 2006). Thus,  $H_2'$  quantifies the deviation of the observed interactions from those expected under a neutral assumption that species' interactions are entirely determined by partner availability. This assumption minimizes the influence of rare interactions by causing frequent interactions to dominate  $H_2'$ .

(2) *Quantitative bipartite modularity* (QuanBiMo) is an algorithm which places species among an *a priori* unespecified number of modules, such that species interact at high frequencies within their module, and show few links and/or low frequency links with species

1 outside their module (Dormann & Strauss, 2014; Schleuning *et al.*, 2014; Fig. 1). Such  
2  
3  
4 304 partition is based on a hierarchical representation of interaction frequencies and optimal  
5  
6 allocation of species into modules. Specifically, the algorithm maximizes the bipartite version  
7  
8  
9 of Newman's modularity ( $Q$ ), so that  $Q = \frac{1}{2N} \sum (A_{ij} - K_{ij}) \delta(m_i, m_j)$ , where  $N$  reflects the total  
10 306  
11 number of interactions,  $A_{ij}$  the normalized number of interactions between species  $i$  and  $j$ ,  $K_{ij}$   
12  
13 the expected interaction probability between species  $i$  and  $j$  drawn from a neutral model of  
14 308  
15 interactions, and the indicator function  $\delta(m_i, m_j)$  equals 1 when species  $i$  and  $j$  are placed in  
16  
17 the same module and 0 otherwise. Modularity ranges from no ( $Q=0$ ) to maximum ( $Q=1$ )  
18  
19 310 modularity. We ran the QuanBiMo algorithm following the methodology established by  
20  
21 Schleuning *et al.* (2014) and the default specifications of the `computeModules` function in  
22  
23 312 bipartite, that is, for each network we chose the partition showing highest modularity from five  
24  
25 independent runs of the algorithm (Dormann & Strauss, 2014; Schleuning *et al.*, 2014).  
26  
27 314 Variations in the likelihood values of modularity were negligible (all  $SD < 0.05$ ).  
28  
29  
30  
31

32 316 As raw values for network metrics may be affected by species frequencies and  
33  
34 network connectance, network estimates for complementary specialization and modularity  
35  
36 318 were corrected using null models (Schleuning *et al.*, 2012, 2014; Dormann & Strauss, 2014).  
37  
38 Null models simulated matrices with the same number of species and interactions as the  
39  
40 320 empirical network, and with a species' interaction probability distribution drawn from observed  
41  
42 species' connectivity (vaznull model in bipartite; Vázquez *et al.*, 2007). Corrected metrics  
43  
44 322 were then calculated as the difference between the value of the empirical network and the  
45  
46 mean value obtained from 1000 and 100 null models for  $H_2'$  and QuanBiMo, respectively (as  
47  
48 in e.g. Schleuning *et al.*, 2012, 2014). As for the calculation of empirical QuanBiMo values,  
49  
50 324 for each of the 100 null matrices we used the maximum value of five independent runs of the  
51  
52 QuanBiMo algorithm (Schleuning *et al.*, 2014). By extracting the network structure achieved  
53  
54 326 under null conditions, corrected metrics quantify how much an empirical community departs  
55  
56  
57  
58  
59  
60

1  
2 328 from an average random one with an equivalent set of species, interactions and interaction  
3 probability distribution. Notice that for instance, as Figure 1 illustrates, a corrected value of  
4 0.3 gives no information on the uncorrected values, only that there is a 0.3 difference  
5  
6 330  
7  
8 between observed and null values, e.g. this value can result both from a raw observed value  
9  
10 332 of 0.4 and a mean null value of 0.1 (Fig. 1 community A), and from a raw observed value of  
11  
12 0.5 and a mean null value of 0.2 (Fig. 1 community B), etc.  
13  
14

15 334

### 17 *Analysis of phylogenetic signal in bipartite interaction networks*

18  
19 336 We measured the phylogenetic signal exhibited by hummingbird-plant networks by  
20  
21 quantifying the degree to which closely-related species share more interaction partners than  
22  
23 338 distantly-related species (Ives & Godfray, 2006). Species' interactions are considered to  
24  
25 exhibit a higher phylogenetic signal, when closely-related species share relatively more  
26  
27 340 interaction partners than distantly-related species (Ives & Godfray, 2006; Vázquez *et al.*,  
28  
29 2009b).  
30  
31

32  
33 342 We used state-of-the-art phylogenies of plants and hummingbirds to create variance-  
34  
35 covariance matrices, which quantify the phylogenetic relatedness of plants or hummingbirds  
36  
37 344 in each community, using the “vcv” function from the R package *ape* (Paradis *et al.*, 2004).  
38  
39 We fitted these vcv matrices to each observed bipartite interaction matrix through a linear  
40  
41 346 model using the “pblm” function from the R package *picante* (Kembel *et al.*, 2010). This  
42  
43 analysis results in two independent measures of the strength of the phylogenetic signal, one  
44  
45 348 for plants ( $d_{\text{plants}}$ ) and another for hummingbirds ( $d_{\text{hummingbirds}}$ ), together with an overall  
46  
47 measure of strength of the model fits for the entire community (measured as mean squared  
48  
49 error of the model, MSE). We evaluated three different models, one assuming no  
50  
51 350 phylogenetic signal ( $d_{\text{plants}}=d_{\text{hummingbirds}}=0$ ; Star model), one assuming a maximum  
52  
53 phylogenetic signal ( $d_{\text{plants}}=d_{\text{hummingbirds}}=1$ ; Brownian model), and a final one which  
54  
55 352  
56  
57  
58  
59  
60



1 incorporated the observed phylogenetic signals combined (estimated  $d_{\text{plants}}$  and  $d_{\text{hummingbirds}}$ ;  
2  
3  
4 354 Data model). We used the bootstrapping option to calculate confidence intervals for  $d_{\text{plants}}$   
5  
6 and  $d_{\text{hummingbirds}}$ . Networks where the these confidence intervals did not overlap zero or when  
7  
8 356 the  $\text{MSE}_{\text{Data}} < \text{MSE}_{\text{Star}}$  were considered to exhibit a significant phylogenetic signal (Ives &  
9  
10 Godfray, 2006; Vázquez *et al.*, 2009b). For a list of all species included in this study and a  
11  
12 358 detailed explanation of the phylogenetic analysis, refer to Appendices S1 and S2,  
13  
14 respectively.  
15

360

### Environmental variables

21 362 We analyzed six variables describing contemporary and historical temperature and  
22  
23 precipitation known or hypothesized to affect the structure of pollination networks. Four of the  
24  
25 364 six variables describe contemporary climatic conditions: *mean annual temperature* (MAT, °C),  
26  
27 *temperature seasonality* (TS,  $\text{standard deviation} \times 100$ ), *mean annual precipitation* (MAP,  
28  
29 mm) and *precipitation seasonality* (PS,  $\text{coefficient of variation}$ ). We obtained these  
30  
31 366 measurements from the WorldClim data set with spatial resolutions of 1 x 1 km  
32  
33 (<http://www.worldclim.org>; see also Hijmans *et al.*, 2005). The historical climate variables, the  
34  
35 368 *velocity of temperature change* and the *velocity of precipitation change*, reflect the speed of  
36  
37 temperature and precipitation change between the Last Glacial Maximum (LGM) and pre-  
38  
39 370 industrial times (VT, m/yr; VP, m/yr), following the definition of Loarie *et al.* (2009). Projections  
40  
41 of the global climate during the LGM and pre-industry were generated by Hadley Centre  
42  
43 372 Coupled Model Version 3 (HadCM3) with a resolution of 3.75 x 2.5 arc degrees (Singarayer  
44  
45 & Valdes, 2010), and were downscaled to 0.1 x 0.1 arc degrees. For each study site, climate  
46  
47 374 estimates were calculated as the average values of all 1 x 1 km grid cells (0.1 x 0.1 arc  
48  
49 degrees for paleoclimate data) within a concentric distance of 10 km from the sampling  
50  
51 376 location. Additionally, we scored whether a network was from the mainland (0) or an island  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2 378 (1).  
3  
4  
5

6 380 *Macroecological models*  
7

8 We used a multi-model approach based on information theory as outlined in Diniz-Filho *et al.*  
9  
10 382 (2008) to simultaneously evaluate the relationships between species richness, phylogenetic  
11 signal and environment and hummingbird-plant network structure. First, for each network  
12  
13 384 metric, we calculated full ordinary least squares (OLS) regression models which included the  
14 following ten predictor variables: 1) species richness (network size), 2) the phylogenetic  
15  
16 386 signal in the interaction pattern of plants' ( $d_{\text{plants}}$ ) and 3) hummingbirds' ( $d_{\text{hummingbirds}}$ ), 4)  
17 annual average temperature, 5) temperature seasonality, 6) annual average precipitation, 7)  
18  
19 388 precipitation seasonality, 8) temperature-change velocity, 9) precipitation-change velocity and  
20  
21 10) insularity. Visually, OLS regression models minimize the distance between observed data  
22  
23 390 points and those predicted by a linear approximation of the data. Second, for each network  
24 metric, we used the Akaike Information Criterion ( $AIC_C$ ) aiming at identifying minimum  
25  
26 392 adequate models (MAMs) among all possible model combinations of our ten predictor  
27 variables. MAMs were defined as models exhibiting a differences in  $AIC_C$  of at least two  
28  
29 394 points lower than other models (i.e. a  $\Delta AIC_C < 2$ ; Burnham & Anderson, 2002; Diniz-Filho *et al.*, 2008). As no single MAM was identified, often around 10 models had  $\Delta AIC_C < 2$ , we  
30  
31 396 instead used a multi-model approach. Specifically, instead of calculating regression  
32 coefficients in a single best model (MAM), we calculated the overall importance of each  
33  
34 398 model ( $w_i$ ) as the relative likelihood of any given model  $i$  over the sum of the likelihoods of the  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48

$$w_i = \frac{\exp\left(\frac{-1}{2}\Delta_i\right)}{\sum_{r=1}^R \exp\left(\frac{-1}{2}\Delta_r\right)}$$

49 entire dataset of models, such that , being  $\Delta_i$  the differences in AIC  
50  
51  
52 400 between the set of  $R$  models, so that the relative strength of each model depends on the  
53  
54  
55  
56  
57  
58  
59  
60  
60

1  
2 402 importance ( $\sum w_i$ ) of each variable for ordinary least squares OLS regression for an averaged  
3  
4 model based on weighted  $w_i$  (Burnham & Anderson, 2002; Diniz-Filho *et al.*, 2008), for which  
5  
6 404 we adopted an importance cut-off value of  $\geq 0.750$ . The standardized regression coefficients  
7  
8 were also reported for the OLS regression model including all ten predictor variables (“full”  
9  
10  
11 406 model). For each network metric, we used partial regressions to separate the total, unique  
12  
13 and shared variation explained by species richness, phylogenetic signal and environmental  
14  
15 408 factors in the “full” models.  
16

17  
18 Previous studies suggest that the structure of mainland and island hummingbird-plant  
19  
20 410 and pollination networks may differ (Olesen *et al.*, 2002; Dalsgaard *et al.*, 2009, 2013), and  
21  
22 the colonization of the Caribbean by hummingbirds has been considerably more recent than  
23  
24  
25 412 mainland America (c. 5 million years ago versus 12-22 million years ago for North and South  
26  
27 America, respectively). Consequently insular hummingbirds have had less time for  
28  
29 414 specialization and coevolution with their nectar plants than their mainland counterparts  
30  
31 (Bleiweiss, 1998; McGuire *et al.*, 2014). Together with the higher levels of generalization and  
32  
33  
34 416 less modularity predicted for insular pollinator communities (Olesen *et al.*, 2002; Dalsgaard *et*  
35  
36 *al.*, 2013), and a high level of periodic disturbances characteristic of Caribbean communities  
37  
38 418 which may hinder high levels of specialization on islands (Graves & Olsen, 1987; Rivera-  
39  
40 Marchand & Ackerman, 2006), the relationship between species richness, phylogenetic  
41  
42 420 signal and environmental factors and network structure may differ between mainland and  
43  
44 insular communities. For instance, the influence of historical climate change may be weaker  
45  
46  
47 422 on islands than on the mainland (Dalsgaard *et al.*, 2013, 2014). Our dataset contains too few  
48  
49 island networks (n=9) to allow for a separate analysis for insular networks, hence, we  
50  
51 424 explored putative differences in mainland and island networks by analyzing a subset of the  
52  
53 dataset composed exclusively by networks from the mainland (Mainland, n=45 networks) and  
54  
55  
56 426 comparing these results to those of the entire dataset (Global dataset, which includes both  
57  
58  
59  
60

1 mainland and insular communities, n=54 networks).

2  
3  
4  
5 428 For all macroecological models, we log<sub>10</sub>-transformed species richness, temperature  
6 seasonality, temperature velocity, and precipitation velocity; we squared temperature, and  
7  
8  
9 430 square-root transformed precipitation. All other variables were left untransformed. In all  
10 spatial models we tested whether significant positive spatial autocorrelation remained in  
11  
12  
13 432 model residuals of the “full” models (i.e., whether p<0.05 in all distance classes, tested using  
14  
15  
16 10 equally-spaced distance classes and applying a permutation test with 10,000 iterations).  
17  
18 434 As no positive spatial autocorrelation was observed we did not build more sophisticated  
19  
20  
21 spatial models. All regression analyses were conducted using the software Spatial Analysis in  
22  
23 436 Macroecology, SAM 4.0 (Rangel *et al.*, 2010).

24  
25  
26 To better support our findings, we performed a number of complementary  
27  
28 438 macroecological models which included sampling effort as a predictor variable (Appendix  
29  
30 S3), observed (uncorrected) metrics of network structure instead of null model corrected  
31  
32 440 metrics (Appendix S4), and climate anomaly as a measure of historical climate stability  
33  
34 instead of climate change velocity (Appendix S5).  
35  
36

37 442

## 38 39 RESULTS

### 40 41 444 *Phylogenetic signal on species interaction patterns*

42  
43 Values for the independent phylogenetic signals of plants ( $d_{\text{plants}}$ ) and hummingbirds  
44  
45 446 ( $d_{\text{hummingbirds}}$ ) were low but above zero in 85 and 65% of the networks, respectively,  
46  
47 suggesting a significant relationship between phylogenetic signal and species interaction  
48  
49 patterns. The confidence intervals (CI) of  $d_{\text{hummingbirds}}$  overlapped with zero in a higher number  
50  
51 448 of networks than did CIs of  $d_{\text{plants}}$  (82 and 57%; Table in Appendix S2), indicating that the  
52  
53  
54 450 significant association between phylogeny and interaction patterns was exhibited mostly by  
55  
56  
57 plant species.  
58  
59  
60

1  
2 452 We examined the overall association between phylogeny and the structure of bipartite  
3  
4 interaction networks by comparing the mean squared error (MSE) of the model fit among  
5  
6 454 models adjusted after observed phylogenetic signal ( $MSE_{data}$ ), models which assumed no  
7  
8 phylogenetic signal ( $MSE_{star}$ ), and models which assume a maximum signal ( $MSE_{Brownian}$ ).  
9  
10 456 Results showed that most networks exhibited a significant phylogenetic signal (in 52  
11  
12 networks  $MSE_{data} < MSE_{star}$ , in 1 networks  $MSE_{data} = MSE_{star}$ , and in 1  $MSE_{data} > MSE_{star}$ ; in all  
13  
14 458 cases  $MSE_{Brownian}$  had clearly the highest values; Table in Appendix S2), e.g. in most  
15  
16 communities the model which fitted best to the data (the one having the lowest error) was  
17  
18 achieved when adjusting the model with observed phylogenetic signals of plants and  
19  
20 460 hummingbirds ( $MSE_{data}$ ). Nevertheless, most differences between  $MSE_{data}$  and  $MSE_{star}$  were  
21  
22 small, and much smaller than between  $MSE_{data}$  and  $MSE_{Brownian}$ , indicating that in general, the  
23  
24 462 association between phylogenetic signal and species interaction pattern was weak.  
25  
26  
27

464

### 30 *Macroecological models*

31  
32  
33 466 Full OLS models on corrected network metrics were able to account for 54%-62% of the  
34  
35 observed variation in network metrics across the sampled communities (Table 1, Table S3,  
36  
37 468 Fig. 3). Species richness and environmental factors showed the strongest association with  
38  
39 network structure, although with varying strengths across the "Global" and "Only Mainland"  
40  
41 470 datasets and network metrics (Table 1, Table S3, Figure S1). On the contrary, the  
42  
43 associations between phylogenetic signal and network structure remained constant,  
44  
45 exhibiting low regression coefficients in both metrics and datasets (Table 1, Table S3).  
46  
47 472 Collectively, complementary specialization exhibited a stronger association with current  
48  
49 temperature and historical temperature stability, and to a lesser extent with species richness;  
50  
51 474 whereas species richness had the strongest association with modularity (Fig. 3; Table S3).  
52  
53

54  
55 476 When the ten predictor variables included in the full OLS models were examined in  
56  
57  
58  
59  
60

1 detail, species richness and hummingbird phylogenetic signal were the only variables  
2  
3  
4 478 showing an important, spatially consistent and widespread association to network structure,  
5  
6 emerging as highly important for both network metrics and datasets. Species richness  
7  
8 480 showed a strong positive relationship with complementary specialization and modularity  
9  
10 (Table 1). On the contrary, a higher phylogenetic signal among hummingbirds was related  
11  
12 482 with a lower complementary specialization and level of modularity. In other words, despite the  
13  
14 association between phylogenetic signal and species' interaction pattern was weak and the  
15  
16 weighted regression coefficients of  $d_{\text{hummingbirds}}$  were low, complementary specialization and  
17 484 modularity consistently increased when closely-related hummingbirds visited distinct arrays  
18  
19 of plant species (Table 1, Table S3, Fig. 3).  
20  
21  
22 486

23  
24 Complementary specialization was also highly associated with the environmental  
25  
26 488 conditions, as the total variation explained by environmental factors was consistently higher  
27  
28 than the variation attributed to species richness and phylogenetic signal in both the "Global"  
29  
30 490 and "Only Mainland" datasets, and approximated to the variation explained by the full models  
31  
32 (Table S3, Fig. 3). The amount of unique variation (i.e. variation explained exclusively by the  
33  
34 predictor-type) explained by environmental factors was similar to the unique variation  
35 492 explained by species richness on the Global dataset (Table S3, Fig. 3). However, when  
36  
37 examining only mainland networks, the unique variation explained by environmental factors  
38  
39 494 was two times greater than the unique variation explained by the other two predictor-types  
40  
41 was two times greater than the unique variation explained by the other two predictor-types  
42  
43 (Table S3, Fig. 3). Among the seven environmental variables examined, only current  
44 496 temperature and historical temperature stability showed a strong relationship with  
45  
46 complementary specialization, with increasing complementary specialization in communities  
47  
48 498 with current warmer temperatures and with a higher historical temperature stability (Table 1).  
49  
50

51  
52  
53 500 Quantitative bipartite modularity showed the strongest association with species  
54  
55 richness, particularly in the global dataset, with environmental conditions having a similar  
56  
57  
58  
59  
60

1  
2 502 strength association among mainland communities (Table S3, Fig. 3).

3  
4 Shared variability among species richness and phylogenetic signal (i.e. variability  
5  
6 504 explained by these two predictor types) was non-existent, i.e. richness and phylogenetic  
7  
8 signal were distinctly associated to specialization and modularity. Shared variability between  
9  
10 506 environmental factors and species richness or phylogenetic signal was low (Table S3).

11  
12 The remaining five environmental variables (insularity, temperature seasonality, mean  
13  
14 508 annual precipitation, precipitation seasonality, and precipitation change velocity) as well as  
15  
16 phylogenetic signal in the interaction pattern of plants, showed no important associations with  
17  
18 510 null model corrected network structure (Table 1).

## 21 22 23 24 512 **DISCUSSION**

25  
26 As previous macroecological studies of interaction networks, we found complementary  
27  
28 514 specialization and modularity to vary along with environmental conditions and species  
29  
30 richness. Additionally, for the first time, we report an association between phylogenetic signal  
31  
32 516 and network structure at macroecological scale. Specifically, species richness and  
33  
34 phylogenetic signal in hummingbird interaction patterns were the two predictor variables that  
35  
36 518 associated most consistently to network structure, with an association between  
37  
38 complementary specialization and current and historical temperature conditions limited to  
39  
40 520 mainland networks.

41  
42 Hummingbird phylogenetic signal had a constant association with both complementary  
43  
44 522 specialization and modularity, and on both the Global and Only Mainland datasets, although  
45  
46 the variability associated with phylogenetic signal was the lowest of all predictor types. A  
47  
48 524 weak but significant relationship between phylogenetic signal and species' interaction pattern  
49  
50 within ecological networks has also been reported by previous studies on food webs, host-  
51  
52 526 parasite networks or plant-pollinator networks (Ives & Godfray, 2006; Vázquez *et al.*, 2009b;  
53  
54  
55  
56  
57  
58  
59  
60

1 Krasnov *et al.*, 2012; Rafferty & Ives, 2013). These studies also identified asymmetries in the  
2  
3  
4 528 phylogenetic signal between trophic levels, with the association between phylogenetic  
5  
6 relatedness and species interaction pattern being stronger at lower trophic levels, i.e. plants  
7  
8  
9 530 in our system (but see Rezende *et al.*, 2007). According to these studies, such asymmetries  
10  
11 may stem from a differential relationship between phylogeny and the interaction pattern of  
12  
13 532 each trophic group, as species from the higher trophic group (“consumer” species) are more  
14  
15 likely to adjust their feeding behavior according to local conditions. Our results corroborate  
16  
17 534 this hypothesis: hummingbird phylogenetic signal showed a weaker relationship with their  
18  
19 interaction pattern than plant species ( $d_{\text{hummingbirds}}$  tends to be lower than  $d_{\text{plants}}$ , and CI  
20  
21 536  $d_{\text{hummingbirds}}$  overlap with zero in more networks). Nevertheless, and unlike for plants which  
22  
23 showed no association to the spatial variation in network structure, hummingbird phylogenetic  
24  
25  
26 538 signal was associated to network structure, with higher levels of complementary  
27  
28 specialization and modularity consistently achieved when closely-related hummingbird  
29  
30 540 species visited distinct sets of flowering plant species. This suggests that resource  
31  
32 partitioning and inter-specific competition among closely-related hummingbirds might play an  
33  
34  
35 542 important role in structuring interactions in hummingbird-plant networks. Indeed, inter-specific  
36  
37 competition is known to be strong among hummingbirds, and has been noted as a potential  
38  
39 544 driver of patterns of hummingbird biodiversity (Brown & Bowers, 1985; Bleiweiss, 1998;  
40  
41 Cotton, 1998).

42  
43  
44 546 Species richness had the strongest association with both metrics and in both the  
45  
46 Global and Only Mainland datasets except for complementary specialization in the mainland.  
47  
48  
49 548 Higher species richness may result in a higher niche availability, thus providing ample  
50  
51 opportunities for biotic specialization. At the same time, higher species richness may increase  
52  
53 550 inter-specific competition, which may explain why high species richness lead to higher levels  
54  
55 of complementary specialization and modularity (Rezende *et al.*, 2009; Dalsgaard *et al.*,  
56  
57  
58  
59  
60



1  
2 552 2011; Krasnov *et al.*, 2012; Junker *et al.*, 2013). Moreover, a higher plant richness may also  
3  
4 translate into a temporally-stable availability of floral resources (or the establishment of a  
5  
6 554 constant minimum local flower supply), [enabling a locally constant hummingbird population](#)  
7  
8 [and, hence, potential for biotic specialization](#) (Montgomerie & Gass, 1981; Stiles, 1985;  
9  
10 556 [Araujo & Sazima, 2003; Cotton, 2007; Abrahamczyk \*et al.\*, 2011](#)).

11  
12 In the mainland, higher levels of [complementary specialization](#) were also found in  
13  
14  
15 558 warmer areas and in areas with higher historical temperature stability. The former may again  
16  
17 be linked to higher inter-specific competition in warmer areas, and particularly in mainland  
18  
19 560 settings, where hummingbird species and phylogenetic richness is highest (Bleiweiss, 1998;  
20  
21 McGuire *et al.*, 2014). This relationship might be stronger on the mainland as insular  
22  
23 562 pollinator faunas are typically depauperate (Olesen *et al.*, 2002). Moreover, in historically  
24  
25 stable climates, hummingbirds have been able to establish long-term associations with  
26  
27 564 plants, which may lead to higher levels of local adaptation, interaction specificity and  
28  
29 specialization (see Dalsgaard *et al.*, 2011 for a study on a subset of the networks from [the](#)  
30  
31 [dataset used here](#)). Such long associations may lead to higher [complementary specialization](#)  
32  
33 in areas with low temperature-change velocity, such as montane habitats and tropical  
34  
35 568 environments, leaving more generalized interaction patterns to areas with low topography  
36  
37 and especially at higher latitudes where changes in [temperature velocity have been greatest](#)  
38  
39 [\(Sandel \*et al.\*, 2011\)](#). Moreover, a higher [complementary specialization](#) in areas of high  
40  
41 570 historical temperature stability is consistent with general hummingbird historical biogeography  
42  
43 and speciation patterns, as the core area of hummingbird speciation and diversity occurs in  
44  
45 Andean highland and in tropical lowland South-America (Brown & Bowers, 1985; Bleiweiss,  
46  
47 574 1998; McGuire *et al.*, 2014). [We note that complementary macroecological analysis](#)  
48  
49 [presented in Appendices S3-S5 generally confirmed the importance of species richness and](#)  
50  
51 [hummingbird phylogenetic signal, and, additionally, showed an association between](#)  
52  
53  
54  
55 576  
56  
57  
58  
59  
60

1 observed (uncorrected) complementary specialization/modularity and insularity and  
2  
3  
4 578 precipitation, with areas of high precipitation and mainland communities showing higher  
5  
6 observed complementary specialization and modularity. Precipitation has previously been  
7  
8 580 shown to influence both phenotypical and network specialization of hummingbird-plant  
9  
10 interactions and of entire pollination communities (Cruden, 1972; Dalsgaard *et al.*, 2009;  
11  
12 582 Martín González *et al.*, 2009). The fact that precipitation and insularity only associated with  
13  
14 observed complementary specialization and observed modularity and not with null model  
15  
16  
17 584 corrected values of these network metrics, may indicate that insularity and precipitation do  
18  
19 not directly influence the interaction pattern of species, but rather associate indirectly with  
20  
21 network structure through species richness and/or interaction probability distributions.  
22  
23  
24  
25

## 26 588 **CONCLUSIONS**

27  
28 Characterizing potential ecological, evolutionary and environmental mechanisms **associated**  
29  
30 590 **with** the structure of ecological communities is a critical first step towards understanding the  
31  
32 determinants of community assembly and how climate-change may affect biodiversity  
33  
34 592 (Woodward *et al.*, 2010; Schleuning *et al.*, 2014). By examining null model corrected network  
35  
36 metrics, we are actively investigating how much observed communities depart from random  
37  
38 ones with an equivalent set of species and interactions. Hence, we were able to investigate  
39  
40 594 not only whether a community presents a structure **which differs significantly from random**,  
41  
42  
43 but also to associate the difference in complementary specialization and modularity to  
44  
45 596 species richness, evolutionary history, **and environmental conditions**.  
46  
47

48  
49 598 We have shown that **complementary specialization and modularity** in hummingbird-  
50  
51 plant **networks associate to** species richness, hummingbird phylogenetic signal and  
52  
53 600 environmental factors acting at varying spatio-temporal scales and in different aspects of  
54  
55 network structure. Notably, species richness and hummingbird phylogenetic signal showed a  
56  
57  
58  
59  
60

1  
2 602 consistent association with network structure, with a more restricted but still important role of  
3  
4 contemporary temperature and historical temperature stability, which were only important  
5  
6 604 among mainland communities. These results are markedly different to the ones for avian  
7  
8 seed-dispersal networks for which species' phylogeny and historical climatic stability were  
9  
10  
11 606 unrelated to modularity and complementary specialization (Schleuning *et al.*, 2012, 2014;  
12  
13 Sebastián González *et al.*, 2015). Such difference suggests that hummingbirds and flowers  
14  
15 608 engage in tighter co-evolutionary associations than frugivore birds and their plants, and  
16  
17 hence, historical and evolutionary factors may have a stronger role in hummingbird-plant  
18  
19 610 assemblages than for frugivore birds-plant assemblages.  
20  
21  
22  
23

## 24 612 **ACKNOWLEDGMENTS**

25  
26 We thank Carsten Dormann and Matthew Helmus for invaluable help regarding the use of the  
27  
28 614 software bipartite and the interpretation of the phylogeny results, respectively. AMMG was  
29  
30 supported by the Spanish Ministry of Education; BD by the Carlsberg Foundation; ACB by the  
31  
32 616 OTICON Fonden; PAC by the David Lack studentship from the British Ornithologists' Union  
33  
34 and Wolfson College, University of Oxford; CL by the CACyPI-Uatx-2013; MAM by the  
35  
36 618 Universidad Estatal a distancia (UNED), Costa Rica, and the Biodiversity and Climate  
37  
38 Research Centre (BIK-F), Frankfurt, Germany; LRL by FAEP and Unicamp; TTI by Frimodt-  
39  
40 620 Heineke Foundation, the Knud Højgaard Foundation and Faculty of Natural Sciences  
41  
42 University of Aarhus. FAPESP supported ACA and MS; FAPEMIG supported PKM and PEO;  
43  
44 622 CNPq supported ACA, GMO, LCR, LRL, MS, PKM, PEO, IGV and JV-B; CAPES supported  
45  
46 FPA, FMGLC, LRL, PKM and LCR; and FUNDECT supported ACA. SW thanks the National  
47  
48 624 Institute of Natural Resources (INRENA) for research permits in Perú and LRL the Ministerio  
49  
50 del Medio Ambiente de Colombia for permission the entrance to Chiribiquete. AMMG, BD,  
51  
52 626 DN-B, PKM, ZW and CR thank the Danish National Research Foundation for its support of  
53  
54  
55  
56  
57  
58  
59  
60

1 the Center for Macroecology, Evolution and Climate.  
2  
3

4 628

## 5 REFERENCES

6  
7 630 Abrahamczyk, S., Kluge, J., Gareca, Y., Reichle, S & Kessler, M. (2011) The influence of

8 climatic seasonality on the diversity of different tropical pollinator groups. *PLoS ONE*, **6**,

9  
10  
11 632 e27115.

12  
13  
14 Araujo, A.C. & Sazima, M. (2003) The assemblage of flowers visited by hummingbirds in the

15  
16 634 "capões" of Southern Pantanal, Mato Grosso do Sul, Brazil. *Flora*, **198**, 427–435.

17  
18 Bleiweiss, R. (1998) Origin of hummingbird faunas. *Biological Journal of the Linnean*

19  
20  
21 636 *Society*, **6**, 77–97.

22  
23 Blüthgen, N., Menzel, F. & Blüthgen, N. (2006) Measuring specialization in species

24  
25 638 interaction networks. *BMC Ecology*, 6:9.

26  
27 [Blüthgen, N. \(2010\). Why network analysis is often disconnected from community ecology: a](#)

28  
29 640 [critique and an ecologist's guide. \*Basic and Applied Ecology\*, \*\*11\*\*\(3\), 185-195.](#)

30  
31  
32 Brown, J.H. & Bowers, M.A. (1985) Community organization in hummingbirds: relationships

33  
34 642 between morphology and ecology. *The Auk*, **102**, 251–269.

35  
36 Burnham, K.P. & Anderson, D.R. (2002) *Model Selection and Multimodel Inference: A*

37  
38 644 *Practical Information-Theoretic Approach* (2nd ed.), Springer-Verlag, New York.

39  
40 Cotton, P.A. (1998) Coevolution in an Amazonian hummingbird-plant community. *Ibis*, **140**,

41  
42  
43 646 639-646.

44  
45 Cotton, P.A. (2007) Seasonal resource tracking by Amazonian hummingbirds. *Ibis*, **149**,

46  
47 648 135–142.

48  
49 Cruden, R.W. (1972) Pollinators in high-elevation ecosystems: relative effectiveness of birds

50  
51  
52 650 and bees. *Science*, **176**, 1439–1440.

53  
54 Dalsgaard, B., Martín González, A.M., Olesen, J.M., Ollerton, J., Timmermann, A., Andersen,

55  
56 652 L. H. & Tossas, A. (2009) Plant–hummingbird interactions in the West Indies: floral

57  
58  
59  
60

- 1 specialisation gradients associated with environment and hummingbird size. *Oecologia*,  
2  
3  
4 654 **159**, 757–766.
- 5  
6 Dalsgaard, B., Magård, E., Fjeldså, J., Martín González, A.M., Rahbek, C., Olesen, J.M., et  
7  
8 656 al. (2011) Specialization in plant-hummingbird networks is associated with species  
9  
10 richness, contemporary precipitation and quaternary climate-change velocity. *PloS ONE*,  
11  
12 658 **6**, e25891.
- 13  
14 Dalsgaard, B., Trøjelsgaard, K., Martín González, A.M., Nogués-Bravo, D., Ollerton, J.,  
15  
16 660 Petanidou, T., et al. (2013) Historical climate-change influences modularity and  
17  
18 nestedness of pollination networks. *Ecography*, **36**, 1331–1340.
- 19  
20  
21 662 Dalsgaard, B., Carstensen, D.W., Fjeldså, J. Maruyama, P.K., Rahbek, C., Sandel, B.,  
22  
23 Sonne, J., Svenning, J.C., Wang, Z. & Sutherland, W.J. (2014) Determinants of bird  
24  
25 664 species richness, endemism, and island network roles in Wallacea and the West Indies:  
26  
27 is geography sufficient or does current and historical climate matter? *Ecology and*  
28  
29 666 *Evolution*, **4**, 4019–4031.
- 30  
31  
32 Diniz-Filho, J.A.F., Rangel, T.F. L. & Bini, L.M. (2008) Model selection and information theory  
33  
34 668 in geographical ecology. *Global Ecology and Biogeography*, **17**, 479–488.
- 35  
36  
37 Dormann, C.F. & Strauss, R. (2014) A method for detecting modules in quantitative bipartite  
38  
39 670 networks: the QuanBiMo algorithm. *Methods in Ecology and Evolution*, **5**, 90–98.
- 40  
41  
42 Dormann, C.F., Fründ, J., Blüthgen, N. & Gruber, B. (2009) Indices, graphs and null models:  
43  
44 672 analyzing bipartite ecological networks. *Open Ecology Journal*, **2**, 7–24.
- 45  
46  
47 Graves, G.R. & Olsen, S.L. (1987) *Chlorostilbon bracei lawrence*, an extinct species of  
48  
49 674 hummingbird from New Providence Island, Bahamas. *Auk*, **104**, 296–302.
- 50  
51  
52 Hijmans, R.J., Cameron, S.E., Parra, J.L., Jones, P.G. & Jarvis, A. (2005) Very high  
53  
54 676 resolution interpolated climate surfaces for global land areas. *International Journal of*  
55  
56 *Climatology*, **25**, 1965–1978.  
57  
58  
59  
60

- 1  
2 678 Ives, A.R. & Godfray, H.C.J. (2006) Phylogenetic analysis of trophic associations. *American*  
3  
4 *Naturalist*, **168**, E1–E14.
- 5  
6 680 Junker, R.R., Blüthgen, N., Brehm, T., Binkenstein, J., Paulus, J., Martin Schaefer, H., et al.  
7  
8 (2013) Specialization on traits as basis for the niche-breadth of flower visitors and as  
9  
10 682 structuring mechanism of ecological networks. *Functional Ecology*, **27**, 329–341.
- 11  
12  
13  
14  
15 684 Kembel, S.W., Cowan, P.D., Helmus, M.R., Cornwell, W.K., Morlon, H., Ackerly, D.D., et al.  
16  
17 (2010) Picante: R tools for integrating phylogenies and ecology. *Bioinformatics*, **26**,  
18  
19 14631464.
- 20 686 Krasnov, B. R., Fortuna, M. A., Mouillot, D., Khokhlova, I. S., Shenbrot, G. I. & Poulin, R.  
21  
22 (2012) Phylogenetic signal in module composition and species connectivity in  
23  
24 688 compartmentalized host-parasite networks. *American Naturalist*, **179**, 501–511.
- 25  
26  
27  
28 690 Loarie, S.R., Duffy, P.B., Hamilton, H., Asner, G.P., Field, C.B. & Ackerly, D.D. (2009) The  
29  
30 velocity of climate change. *Nature*, **462**, 1052–1055.
- 31  
32  
33  
34  
35  
36  
37 694 Martín González, A.M., Dalsgaard, B., Ollerton, J., Timmermann, A., Olesen, J.M., Andersen,  
38  
39 [L. & Tossas, A. \(2009\) Effects of climate on pollination networks in the West Indies.](#)  
40  
41 [Journal of Tropical Ecology](#), **25**, 493–506.
- 42  
43  
44  
45  
46 696 Martín González, A.M., Allesina, S., Rodrigo, A., & Bosch, J. (2012). Drivers of  
47  
48 [compartmentalization in a Mediterranean pollination network.](#) *Oikos*, **121**(12), 2001–2013.
- 49  
50  
51  
52  
53  
54  
55 698 Maruyama, P.K., Vizentin-Bugoni, J., Oliveira, G.M., Oliveira, P.E. & Dalsgaard, B. (2014)  
56  
57 Morphological and spatio-temporal mismatches shape a Neotropical savanna plant-  
58  
59 hummingbird network. *Biotropica*, **46**, 740–747.
- 60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000

- 1 tropical and temperate communities. *Oecologia*, **50**, 162–165.
- 2
- 3
- 4 704 Olesen, J.M., Eskildsen, L.I. & Venkatasamy, S. (2002) Invasion of pollination networks on
- 5 oceanic islands: importance of invader complexes and endemic super generalists.
- 6
- 7
- 8 706 *Diversity and Distributions*, **8**, 181–192.
- 9
- 10 Paradis, E., Claude, J., & Strimmer, K. (2004) APE: analyses of phylogenetics and evolution
- 11
- 12 708 in R language. *Bioinformatics*, **20**, 289–290.
- 13
- 14 Rafferty, N.E. & Ives, A.R. (2013) Phylogenetic trait-based analyses of ecological networks.
- 15
- 16 710 *Ecology*, **94**, 2321–2333.
- 17
- 18
- 19 Rangel, T.F., Diniz-Filho, J.A.F. & Bini, L.M. (2010) SAM: a comprehensive application for
- 20
- 21 712 spatial analysis in macroecology. *Ecography*, **33**, 46–50.
- 22
- 23
- 24 Rezende, E.L., Albert, E.M., Fortuna, M.A. & Bascompte, J. (2009) Compartments in a
- 25
- 26 714 marine food web associated with phylogeny, body mass, and habitat structure. *Ecology*
- 27
- 28 *Letters*, **12**, 779–788.
- 29
- 30 716 Rezende, E.L., Jordano, P. & Bascompte, J. (2007) Effects of phenotypic complementarity
- 31
- 32 and phylogeny on the nested structure of mutualistic networks. *Oikos*, **116**, 1919–1929.
- 33
- 34 718 Rivera-Marchand, B. & Ackerman, J.D. (2006) Bat pollination breakdown in the Caribbean
- 35
- 36 columnar cactus *Pilosocereus royenii*. *Biotropica*, **38**, 635–642.
- 37
- 38
- 39 720 Sandel, B., Arge, L., Dalsgaard, B., Davies, R.G., Gaston, K.J., Sutherland, W.J. & Svenning,
- 40
- 41 J.-C. (2011) The influence of late Quaternary climate-change velocity on species
- 42
- 43 722 endemism. *Science*, **334**, 660–664.
- 44
- 45
- 46 Schleuning, M., Fründ, J., Klein, A. M., Abrahamczyk, S., Alarcón, R., Albrecht, M., et al.
- 47
- 48 724 (2012) Specialization of mutualistic interaction networks decreases toward tropical
- 49
- 50 latitudes. *Current Biology*, **22**, 1925–1931.
- 51
- 52
- 53 726 Schleuning, M., Ingmann, L., Strauss, R., Fritz, S.A., Dalsgaard, B., Matthias Dehling, D., et
- 54
- 55 al. (2014) Ecological, historical and evolutionary determinants of modularity in weighted
- 56
- 57
- 58
- 59
- 60

- 1  
2 728 seed-dispersal networks. *Ecology Letters*, **17**, 454–463.
- 3  
4 [Sebastián González , E., Dalsgaard, B., Sandel, B. & Guimarães, P. \(2015\). Macroecological](#)  
5  
6 730 [trends in nestedness and modularity of seed-dispersal networks: human impact matters.](#)  
7  
8 [Global Ecology and Biogeography](#), **24**, 293–303.
- 9  
10  
11 732 Singarayer, J.S., & Valdes, P. J. (2010) High-latitude climate sensitivity to ice-sheet forcing  
12  
13 over the last 120 kyr. *Quaternary Science Reviews*, **29**, 43–55.
- 14  
15 734 Stiles, G.F. (1985) Seasonal patterns and coevolution in the hummingbird-flower community  
16  
17 of a Costa Rican subtropical forest. *Ornithological Monographs*, **36**, 757–785.
- 18  
19 736 Stouffer, D.B. & Bascompte, J. (2011) Compartmentalization increases food-web  
20  
21 persistence. *Proceedings of the National Academy of Science USA*, **108**, 3648–3652.
- 22  
23  
24 738 [Thébault, E., & Fontaine, C. \(2010\). Stability of ecological communities and the architecture](#)  
25  
26 [of mutualistic and trophic networks. Science](#), **329**(5993), 853–856.
- 27  
28 740 Trøjelsgaard, K. & Olesen, J.M. (2013) Macroecology of pollination networks. *Global*  
29  
30 *Ecology Biogeography*, **22**, 149–162.
- 31  
32  
33 742 Vázquez, D.P., Melian, C.J., Williams, N.M., Blüthgen, N., Krasnov, B.R. & Poulin, R. (2007)  
34  
35 Species abundance and asymmetric interaction strength in ecological networks. *Oikos*,  
36  
37 744 **116**, 1120–1127.
- 38  
39 Vázquez, D.P., Blüthgen, N., Cagnolo, L. & Chacoff, N.P. (2009a) Uniting pattern and  
40  
41 746 process in plant–animal mutualistic networks: a review. *Annals of Botany*, **103**, 1445–  
42  
43 1457.
- 44  
45 748 Vázquez, D.P., Chacoff, N.P. & Cagnolo, L. (2009b) Evaluating multiple determinants of the  
46  
47 structure of plant-animal mutualistic networks. *Ecology*, **90**, 2039–2046.
- 48  
49  
50 750 Webb, C.O., Ackerly, D.D., McPeck, M.A. & Donoghue, M.J. (2002) Phylogenetics and  
51  
52 community ecology. *Annual Review of Ecology and Systematics*, **33**, 475–505.
- 53  
54  
55 752 Williams, R.J. (2010) *Network3D Software*. Microsoft Research, Cambridge, UK.
- 56  
57  
58  
59  
60



1 Woodward, G., Benstead, J.P., Beveridge, O.S., Blanchard, J., Brey, T., Brown, L., et al.

2  
3  
4 754 (2010) Ecological networks in a changing climate. *Advanced Ecological Research*, **42**,  
5  
6 71–138.

7  
8 756 Yoon, I., Williams, R.J., Levine, E., Yoon, S., Dunne, J.A. & Martinez, N.D. (2004) *Webs on*  
9  
10 *the Web (WoW): 3D visualization of ecological networks on the WWW for collaborative*  
11  
12 *research and education*. Proceedings of the IS&T/SPIE Symposium on Electronic  
13 758  
14 Imaging, Visualization and Data Analysis, 5295, 124-132.

15  
16  
17 760

18  
19 **Biosketch:** This work results from an ongoing collaboration among numerous researchers  
20  
21 762 from different scientific backgrounds and institutions, who have gathered an extensive  
22  
23 dataset of hummingbird plant interaction networks and complementary data on species traits,  
24  
25 phylogeny, climate, etc. Such multidisciplinary information is allowing us to study in depth  
26 764  
27 different aspects of hummingbird-plant interaction patterns at macroecological scales, and to  
28  
29 identify potential mechanisms responsible for them. Our ultimate goal is to describe general  
30 766  
31 macroecological patterns in biotic interactions using hummingbird-plant interactions as a  
32  
33  
34  
35 768 model system.

36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Table 1.** Relationship between species richness, phylogenetic signal and environment, and complementary specialization ( $H_2'$ ) and modularity (QuanBiMo) across the Americas (Global dataset) and the mainland, i.e. when excluding insular communities (Only Mainland). Complementary specialization measures the exclusiveness of the interactions in the community, whereas modularity quantifies whether species interact more frequently with subsets of available species within a community. For each network metric, the standardized regression coefficients are reported for ordinary least squares regression for a model including all variables ("Full") and for a model averaged across all possible models using Akaike Weights (AICc  $w_i$  - "Averaged"). For the averaged models, the relative importance of each predictor variable (" $\Sigma w_i$ ") is given by the Akaike Weights (AICc  $w_i$ ). Predictors with high importance are marked in bold. Phylogenetic signal among plants and hummingbirds,  $d_p$  and  $d_h$ , respectively; Mean annual temperature, MAT; Temperature seasonality, TS; Mean annual precipitation, MAP; Precipitation seasonality, PS; Velocity of temperature change since the last Glacial Maximum, MAT velocity; Velocity of precipitation change since the last Glacial Maximum, MAP velocity. The significance of Moran's  $I$  was tested with 10 distance classes and a permutation test with 10,000 iterations. In all models multicollinearity was not an issue, i.e. VIF  $\leq$  3.4, CN  $\leq$  4.0, and there was no positive spatial autocorrelation, i.e. Moran's  $I$  was non-significant.

		Species richness			Phylogenetic signal			Environmental factors						
		Model	AICc	R <sup>2</sup>	Network size	$d_p$	$d_h$	Insularity	MAT	TS	MAP	PS	MAT velocity	MAP velocity
Global dataset (54 networks)	$H_2'$	Full	-54.368	0.543	<b>0.461</b>	-0.087	<b>-0.303</b>	-0.222	0.284	0.176	0.045	-0.209	-0.475	0.190
		Averaged			<b>0.520</b>	-0.143	<b>-0.361</b>	-0.121	0.137	0.121	0.085	-0.217	-0.270	0.106
		$\Sigma w_i$			<b>0.999</b>	0.380	<b>0.962</b>	0.283	0.308	0.311	0.259	0.603	0.671	0.290
	QuanBiMo	Full	-99.867	0.638	<b>0.638</b>	-0.145	<b>-0.232</b>	-0.187	0.188	0.081	-0.003	-0.204	-0.132	-0.098
		Averaged			<b>0.652</b>	-0.183	<b>-0.283</b>	-0.137	0.080	0.058	0.097	-0.184	-0.098	-0.134
		$\Sigma w_i$			<b>1.000</b>	0.619	<b>0.915</b>	0.341	0.257	0.237	0.286	0.567	0.288	0.392
Only Mainland (45 networks)	$H_2'$	Full	-55.244	0.581	<b>0.326</b>	-0.042	<b>-0.332</b>	na	<b>0.477</b>	0.125	-0.085	-0.261	<b>-0.714</b>	0.307
		Averaged			<b>0.324</b>	-0.048	<b>-0.358</b>	na	<b>0.451</b>	0.110	-0.044	-0.241	<b>-0.653</b>	0.359
		$\Sigma w_i$			<b>0.807</b>	0.198	<b>0.919</b>	na	<b>0.854</b>	0.258	0.227	0.606	<b>0.931</b>	0.776
	QuanBiMo	Full	-76.885	0.567	<b>0.539</b>	-0.174	<b>-0.283</b>	na	0.289	0.046	-0.079	-0.248	-0.289	-0.093
		Averaged			<b>0.544</b>	-0.184	<b>-0.300</b>	na	0.231	0.021	0.096	-0.232	-0.237	-0.127
		$\Sigma w_i$			<b>0.999</b>	0.476	<b>0.805</b>	na	0.414	0.201	0.257	0.592	0.427	0.304


1  
2  
3  
4 776 **Figure 1.** Cartoon illustrations depicting two different communities and the characteristics  
5 and relationship between complementary specialization ( $H_2'$ ) and quantitative bipartite  
6 modularity (QuanBiMo). Both communities depict 700 interaction events between 10 plants  
7  
8 778 and 5 pollinators. Plants are labeled as numbers 1-10 in the matrix and as yellow nodes in  
9 the network representation, and pollinators by letters A-E in the matrix and as blue nodes in  
10  
11 780 the network. Differences in interaction frequencies between plants and pollinators are  
12 illustrated as numbers of interaction events in the matrix format and as varying line widths in  
13  
14 782 the network illustration. Complementary specialization measures the exclusiveness in  
15 species' interactions, whereas modularity quantifies whether species interact more frequently  
16 with subsets of available species within a community. Community A exhibits a moderate  
17  
18 784 complementary specialization, with different species-pairs exhibiting various degrees of  
19 complementary specialization. For instance, species-pair A-1 shows a high complementary  
20  
21 786 specialization, as hummingbird A visits exclusively and with high frequency (155 times) plant  
22 1, which in turn is only visited once by another hummingbird. On the other hand, the  
23  
24 788 complementary specialization or pair E-4 is lower than for the previous pair despite these  
25 species interact with the same frequency because hummingbird E also interacts with other  
26  
27 790 plant species, i.e. the interaction E-4 is less exclusive. Interactions in community B are also  
28 somewhat specialized (species favored interactions with few of the available partners), but  
29  
30 792 the exclusiveness of these interactions is lower than for community A. Both communities  
31 show very similar values for corrected modularity and can be divided into three distinct  
32  
33 794 modules, although the modules are composed of a different array of species. By having  
34 different degree of complementary specialization but similar values of modularity, these  
35  
36 796 communities show that although these metrics are positively related, measure  
37 complementary but different aspects of specialization. Network drawings were created using  
38  
39 800 Network3D and energized with the 3D Force-directed algorithm to enhance visualization of  
40 modularity patterns (Yoon et al. 2004, Williams 2010).  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 802 **Figure 2.** Map of the American continent showing the location of the 54 study sites and a  
5 number of example networks located along a species richness gradient. Localities with  
6  
7  
8 804 darker shades of green denote networks with a higher richness. Some networks have been  
9 slightly moved horizontally to maximize visualization (exact coordinates of the localities on  
10  
11  
12 806 Table S1). For each illustrated network, the reference number and a concise description of  
13 the vegetation type is given, along with a network drawing. For the network drawings, blue  
14  
15  
16 808 and yellow nodes depict hummingbird and plant species, respectively, while line width  
17 depicts log+1 frequency of interaction among species. Notice that species-rich networks in  
18  
19  
20 810 general present more complex structures, with networks 11, 21 and 50 exhibiting the lowest  
21 corrected complementary specialization; networks 50, 11 and 21 the lowest corrected  
22  
23  
24 812 modularity; networks 53, 12 and 4 the highest corrected complementary specialization; and  
25 networks 10, 53 and 4 the highest corrected modularity, respectively. Complementary  
26  
27  
28 814 specialization measures the exclusiveness in species' interactions, whereas modularity  
29 quantifies whether species interact more frequently with subsets of available species within a  
30  
31  
32 816 community. Network drawings were created using Network3D and energized with the 3D  
33 Force-directed algorithm to enhance visualization of modularity patterns (Yoon et al. 2004,  
34  
35  
36 818 Williams 2010). The grey shading of the background illustrates altitude, with darker shades  
37 depicting higher altitudes.  
38  
39  
40  
41

42 820 **Figure 3.** Coefficients of determination ( $R^2$ ) for complementary specialization and modularity  
43 obtained from partial regression of full models, i.e. in models including all ten predictor  
44  
45  
46 822 variables (see Table 1 for standardized coefficients of each variable and more details of  
47 model fit, and Table S3 for the  $R^2$  values used in this figure). Complementary specialization  
48  
49  
50 824 measures the exclusiveness in species' interactions, whereas modularity quantifies whether  
51 species interact more frequently with subsets of available species within a community. We  
52  
53  
54 826 represent values for all networks in the study (Global dataset; n=54) and excluding insular  
55 communities (Only Mainland; n=45). Bars illustrate the association between the different  
56  
57  
58  
59  
60


1  
2  
3 828 "predictor-types" and network structure. Predictor-types refers to (a) species richness (one  
4 variable), (b) phylogenetic signal (two variables: phylogenetic signal in the interaction pattern  
5 of plants and hummingbirds), and (c) environmental factors (eight variables: insularity,  
6  
7 830 average annual temperature, temperature seasonality, total annual precipitation, precipitation  
8 seasonality, temperature and precipitation-change velocity between the Last Glacial  
9  
10 832 Maximum and the present). The amount of variation explained by each pooled predictor-type  
11 is color-coded with different shades of blue: bars colored in darkest blue depict the overall  
12 variation explained by all factors together, whereas medium-blue illustrate the total variation  
13 explained by that predictor-type, and light-blue the unique variation explained by each  
14 predictor-type and not shared by other variable-types.  $H_2'$ , complementary specialization;  
15  
16 834  
17  
18  
19  
20 836  
21  
22  
23  
24 838 QuanBiMo, quantitative bipartite modularity.  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Figure 1.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60


Figure 2.



842

iew

Figure 3.



844



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



**Appendix S1.** List of all plant and hummingbird species present in the studied networks.

**Plant species**

Family	Plant Species
Acanthaceae	<i>Anisacanthus brasiliensis</i> Lindau
Acanthaceae	<i>Aphelandra aurantiaca</i> (Scheidw.) Lindl.
Acanthaceae	<i>Aphelandra colorata</i> (Vell.) Wassh.
Acanthaceae	<i>Aphelandra macrostachya</i> Nees
Acanthaceae	<i>Aphelandra</i> sp. 1
Acanthaceae	<i>Aphelandra</i> sp. 2
Acanthaceae	<i>Aphelandra storkii</i> Leonard
Acanthaceae	<i>Dicliptera pohliana</i> Nees
Acanthaceae	<i>Dicliptera squarrosa</i> Nees
Acanthaceae	<i>Geissomeria longiflora</i> Lindl.
Acanthaceae	<i>Geissomeria schottiana</i> Nees
Acanthaceae	<i>Geissomeria</i> sp.
Acanthaceae	<i>Justicia brasiliana</i> Roth
Acanthaceae	<i>Justicia candicans</i> (Nees) L.D.Benson
Acanthaceae	<i>Justicia carnea</i> Lindl.
Acanthaceae	<i>Justicia cuatrecasasii</i> Wassh.
Acanthaceae	<i>Justicia scheidweileri</i> V.A.W. Graham
Acanthaceae	<i>Justicia sebastianopolitanae</i> Profice
Acanthaceae	<i>Justicia secunda</i> Vahl
Acanthaceae	<i>Justicia</i> sp. 1
Acanthaceae	<i>Justicia</i> sp. 2
Acanthaceae	<i>Justicia tenuistachys</i> (Rusby) Wassh. & J.R.I. Wood
Acanthaceae	<i>Mendoncia</i> sp.
Acanthaceae	<i>Mendoncia coccinea</i> Ruiz & Pav.
Acanthaceae	<i>Odontonema barleriodes</i> Kuntze
Acanthaceae	<i>Pachystachys coccinea</i> (Aubl.) Nees
Acanthaceae	<i>Ruellia asperula</i> (Mart. & Nees) Lindau
Acanthaceae	<i>Ruellia brevifolia</i> (Pohl) C.Ezcurra
Acanthaceae	<i>Ruellia inflata</i> Rich.
Acanthaceae	<i>Ruellia</i> sp.
Acanthaceae	<i>Ruellia subsessilis</i> Lindau
Acanthaceae	<i>Sanchezia munita</i> (Nees) Planch.
Acanthaceae	<i>Sanchezia oblonga</i> Ruiz & Pav.
Acanthaceae	<i>Sanchezia putumayensis</i> Leonard
Acanthaceae	<i>Sanchezia scandens</i> (Lindau) Leonard & L.B. Sm.
Acanthaceae	<i>Staurogyne itatiaiae</i> (Wawra) Leonard
Acanthaceae	<i>Suessenguthia multisetosa</i> (Rusby) Wassh. & J.R.I. Wood
Adoxaceae	<i>Sambucus</i> sp.
Alstroemiaceae	<i>Alstroemeria inodora</i> Herb.
Alstroemiaceae	<i>Alstroemeria isabellana</i> Herb.
Alstroemiaceae	<i>Alstroemeria rupestris</i> M.C.Assis
Alstroemiaceae	<i>Alstroemeria</i> sp.

Family	Plant Species
Alstroemeriaceae	<i>Bomarea cf. carderi</i> Mast.
Alstroemeriaceae	<i>Bomarea costaricensis</i> Kraenzl.
Alstroemeriaceae	<i>Bomarea edulis</i> (Tussac) Herb.
Alstroemeriaceae	<i>Bomarea hirsuta</i> (Kunth) Herb.
Alstroemeriaceae	<i>Bomarea linifolia</i> (Kunth) Baker
Alstroemeriaceae	<i>Bomarea pardina</i> Herb.
Alstroemeriaceae	<i>Bomarea</i> sp. 1
Alstroemeriaceae	<i>Bomarea</i> sp. 2
Amaryllidaceae	<i>Hippeastrum aulicum</i> (Ker Gawl.) Herb.
Amaryllidaceae	<i>Hippeastrum aviflorum</i> (Ravenna) Dutilh
Amaryllidaceae	<i>Hippeastrum puniceum</i> (Lam.) Voss
Amaryllidaceae	<i>Hippeastrum cipoanum</i> (Ravenna) Meerow
Apocynaceae	<i>Allamanda blanchetii</i> A.DC.
Apocynaceae	<i>Blepharodon pictum</i> (Vahl) W.D.Stevens
Apocynaceae	<i>Ditassa retusa</i> Mart.
Apocynaceae	<i>Mandevilla aff. mollissima</i> (Kunth) K.Schum.
Apocynaceae	<i>Mandevilla funiformis</i> (Vell.) K.Schum.
Apocynaceae	<i>Mandevilla hirsuta</i> (Rich.) K.Schum.
Apocynaceae	<i>Mandevilla scabra</i> (Hoffmanns. ex Roem. & Schult.) K.Schum.
Apocynaceae	<i>Nephradenia asparagoides</i> (Decne.) E.Fourn.
Apocynaceae	<i>Pentalinon luteum</i> (L.) B.F.Hansen & Wunderlin
Apocynaceae	<i>Prestonia coalita</i> (Vell.) Woodson
Apocynaceae	<i>Stipecoma peltigera</i> (Stadelm.) Müll.Arg.
Apocynaceae	<i>Tabernaemontana alba</i> Mill.
Apocynaceae	<i>Tabernaemontana cymosa</i> Jacq.
Apocynaceae	<i>Tabernaemontana macrocalyx</i> Müll.Arg.
Araceae	<i>Philodendron fragrantissimum</i> (Hook.) G.Don
Asparagaceae	<i>Furcraea</i> sp.
Asteraceae	<i>Acritopappus longifolius</i> (Gardner) R.M.King & H.Rob.
Asteraceae	<i>Asteraceae</i> sp.
Asteraceae	<i>Baccharis salicina</i> Torr. & A.Gray
Asteraceae	<i>Barnadesia horrida</i> Muschl.
Asteraceae	<i>Barnadesia spinosa</i> L.f.
Asteraceae	<i>Chronopappus bifrons</i> (DC. ex Pers.) DC.
Asteraceae	<i>Cirsium subcoriaceum</i> (Less.) Sch.Bip. ex Sch.Bip.
Asteraceae	<i>Conocliniopsis prasiifolia</i> (DC.) R.M.King & H.Rob.
Asteraceae	<i>Dasyphyllum candolleanum</i> (Gardner) Cabrera
Asteraceae	<i>Diplostephium</i> sp.
Asteraceae	<i>Eremanthus crotonoides</i> (DC.) Sch.Bip.
Asteraceae	<i>Eremanthus erythropappus</i> (DC.) MacLeish
Asteraceae	<i>Espeletia</i> sp.
Asteraceae	<i>Gynoxys longifolia</i> Sch.Bip. ex Wedd.
Asteraceae	<i>Hololepis pedunculata</i> (DC. ex Pers.) DC.
Asteraceae	<i>Lepidaploa cotoneaster</i> (Willd. ex Spreng.) H.Rob.
Asteraceae	<i>Lepidaploa</i> sp.
Asteraceae	<i>Lessingianthus linearis</i> (Spreng.) H.Rob.
Asteraceae	<i>Moquinia kingii</i> (H.Rob.) Gambero

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Family	Plant Species
Asteraceae	<i>Mutisia speciosa</i> Aiton ex Hook.
Asteraceae	<i>Neomirandea eximia</i> (B.L.Rob.) R.M.King & H.Rob.
Asteraceae	<i>Peltophorum dubium</i> (Spreng.) Taub.
Asteraceae	<i>Piptocarpha notata</i> (Less.) Baker
Asteraceae	<i>Piptolepis leptospermoides</i> (Mart. ex DC.) Sch.Bip.
Asteraceae	<i>Pithecoseris pacourinoides</i> Mart. ex DC.
Asteraceae	<i>Trixis vauthieri</i> DC.
Asteraceae	<i>Vernonanthura phosphorica</i> (Vell.) H.Rob.
Balsaminaceae	<i>Impatiens</i> sp.
Balsaminaceae	<i>Impatiens walleriana</i> Hook.f.
Bignoniaceae	<i>Adenocalymma campicola</i> (Pilg.) L.G.Lohmann
Bignoniaceae	<i>Bignonia binata</i> Thunb.
Bignoniaceae	<i>Bignoniaceae</i> sp.
Bignoniaceae	<i>Campsis</i> cf. <i>Grandiflora</i> (Thunb.) K.Schum.
Bignoniaceae	<i>Cuspidaria inaequalis</i> (DC. ex Splitg.) L.G.Lohmann
Bignoniaceae	<i>Cuspidaria lateriflora</i> (Mart.) DC.
Bignoniaceae	<i>Cuspidaria simplicifolia</i> DC.
Bignoniaceae	<i>Dolichandra unguis-cati</i> (L.) L.G.Lohmann
Bignoniaceae	<i>Fridericia erubescens</i> (DC.) L.G.Lohmann
Bignoniaceae	<i>Fridericia florida</i> (DC.) L.G.Lohmann
Bignoniaceae	<i>Fridericia speciosa</i> Mart.
Bignoniaceae	<i>Fridericia triplinervia</i> (Mart. ex DC.) L.G.Lohmann
Bignoniaceae	<i>Handroanthus chrysotrichus</i> (Mart. ex DC.) Mattos
Bignoniaceae	<i>Handroanthus heptaphyllus</i> (Vell.) Mattos
Bignoniaceae	<i>Handroanthus impetiginosus</i> (Mart. ex DC.) Mattos
Bignoniaceae	<i>Handroanthus selachidentatus</i> (A.H.Gentry) S.O.Grose
Bignoniaceae	<i>Jacaranda irwinii</i> A.H.Gentry
Bignoniaceae	<i>Jacaranda mimosifolia</i> D.Don
Bignoniaceae	<i>Jacaranda puberula</i> Cham.
Bignoniaceae	<i>Jacaratia spinosa</i> (Aubl.) A.DC.
Bignoniaceae	<i>Lundia cordata</i> (Vell.) DC.
Bignoniaceae	<i>Pyrostegia venusta</i> (Ker Gawl.) Miers
Bignoniaceae	<i>Tabebuia aurea</i> (Silva Manso) Benth. & Hook.f. ex S.Moore
Bignoniaceae	<i>Tabebuia heterophylla</i> (DC.) Britton
Bignoniaceae	<i>Tabebuia orinocensis</i> (Sandwith) A.H.Gentry
Bignoniaceae	<i>Tabebuia</i> sp.
Bignoniaceae	<i>Tabebuia stenocalyx</i> Sprague & Stapf
Bignoniaceae	<i>Tecoma stans</i> (L.) Juss. ex Kunth
Bignoniaceae	<i>Zeyheria montana</i> Mart.
Boraginacea	<i>Cordia bullata</i> var. <i>globosa</i> (Jacq.) Govaerts
Boraginacea	<i>Cordia multispicata</i> Cham.
Boraginacea	<i>Cordia seleriana</i> Fernald
Boraginacea	<i>Cordia superba</i> Cham.
Boraginacea	<i>Cordia bicolor</i> A.DC.
Boraginacea	<i>Cordia curassavica</i> (Jacq.) Roem. & Schult.
Boraginacea	<i>Cordia glabrata</i> (Mart.) A.DC.
Bromeliaceae	<i>Aechmea aquilega</i> (Salisb.) Griseb.

Family	Plant Species
Bromeliaceae	<i>Aechmea blumenavii</i> Reitz
Bromeliaceae	<i>Aechmea bromeliifolia</i> (Rudge) Baker
Bromeliaceae	<i>Aechmea chantinii</i> (Carrière) Baker
Bromeliaceae	<i>Aechmea contracta</i> (Mart. ex Schult. & Schult.f.) Baker
Bromeliaceae	<i>Aechmea corymbosa</i> (Mart. ex Schult. & Schult.f.) Mez
Bromeliaceae	<i>Aechmea cylindrata</i> Lindm.
Bromeliaceae	<i>Aechmea dichlamydea</i> Baker
Bromeliaceae	<i>Aechmea distichantha</i> Lem.
Bromeliaceae	<i>Aechmea fendleri</i> André ex Mez
Bromeliaceae	<i>Aechmea gamosepala</i> Wittm.
Bromeliaceae	<i>Aechmea longifolia</i> (Rudge) L.B.Sm. & M.A.Spencer
Bromeliaceae	<i>Aechmea mariae-reginae</i> H.Wendl.
Bromeliaceae	<i>Aechmea nudicaulis</i> (L.) Griseb.
Bromeliaceae	<i>Aechmea organensis</i> Wawra
Bromeliaceae	<i>Aechmea pectinata</i> Baker
Bromeliaceae	<i>Aechmea poitaei</i> (Baker) L.B.Sm. & M.A.Spencer
Bromeliaceae	<i>Aechmea recurvata</i> (Klotzsch) L.B.Sm.
Bromeliaceae	<i>Aechmea rubiginosa</i> Mez
Bromeliaceae	<i>Aechmea</i> sp.
Bromeliaceae	<i>Aechmea van-houtteana</i> (Van Houtte) Mez
Bromeliaceae	<i>Aechmea williamsii</i> (L.B.Sm.) L.B.Sm. & M.A.Spencer
Bromeliaceae	<i>Ananas ananassoides</i> (Baker) L.B.Sm.
Bromeliaceae	<i>Ananas comosus</i> (L.) Merr.
Bromeliaceae	<i>Billbergia amoena</i> (Lodd.) Lindl.
Bromeliaceae	<i>Billbergia decora</i> Poepp. & Endl.
Bromeliaceae	<i>Billbergia distachya</i> (Vell.) Mez
Bromeliaceae	<i>Billbergia pyramidalis</i> (Sims) Lindl.
Bromeliaceae	<i>Billbergia vittata</i> Brongn. ex C.Morel
Bromeliaceae	<i>Bromelia antiacantha</i> Bertol.
Bromeliaceae	<i>Bromelia balansae</i> Mez
Bromeliaceae	<i>Bromelia</i> sp.
Bromeliaceae	<i>Bromeliaceae</i> sp. 1
Bromeliaceae	<i>Bromeliaceae</i> sp. 2
Bromeliaceae	<i>Canistrum cyathiforme</i> (Vell.) Mez
Bromeliaceae	<i>Canistrum fragrans</i> (Linden) Mabb.
Bromeliaceae	<i>Canistrum giganteum</i> (Baker) L.B.Sm.
Bromeliaceae	<i>Canistrum perplexum</i> L.B.Sm.
Bromeliaceae	<i>Dyckia leptostachya</i> Baker
Bromeliaceae	<i>Dyckia</i> sp.
Bromeliaceae	<i>Encholirium brachypodium</i> L.B.Sm. & Read
Bromeliaceae	<i>Encholirium spectabile</i> Mart. ex Schult. & Schult.f.
Bromeliaceae	<i>Encholirium subsecundum</i> (Baker) Mez
Bromeliaceae	<i>Guzmania bakeri</i> (Wittm.) Mez
Bromeliaceae	<i>Guzmania berteroniana</i> (Schult. & Schult. f.) Mez
Bromeliaceae	<i>Guzmania candelabrum</i> (André) André ex Mez
Bromeliaceae	<i>Guzmania squarrosa</i> (Mez & Sodiro) L.B.Sm. & Pittendr.
Bromeliaceae	<i>Guzmania danielii</i> L.B.Sm.

	Family	Plant Species
1		
2		
3		
4	Bromeliaceae	<i>Guzmania jaramilloi</i> H.E.Luther
5	Bromeliaceae	<i>Guzmania monostachia</i> (L.) Rusby ex Mez
6	Bromeliaceae	<i>Guzmania nicaraguensis</i> Mez & C.F.Baker
7	Bromeliaceae	<i>Guzmania</i> sp. 1
8	Bromeliaceae	<i>Guzmania</i> sp. 2
9	Bromeliaceae	<i>Guzmania</i> sp. 3
10	Bromeliaceae	<i>Guzmania teuscheri</i> L.B.Sm.
11	Bromeliaceae	<i>Hohenbergia cf. catingae</i> Ule
12	Bromeliaceae	<i>Hohenbergia ridleyi</i> (Baker) Mez
13	Bromeliaceae	<i>Neoglaziovia variegata</i> (Arruda) Mez
14	Bromeliaceae	<i>Neoregelia bahiana</i> (Ule) L.B.Sm.
15	Bromeliaceae	<i>Nidularium rutilans</i> E. Morren
16	Bromeliaceae	<i>Nidularium innocentii</i> Lem.
17	Bromeliaceae	<i>Nidularium itatiaiae</i> L.B.Sm.
18	Bromeliaceae	<i>Nidularium marigoii</i> Leme
19	Bromeliaceae	<i>Nidularium procerum</i> Lindm.
20	Bromeliaceae	<i>Nidularium seidelii</i> L.B.Sm. & Reitz
21	Bromeliaceae	<i>Nidularium</i> sp.
22	Bromeliaceae	<i>Orthophytum albopictum</i> Philcox
23	Bromeliaceae	<i>Orthophytum lemei</i> E.Pereira & I.A.Penna
24	Bromeliaceae	<i>Pitcairnia brittoniana</i> (Mez) Mez
25	Bromeliaceae	<i>Pitcairnia flammea</i> Lindl.
26	Bromeliaceae	<i>Pitcairnia nigra</i> (Carrière) André
27	Bromeliaceae	<i>Pitcairnia</i> sp.
28	Bromeliaceae	<i>Pitcairnia sprucei</i> Baker
29	Bromeliaceae	<i>Puya clava-herculis</i> Mez & Sodiro
30	Bromeliaceae	<i>Puya ferruginea</i> (Ruiz & Pav.) L.B.Sm.
31	Bromeliaceae	<i>Quesnelia augusto-coburgii</i> Wawra
32	Bromeliaceae	<i>Quesnelia</i> sp.
33	Bromeliaceae	<i>Tillandsia aeranthos</i> (Loisel.) L.B.Sm.
34	Bromeliaceae	<i>Tillandsia aff. turneri</i> Baker
35	Bromeliaceae	<i>Tillandsia bartramii</i> Elliott
36	Bromeliaceae	<i>Tillandsia complanata</i> Benth.
37	Bromeliaceae	<i>Tillandsia dasyliirifolia</i> Baker
38	Bromeliaceae	<i>Tillandsia fasciculata</i> Sw.
39	Bromeliaceae	<i>Tillandsia gardneri</i> Lindl.
40	Bromeliaceae	<i>Tillandsia orbicularis</i> L.B.Sm.
41	Bromeliaceae	<i>Tillandsia paucifolia</i> Baker
42	Bromeliaceae	<i>Tillandsia</i> sp. 1
43	Bromeliaceae	<i>Tillandsia</i> sp. 2
44	Bromeliaceae	<i>Tillandsia stricta</i> Sol. ex Ker Gawl.
45	Bromeliaceae	<i>Tillandsia tetrantha</i> Ruiz & Pav.
46	Bromeliaceae	<i>Tillandsia utriculata</i> L.
47	Bromeliaceae	<i>Vriesea altodaserrae</i> L.B.Sm.
48	Bromeliaceae	<i>Vriesea carinata</i> Wawra
49	Bromeliaceae	<i>Vriesea chrysostachys</i> E.Morren
50	Bromeliaceae	<i>Vriesea ensiformis</i> (Vell.) Beer
51		
52		
53		
54		
55		
56		
57		
58		
59		
60		

Family	Plant Species
Bromeliaceae	<i>Vriesea erythrodactylon</i> (E.Morren) E.Morren ex Mez
Bromeliaceae	<i>Vriesea incurvata</i> Gaudich.
Bromeliaceae	<i>Vriesea jonghei</i> (K.Koch) E.Morren
Bromeliaceae	<i>Vriesea longicaulis</i> (Baker) Mez
Bromeliaceae	<i>Vriesea medusa</i> Versieux
Bromeliaceae	<i>Vriesea ororiensis</i> (Mez) L.B.Sm. & Pittendr.
Bromeliaceae	<i>Vriesea philippocoburgii</i> Wawra
Bromeliaceae	<i>Vriesea procera</i> (Mart. ex Schult. & Schult.f.) Wittm.
Bromeliaceae	<i>Vriesea procera</i> var. <i>tenuis</i> L.B.Sm.
Bromeliaceae	<i>Vriesea rodigasiana</i> E.Morren
Bromeliaceae	<i>Vriesea sceptrum</i> Mez
Bromeliaceae	<i>Vriesea simplex</i> (Vell.) Beer
Bromeliaceae	<i>Vriesea</i> sp. 1
Bromeliaceae	<i>Vriesea</i> sp. 2
Bromeliaceae	<i>Vriesea</i> sp. 3
Bromeliaceae	<i>Vriesea vagans</i> (L.B.Sm.) L.B.Sm.
Burseraceae	<i>Protium</i> sp.
Cactaceae	<i>Cipocereus minensis</i> (Werderm.) F.Ritter
Cactaceae	<i>Consolea rubescens</i> (Salm-Dyck ex DC.) Lem.
Cactaceae	<i>Melocactus concinnus</i> Buining & Brederoo
Cactaceae	<i>Melocactus ernestii</i> Vaupel
Cactaceae	<i>Melocactus intortus</i> (Mill.) Urb.
Cactaceae	<i>Melocactus zehntneri</i> (Britton & Rose) Luetzelb.
Cactaceae	<i>Micranthocereus flaviflorus</i> Buining & Brederoo
Cactaceae	<i>Micranthocereus purpureus</i> (Gürke) F. Ritter
Cactaceae	<i>Opuntia excelsa</i> Sánchez-Mej.
Cactaceae	<i>Opuntia karwinskiana</i> Salm-Dyck
Cactaceae	<i>Opuntia wernerii</i> Eggl
Cactaceae	<i>Pilosocereus aurisetus</i> (Werderm.) Byles & G.D. Rowley
Cactaceae	<i>Pilosocereus gounellei</i> (F.A.C.Weber ex K.Schum.) Byles & G.D.Rowley
Cactaceae	<i>Pilosocereus pachycladus</i> F. Ritter
Cactaceae	<i>Pilosocereus royenii</i> (L.) Byles & G.D.Rowley
Cactaceae	<i>Pilosocereus</i> sp.
Cactaceae	<i>Tacinga inamoena</i> (K.Schum.) N.P.Taylor & Stuppy
Cactaceae	<i>Tacinga palmadora</i> (Britton & Rose) N.P.Taylor & Stuppy
Campanulaceae	<i>Burmeistera cyclostigmata</i> Donn.Sm.
Campanulaceae	<i>Burmeistera globosa</i> E.Wimm.
Campanulaceae	<i>Burmeistera parviflora</i> E.Wimm. ex Standl.
Campanulaceae	<i>Burmeistera</i> sp.
Campanulaceae	<i>Centropogon cornutus</i> (L.) Druce
Campanulaceae	<i>Centropogon costaricae</i> (Vatke) McVaugh
Campanulaceae	<i>Centropogon granulatus</i> C.Presl
Campanulaceae	<i>Centropogon solanifolius</i> Benth.
Campanulaceae	<i>Centropogon</i> sp. 1
Campanulaceae	<i>Centropogon</i> sp. 2
Campanulaceae	<i>Centropogon talamancensis</i> Wilbur
Campanulaceae	<i>Centropogon valerii</i> Standl.

1		
2		
3	<b>Family</b>	<b>Plant Species</b>
4	Campanulaceae	<i>Lobelia cirsiifolia</i> Lam.
5	Campanulaceae	<i>Lobelia fistulosa</i> Vell.
6	Campanulaceae	<i>Lobelia stricta</i> Sw.
7	Campanulaceae	<i>Siphocampylus bogotensis</i> E.Wimm.
8	Campanulaceae	<i>Siphocampylus cf. convolvulaceus</i> (Cham.) G.Don
9	Campanulaceae	<i>Siphocampylus fimbriatus</i> Regel
10	Campanulaceae	<i>Siphocampylus giganteus</i> (Cav.) G.Don
11	Campanulaceae	<i>Siphocampylus imbricatus</i> (Cham.) G.Don
12	Campanulaceae	<i>Siphocampylus longipedunculatus</i> Pohl
13	Campanulaceae	<i>Siphocampylus paramicola</i> McVaugh
14	Campanulaceae	<i>Siphocampylus sp.</i>
15	Campanulaceae	<i>Siphocampylus sulfureus</i> E.Wimm.
16	Campanulaceae	<i>Siphocampylus westinianus</i> (Thunb.) Pohl
17	Cannaceae	<i>Canna indica</i> L.
18	Cannaceae	<i>Canna paniculata</i> Ruiz & Pav.
19	Cannaceae	<i>Canna sp.</i>
20	Capparaceae	<i>Crateva tapia</i> L.
21	Caprifoliaceae	<i>Lonicera japonica</i> Thunb.
22	Caryocaraceae	<i>Caryocar brasiliense</i> A.St.-Hil.
23	Chrysobalanaceae	<i>Couepia schottii</i> Fritsch
24	Clusiaceae	<i>Clusia mangle</i> Rich. ex Planch. & Triana
25	Clusiaceae	<i>Clusia melchiorii</i> Gleason
26	Clusiaceae	<i>Clusia sp. 1</i>
27	Clusiaceae	<i>Clusia sp. 2</i>
28	Clusiaceae	<i>Symphonia globulifera</i> L.f.
29	Columelliaceae	<i>Desfontainia spinosa</i> Ruiz & Pav.
30	Combretaceae	<i>Combretum farinosum</i> Kunth
31	Combretaceae	<i>Combretum llewelynii</i> J.F.Macbr.
32	Convolvulaceae	<i>Aniseia martinicensis</i> (Jacq.) Choisy
33	Convolvulaceae	<i>Ipomoea aquatica</i> Forssk.
34	Convolvulaceae	<i>Ipomoea bracteata</i> Cav.
35	Convolvulaceae	<i>Ipomoea quamoclit</i> L.
36	Convolvulaceae	<i>Ipomoea sp. 1</i>
37	Convolvulaceae	<i>Ipomoea sp. 2</i>
38	Convolvulaceae	<i>Ipomoea sp. 3</i>
39	Convolvulaceae	<i>Ipomoea subincana</i> Meisn.
40	Convolvulaceae	<i>Jacquemontia sp. 1</i>
41	Convolvulaceae	<i>Jacquemontia sp. 2</i>
42	Convolvulaceae	<i>Merremia macrocalyx</i> (Ruiz & Pav.) O'Donell
43	Costaceae	<i>Costus curvibracteatus</i> Maas
44	Costaceae	<i>Costus lasius</i> Loes.
45	Costaceae	<i>Costus pulverulentus</i> C.Presl
46	Costaceae	<i>Costus scaber</i> Ruiz & Pav.
47	Costaceae	<i>Costus spiralis</i> (Jacq.) Roscoe
48	Costaceae	<i>Dimerocostus strobilaceus</i> Kuntze
49	Crassulaceae	<i>Kalanchoe sp.</i>
50	Cucurbitaceae	<i>Apodanthera villosa</i> C.Jeffrey
51		
52		
53		
54		
55		
56		
57		
58		
59		
60		

Family	Plant Species
Cucurbitaceae	<i>Gurania acuminata</i> Cogn.
Cucurbitaceae	<i>Gurania coccinea</i> Cogn.
Cucurbitaceae	<i>Gurania lobata</i> (L.) Pruski
Cucurbitaceae	<i>Gurania rhizantha</i> (Poepp. & Endl.) C.Jeffrey
Cucurbitaceae	<i>Gurania rufipila</i> Cogn.
Cucurbitaceae	<i>Psiguria ternata</i> (M.Roem.) C.Jeffrey
Ericaceae	<i>Agarista coriifolia</i> (Thunb.) Hook. ex Nied.
Ericaceae	<i>Agarista</i> sp.
Ericaceae	<i>Bejaria aestuans</i> Mutis ex L.
Ericaceae	<i>Cavendishia bracteata</i> (Ruiz & Pav. ex J.St.Hil.) Hoerold
Ericaceae	<i>Cavendishia callista</i> Donn.Sm.
Ericaceae	<i>Cavendishia complectens</i> Hemsl.
Ericaceae	<i>Cavendishia grandifolia</i> Herold
Ericaceae	<i>Cavendishia guatapeensis</i> Mansf.
Ericaceae	<i>Cavendishia pubescens</i> (Kunth) Hemsl.
Ericaceae	<i>Cavendishia quereme</i> (Kunth) Benth. & Hook.f.
Ericaceae	<i>Cavendishia</i> sp.
Ericaceae	<i>Cavendishia tarapotana</i> Benth. & Hook.f.
Ericaceae	<i>Disterigma codonanthum</i> S.F.Blake
Ericaceae	<i>Disterigma humboldtii</i> (Klotzsch) Nied.
Ericaceae	<i>Disterigma</i> sp. 1
Ericaceae	<i>Disterigma</i> sp. 2
Ericaceae	<i>Ericaceae</i> sp. 1
Ericaceae	<i>Ericaceae</i> sp. 2
Ericaceae	<i>Gaultheria erecta</i> Vent.
Ericaceae	<i>Gaultheria gracilis</i> Small
Ericaceae	<i>Gaultheria insipida</i> Benth.
Ericaceae	<i>Gaultheria myrsinoides</i> Kunth
Ericaceae	<i>Gaultheria</i> sp.
Ericaceae	<i>Gaylussacia brasiliensis</i> (Spreng.) Meisn.
Ericaceae	<i>Gaylussacia montana</i> (Pohl) Sleumer
Ericaceae	<i>Gaylussacia oleifolia</i> Dunal
Ericaceae	<i>Gaylussacia virgata</i> Mart. ex Meisn.
Ericaceae	<i>Gonocalyx pterocarpus</i> (Donn.Sm.) Luteyn
Ericaceae	<i>Macleania pentaptera</i> Hoerold
Ericaceae	<i>Macleania recumbens</i> A.C. Sm.
Ericaceae	<i>Macleania rupestris</i> (Kunth) A.C.Sm.
Ericaceae	<i>Psammisia aberrans</i> A.C. Sm.
Ericaceae	<i>Psammisia ecuadorensis</i> Hoerold
Ericaceae	<i>Psammisia falcata</i> Klotzsch
Ericaceae	<i>Psammisia oreogenes</i> Sleumer
Ericaceae	<i>Psammisia pauciflora</i> Griseb. ex A.C. Sm.
Ericaceae	<i>Psammisia penduliflora</i> (Dunal) Klotzsch
Ericaceae	<i>Psammisia ramiflora</i> Klotzsch
Ericaceae	<i>Psammisia sodiroi</i> Hoerold
Ericaceae	<i>Psammisia ulbrichiana</i> Hoerold
Ericaceae	<i>Satyria meiantha</i> Donn.Sm.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Family	Plant Species
Ericaceae	<i>Satyria panurensis</i> (Benth. ex Meisn.) Benth. & Hook.f. ex Nied.
Ericaceae	<i>Thibaudia costaricensis</i> Hoerold
Ericaceae	<i>Thibaudia rigidiflora</i> A.C.Sm.
Ericaceae	<i>Vaccinium consanguineum</i> Klotzsch
Ericaceae	<i>Vaccinium floribundum</i> Kunth
Ericaceae	<i>Vaccinium racemosum</i> (Vahl) Wilbur & Luteyn
Ericaceae	<i>Vaccinium poasanum</i> Donn.Sm.
Erythroxylaceae	<i>Erythroxylum vaccinifolium</i> Mart.
Euphorbiaceae	<i>Cnidoscopus halteris</i> Fern.Casas
Euphorbiaceae	<i>Cnidoscopus pubescens</i> Pohl
Euphorbiaceae	<i>Croton blanchetianus</i> Baill.
Euphorbiaceae	<i>Jatropha mollissima</i> (Pohl) Baill.
Euphorbiaceae	<i>Jatropha mutabilis</i> (Pohl) Baill.
Euphorbiaceae	<i>Manihot jacobinensis</i> Müll.Arg.
Euphorbiaceae	<i>Manihot reniformis</i> Pohl
Fabaceae	<i>Abarema brachystachya</i> (DC.) Barneby & J.W.Grimes
Fabaceae	<i>Albizia saman</i> (Jacq.) Merr.
Fabaceae	<i>Anadenanthera colubrina</i> (Vell.) Brenan
Fabaceae	<i>Bauhinia brevipes</i> Vogel
Fabaceae	<i>Bauhinia cheilantha</i> (Bong.) Steud.
Fabaceae	<i>Bauhinia forficata</i> Link
Fabaceae	<i>Bauhinia longifolia</i> (Bong.) Steud.
Fabaceae	<i>Bauhinia rufa</i> (Bong.) Steud.
Fabaceae	<i>Bauhinia unguolata</i> L.
Fabaceae	<i>Bauhinia variegata</i> L.
Fabaceae	<i>Bowdichia virgilioides</i> Kunth
Fabaceae	<i>Brownea coccinea</i> subsp. <i>capitella</i> (Jacq.) D.Velasquez & Agostini
Fabaceae	<i>Caesalpinia pyramidalis</i> Tul.
Fabaceae	<i>Calliandra brevipes</i> Benth.
Fabaceae	<i>Calliandra carbonaria</i> Benth.
Fabaceae	<i>Calliandra guildingii</i> Benth.
Fabaceae	<i>Calliandra hygrophila</i> Mackinder & G.P.Lewis
Fabaceae	<i>Calliandra mucugeana</i> Renvoize
Fabaceae	<i>Calliandra purdiaei</i> Benth.
Fabaceae	<i>Calliandra tweedii</i> Benth.
Fabaceae	<i>Calliandra viscidula</i> Benth.
Fabaceae	<i>Camptosema coriaceum</i> Benth.
Fabaceae	<i>Camptosema ellipticum</i> (Desv.) Burkart
Fabaceae	<i>Camptosema scarlatinum</i> (Benth.) Burkart
Fabaceae	<i>Centrosema brasilianum</i> (L.) Benth.
Fabaceae	<i>Centrosema coriaceum</i> Benth.
Fabaceae	<i>Chaetocalyx subulatus</i> Mackinder
Fabaceae	<i>Clathrotropis brachypetala</i> (Tul.) Kleinhoonte
Fabaceae	<i>Collaea speciosa</i> (Loisel.) DC.
Fabaceae	<i>Dahlstedtia pinnata</i> (Benth.) Malme
Fabaceae	<i>Dioclea grandiflora</i> Benth.
Fabaceae	<i>Dioclea</i> sp.

Family	Plant Species
Fabaceae	<i>Erythrina corallodendron</i> L.
Fabaceae	<i>Erythrina crista-galli</i> L.
Fabaceae	<i>Erythrina fusca</i> Lour.
Fabaceae	<i>Erythrina lanata</i> Rose
Fabaceae	<i>Erythrina poeppigiana</i> (Walp.) O.F.Cook
Fabaceae	<i>Erythrina</i> sp.
Fabaceae	<i>Erythrina speciosa</i> Andrews
Fabaceae	<i>Fabaceae</i> sp.
Fabaceae	<i>Inga edulis</i> Mart.
Fabaceae	<i>Inga ingoides</i> (Rich.) Willd.
Fabaceae	<i>Inga leiocalycina</i> Benth.
Fabaceae	<i>Inga luschnathiana</i> Benth.
Fabaceae	<i>Inga sessilis</i> (Vell.) Mart.
Fabaceae	<i>Inga</i> sp. 1
Fabaceae	<i>Inga</i> sp. 2
Fabaceae	<i>Inga</i> sp. 3
Fabaceae	<i>Inga venosa</i> Griseb.
Fabaceae	<i>Inga vera</i> Willd.
Fabaceae	<i>Lonchocarpus benthamianus</i> Pittier
Fabaceae	<i>Lysiloma latisiliquum</i> (L.) Benth.
Fabaceae	<i>Mucuna rostrata</i> Benth.
Fabaceae	<i>Mysanthus uleanus</i> (Harms) G.P.Lewis & A.Delgado
Fabaceae	<i>Neorudolphia volubilis</i> (Willd.) Britton
Fabaceae	<i>Periandra coccinea</i> (Schrad.) Benth.
Fabaceae	<i>Periandra mediterranea</i> (Vell.) Taub.
Fabaceae	<i>Phaseolus coccineus</i> L.
Fabaceae	<i>Abarema jupunba</i> (Willd.) Britton & Killip
Fabaceae	<i>Senna reticulata</i> (Willd.) H.S.Irwin & Barneby
Fabaceae	<i>Tachigali paniculata</i> Aubl.
Fabaceae	<i>Tephrosia noctiflora</i> Baker
Fabaceae	<i>Tipuana tipu</i> (Benth.) Kuntze
Fabaceae	<i>Viguieranthus alternans</i> (Benth.) Villiers
Fabaceae	<i>Zygia lathetica</i> Barneby & J.W. Grimes
Gentianaceae	<i>Chelonanthus alatus</i> (Aubl.) Pulle
Gentianaceae	<i>Calolisianthus pendulus</i> (Mart.) Gilg
Gentianaceae	<i>Lisianthus roseus</i> Sessé & Moc.
Gentianaceae	<i>Macrocarpaea cf. rubra</i> Malme
Gentianaceae	<i>Macrocarpaea</i> sp.
Gentianaceae	<i>Macrocarpaea valerioi</i> Standl.
Gentianaceae	<i>Symbolanthus tricolor</i> Gilg
Gentianaceae	<i>Tachia occidentalis</i> Maguire & Weaver
Gesneriaceae	<i>Alloplectus</i> sp. 1
Gesneriaceae	<i>Alloplectus</i> sp. 2
Gesneriaceae	<i>Besleria cf. solanoides</i> Kunth
Gesneriaceae	<i>Besleria columneoides</i> Hanst.
Gesneriaceae	<i>Besleria longimucronata</i> Hoehne
Gesneriaceae	<i>Besleria notabilis</i> C.V.Morton

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Family	Plant Species
Gesneriaceae	<i>Besleria rotundifolia</i> Rusby
Gesneriaceae	<i>Besleria solanoides</i> Kunth
Gesneriaceae	<i>Besleria petiolaris</i> Griseb.
Gesneriaceae	<i>Columnea ciliata</i> (Wiehler) L.P.Kvist & L.E.Skog
Gesneriaceae	<i>Columnea ericae</i> Mansf.
Gesneriaceae	<i>Columnea magnifica</i> Klotzsch ex Oerst.
Gesneriaceae	<i>Columnea medicinalis</i> (Wiehler) L.E.Skog & L.P.Kvist
Gesneriaceae	<i>Columnea microcalyx</i> Hanst.
Gesneriaceae	<i>Columnea purpurata</i> Hanst.
Gesneriaceae	<i>Columnea querceti</i> Oerst.
Gesneriaceae	<i>Columnea</i> sp.
Gesneriaceae	<i>Columnea strigosa</i> Benth.
Gesneriaceae	<i>Drymonia coccinea</i> (Aubl.) Wiehler
Gesneriaceae	<i>Drymonia conchocalyx</i> Hanst.
Gesneriaceae	<i>Drymonia semicordata</i> (Poepp.) Wiehler
Gesneriaceae	<i>Gasteranthus</i> sp.
Gesneriaceae	<i>Gesneriaceae</i> sp. 1
Gesneriaceae	<i>Gesneriaceae</i> sp. 2
Gesneriaceae	<i>Gesneriaceae</i> sp. 3
Gesneriaceae	<i>Gesneriaceae</i> sp. 4
Gesneriaceae	<i>Glossoloma bolivianum</i> (Britton ex Rusby) J.L.Clark
Gesneriaceae	<i>Hypocyrtia fritschii</i> (Hoehne) Handro
Gesneriaceae	<i>Kohleria spicata</i> (Kunth) Oerst.
Gesneriaceae	<i>Kohleria tigridia</i> (Ohlend.) Roalson & Boggan
Gesneriaceae	<i>Nematanthus aff. fritschii</i> Hoehne
Gesneriaceae	<i>Nematanthus australis</i> Chautems
Gesneriaceae	<i>Nematanthus crassifolius</i> (Schott) Wiehler
Gesneriaceae	<i>Nematanthus fluminensis</i> (Vell.) Fritsch
Gesneriaceae	<i>Nematanthus fornix</i> (Vell.) Chautems
Gesneriaceae	<i>Nematanthus gregarius</i> D.L. Denham
Gesneriaceae	<i>Nematanthus lanceolatus</i> (Poir.) Chautems
Gesneriaceae	<i>Nematanthus savannarum</i> (C.V.Morton) J.L.Clark
Gesneriaceae	<i>Nematanthus strigillosus</i> (Mart.) H.E. Moore
Gesneriaceae	<i>Paliavana sericiflora</i> Benth.
Gesneriaceae	<i>Sinningia cooperi</i> (J. Paxton) Wiehler
Gesneriaceae	<i>Sinningia douglasii</i> (Lindl.) Chautems
Gesneriaceae	<i>Sinningia elatior</i> (Kunth) Chautems
Gesneriaceae	<i>Sinningia gigantifolia</i> Chautems
Gesneriaceae	<i>Sinningia glazioviana</i> (Fritsch) Chautems
Gesneriaceae	<i>Sinningia rupicola</i> (Mart.) Wiehler
Heliconiaceae	<i>Heliconia acuminata</i> A.Rich.
Heliconiaceae	<i>Heliconia angusta</i> Vell.
Heliconiaceae	<i>Heliconia atropurpurea</i> G.S.Daniels & F.G.Stiles
Heliconiaceae	<i>Heliconia bihai</i> (L.) L.
Heliconiaceae	<i>Heliconia burleana</i> Abalo & G.Morales
Heliconiaceae	<i>Heliconia subulata</i> Ruiz & Pav. <i>subsp. subulata</i>
Heliconiaceae	<i>Heliconia episcopalis</i> Vell.

Family	Plant Species
Heliconiaceae	<i>Heliconia farinosa</i> Raddi
Heliconiaceae	<i>Heliconia hirsuta</i> L.f.
Heliconiaceae	<i>Heliconia imbricata</i> (Kuntze) Baker
Heliconiaceae	<i>Heliconia juruana</i> Loes.
Heliconiaceae	<i>Heliconia lankesteri</i> Standl.
Heliconiaceae	<i>Heliconia latispatha</i> Benth.
Heliconiaceae	<i>Heliconia marginata</i> (Griggs) Pittier
Heliconiaceae	<i>Heliconia mariae</i> Hook.f.
Heliconiaceae	<i>Heliconia mathiasiae</i> G.S.Daniels & F.G.Stiles
Heliconiaceae	<i>Heliconia metallica</i> Planch. & Linden ex Hook.
Heliconiaceae	<i>Heliconia pogonantha</i> Cufod.
Heliconiaceae	<i>Heliconia psittacorum</i> L.f.
Heliconiaceae	<i>Heliconia rostrata</i> Ruiz & Pav.
Heliconiaceae	<i>Heliconia schumanniana</i> Loes.
Heliconiaceae	<i>Heliconia</i> sp. 1
Heliconiaceae	<i>Heliconia</i> sp. 2
Heliconiaceae	<i>Heliconia</i> sp. 3
Heliconiaceae	<i>Heliconia</i> sp. 4
Heliconiaceae	<i>Heliconia spathocircinata</i> Aristeg.
Heliconiaceae	<i>Heliconia stricta</i> Huber
Heliconiaceae	<i>Heliconia subulata</i> Ruiz & Pav.
Heliconiaceae	<i>Heliconia vaginalis</i> Benth.
Heliconiaceae	<i>Heliconia wagneriana</i> Petersen
Humiriaceae	<i>Humiria balsamifera</i> Aubl.
Iridacea	<i>Iridacea</i> sp.
Lamiaceae	<i>Aegiphila obducta</i> Vell.
Lamiaceae	<i>Aegiphila perplexa</i> Moldenke
Lamiaceae	<i>Amasonia campestris</i> (Aubl.) Moldenke
Lamiaceae	<i>Volkameria aculeata</i> L.
Lamiaceae	<i>Eriope latifolia</i> (Mart. ex Benth.) Harley
Lamiaceae	<i>Hyptidendron asperrimum</i> (Spreng.) Harley
Lamiaceae	<i>Hyptis atrorubens</i> Poit.
Lamiaceae	<i>Hyptis cuniloides</i> Epling
Lamiaceae	<i>Hyptis leptostachys</i> Epling
Lamiaceae	<i>Hyptis pectinata</i> (L.) Poit.
Lamiaceae	<i>Hyptis</i> sp.
Lamiaceae	<i>Lamiaceae</i> sp.
Lamiaceae	<i>Leonotis nepetifolia</i> (L.) R.Br.
Lamiaceae	<i>Prunella vulgaris</i> L.
Lamiaceae	<i>Salvia arenaria</i> A.St.-Hil. ex Benth.
Lamiaceae	<i>Salvia articulata</i> Epling
Lamiaceae	<i>Salvia carnea</i> Kunth
Lamiaceae	<i>Salvia elegans</i> Vahl
Lamiaceae	<i>Salvia mocinoi</i> Benth.
Lamiaceae	<i>Salvia sellowiana</i> Benth.
Lamiaceae	<i>Salvia</i> sp.
Lamiaceae	<i>Vitex cymosa</i> Bertero ex Spreng.

1		
2		
3	<b>Family</b>	<b>Plant Species</b>
4	Lamiaceae	<i>Vitex divaricata</i> Sw.
5	Lecythidaceae	<i>Lecythidoideae</i> sp.
6	Loganiaceae	<i>Spigelia pulchella</i> Mart.
7	Loranthaceae	<i>Gaiadendron punctatum</i> (Ruiz & Pav.) G.Don
8	Loranthaceae	<i>Ligaria teretiflora</i> (Rizzini) Kuijt
9	Loranthaceae	<i>Loranthaceae</i> sp.
10	Loranthaceae	<i>Loranthus bicalyculatus</i> Mart. ex Schult. & Schult.f.
11	Loranthaceae	<i>Psittacanthus cordatus</i> (Hoffmanns. ex Schult. f.) Blume
12	Loranthaceae	<i>Psittacanthus cucullaris</i> (Lam.) G.Don
13	Loranthaceae	<i>Psittacanthus dichrous</i> Mart.
14	Loranthaceae	<i>Psittacanthus flavo-viridis</i> Eichler
15	Loranthaceae	<i>Loranthus furcatus</i> Mart. ex Schult. & Schult.f.
16	Loranthaceae	<i>Psittacanthus lasianthus</i> Sandwith
17	Loranthaceae	<i>Tripodanthus acutifolius</i> (Ruiz & Pav.) Tiegh.
18	Loranthaceae	<i>Tristerix longebracteatus</i> (Desr.) Barlow & Wiens
19	Lythraceae	<i>Cuphea dipetala</i> (L.f.) Koehne
20	Lythraceae	<i>Cuphea ericoides</i> Cham. & Schltld.
21	Lythraceae	<i>Cuphea melvilla</i> Lindl.
22	Lythraceae	<i>Lafoensia glyptocarpa</i> Koehne
23	Lythraceae	<i>Lafoensia</i> sp.
24	Malvaceae	<i>Abutilon</i> aff. <i>regnellii</i> Miq.
25	Malvaceae	<i>Abutilon bedfordianum</i> (Hook.) A.St.-Hil.
26	Malvaceae	<i>Abutilon darwinii</i> Hook.f.
27	Malvaceae	<i>Abutilon</i> sp.
28	Malvaceae	<i>Ceiba aesculifolia</i> (Kunth) Britten & Baker f.
29	Malvaceae	<i>Ceiba glaziovii</i> (Kuntze) K.Schum.
30	Malvaceae	<i>Ceiba pentandra</i> (L.) Gaertn.
31	Malvaceae	<i>Ceiba speciosa</i> (A.St.-Hil.) Ravenna
32	Malvaceae	<i>Dombeya wallichii</i> (Lindl.) K.Schum
33	Malvaceae	<i>Eriotheca gracilipes</i> (K.Schum.) A.Robyns
34	Malvaceae	<i>Eriotheca pentaphylla</i> (Vell.) A.Robyns
35	Malvaceae	<i>Guazuma ulmifolia</i> Lam.
36	Malvaceae	<i>Helicteres brevispira</i> A.Juss.
37	Malvaceae	<i>Helicteres eichleri</i> K. Schum.
38	Malvaceae	<i>Helicteres guazumifolia</i> Kunth
39	Malvaceae	<i>Helicteres sacarolha</i> A.Juss.
40	Malvaceae	<i>Helicteres velutina</i> K.Schum.
41	Malvaceae	<i>Hibiscus rosa-sinensis</i> L.
42	Malvaceae	<i>Luehea divaricata</i> Mart.
43	Malvaceae	<i>Malvaceae</i> sp.
44	Malvaceae	<i>Malvaviscus arboreus</i> Cav.
45	Malvaceae	<i>Malvaviscus palmanus</i> Pittier & Donn.Sm.
46	Malvaceae	<i>Melochia tomentosa</i> L.
47	Malvaceae	<i>Pavonia luetzelburgii</i> Ulbr.
48	Malvaceae	<i>Pavonia</i> sp.
49	Malvaceae	<i>Pavonia viscosa</i> A. St.-Hil.
50	Malvaceae	<i>Quararibea lasiocalyx</i> (K.Schum.) Vischer
51		
52		
53		
54		
55		
56		
57		
58		
59		
60		

Family	Plant Species
Malvaceae	<i>Spirotheca rivieri</i> (Decne.) Ulbr.
Malvaceae	<i>Urena lobata</i> L.
Malvaceae	<i>Waltheria cinerescens</i> A. St.-Hil.
Malvaceae	<i>Waltheria rotundifolia</i> Schrank
Marantaceae	<i>Calathea capitata</i> (Ruiz & Pav.) Lindl.
Marantaceae	<i>Calathea gymnocarpa</i> H.A.Kenn.
Marantaceae	<i>Calathea inocephala</i> (Kuntze) T.Durand & B.D.Jacks.
Marantaceae	<i>Calathea lasiostachya</i> Donn.Sm.
Marantaceae	<i>Calathea lutea</i> (Aubl.) E.Mey. ex Schult.
Marantaceae	<i>Calathea zingiberina</i> Körn.
Marantaceae	<i>Ischnosiphon arouma</i> (Aubl.) Körn.
Marantaceae	<i>Ischnosiphon hirsutus</i> Petersen
Marantaceae	<i>Ischnosiphon lasiocoleus</i> K.Schum. ex Loes.
Marantaceae	<i>Maranta furcata</i> Nees & Mart.
Marantaceae	<i>Monotagma secundum</i> (Petersen) K.Schum.
Marantaceae	<i>Stromanthe thalia</i> (Vell.) J.M.A.Braga
Marcgraviaceae	<i>Marcgravia myriostigma</i> Triana & Planch.
Marcgraviaceae	<i>Marcgravia polyantha</i> Delpino
Marcgraviaceae	<i>Norantea guianensis</i> Aubl.
Marcgraviaceae	<i>Sarcopera</i> sp.
Marcgraviaceae	<i>Schwartzia brasiliensis</i> (Choisy) Bedell ex Gir.-Cañas
Melastomataceae	<i>Acinodendron sintenisii</i> (Cogn.) Kuntze
Melastomataceae	<i>Brachyotum ledifolium</i> (Desr.) Triana
Melastomataceae	<i>Brachyotum lindenii</i> Cogn.
Melastomataceae	<i>Brachyotum nutans</i> Gleason
Melastomataceae	<i>Brachyotum</i> sp.
Melastomataceae	<i>Charianthus grenadensis</i> Penneys & Judd
Melastomataceae	<i>Huilaea minor</i> (L. Uribe) Lozano & N. Ruiz
Melastomataceae	<i>Melastomataceae</i> sp.
Melastomataceae	<i>Miconia purpurea</i> (D. Don) Judd & Skean
Melastomataceae	<i>Miconia</i> sp. 1
Melastomataceae	<i>Miconia</i> sp. 2
Musaceae	<i>Musa × paradisiaca</i> L.
Musaceae	<i>Musa</i> sp. 1
Musaceae	<i>Musa</i> sp. 2
Myrtaceae	<i>Callistemon speciosus</i> (Sims) Sweet
Myrtaceae	<i>Eucalyptus</i> sp.
Myrtaceae	<i>Melaleuca leucadendra</i> (L.) L.
Myrtaceae	<i>Myrcia lasiantha</i> DC.
Myrtaceae	<i>Myrcianthes oreophila</i> (Diels) McVaugh
Myrtaceae	<i>Psidium</i> sp.
Myrtaceae	<i>Syzygium jambos</i> (L.) Alston
Myrtaceae	<i>Syzygium malaccense</i> (L.) Merr. & L.M.Perry
Nyctaginaceae	<i>Bougainvillea</i> sp.
Onagraceae	<i>Fuchsia corollata</i> Benth.
Onagraceae	<i>Fuchsia macrostigma</i> Benth.
Onagraceae	<i>Fuchsia microphylla</i> Kunth

1		
2	<b>Family</b>	<b>Plant Species</b>
3		
4	Onagraceae	<i>Fuchsia regia</i> (Vand. ex Vell.) Munz
5	Onagraceae	<i>Fuchsia splendens</i> Zucc.
6	Orchidaceae	<i>Cattleya elongata</i> Barb.Rodr.
7	Orchidaceae	<i>Corymborkis flava</i> (Sw.) Kuntze
8	Orchidaceae	<i>Elleanthus aurantiacus</i> (Lindl.) Rchb.f.
9	Orchidaceae	<i>Elleanthus smithii</i> Schltr.
10	Orchidaceae	<i>Elleanthus</i> sp.
11	Orchidaceae	<i>Orchidaceae</i> sp.
12	Orobanchaceae	<i>Agalinis angustifolia</i> (Mart.) D'Arcy
13	Orobanchaceae	<i>Castilleja angustifolia</i> (Nutt.) G.Don
14	Orobanchaceae	<i>Castilleja fissifolia</i> L.f.
15	Orobanchaceae	<i>Castilleja irasuensis</i> Oerst.
16	Orobanchaceae	<i>Castilleja miniata</i> Douglas ex Hook.
17	Orobanchaceae	<i>Castilleja scorzonerifolia</i> Kunth
18	Orobanchaceae	<i>Castilleja tenuiflora</i> Benth.
19	Orobanchaceae	<i>Esterhazyia splendida</i> J.C.Mikan
20	Orobanchaceae	<i>Vellosiella dracocephaloides</i> (Vell.) Baill.
21	Oxalidaceae	<i>Oxalis psoraleoides</i> Kunth
22	Passifloraceae	<i>Passiflora</i> cf. <i>skiantha</i> Huber
23	Passifloraceae	<i>Passiflora coccinea</i> Aubl.
24	Passifloraceae	<i>Passiflora edmundoi</i> Sacco
25	Passifloraceae	<i>Passiflora kermesina</i> Link & Otto
26	Passifloraceae	<i>Passiflora luetzelburgii</i> Harms
27	Passifloraceae	<i>Passiflora mixta</i> L.f.
28	Passifloraceae	<i>Passiflora quadriglandulosa</i> Rodschied
29	Passifloraceae	<i>Passiflora</i> sp.
30	Passifloraceae	<i>Passiflora speciosa</i> Gardner
31	Passifloraceae	<i>Passiflora spinosa</i> (Poepp. & Endl.) Mast.
32	Passifloraceae	<i>Passiflora vitifolia</i> Kunth
33	Passifloraceae	<i>Passifloraceae</i> sp.
34	Passifloraceae	<i>Turnera ulmifolia</i> L.
35	Phytolaccaceae	<i>Phytolacca rivinoides</i> Kunth & C.D.Bouché
36	Plantaginaceae	<i>Penstemon gentianoides</i> (Kunth) Poir.
37	Plantaginaceae	<i>Penstemon labrosus</i> (A. Gray) Mast. ex Hook. f.
38	Plantaginaceae	<i>Penstemon roseus</i> (Cerv. ex Sweet) G. Don
39	Plumbaginaceae	<i>Plumbago zeylanica</i> L.
40	Polygalaceae	<i>Monnina</i> sp.
41	Polygonaceae	<i>Antigonon leptopus</i> Hook. & Arn.
42	Proteaceae	<i>Grevillea banksii</i> R.Br.
43	Ranunculaceae	<i>Aquilegia formosa</i> Fisch. ex DC.
44	Ranunculaceae	<i>Delphinium barbeyi</i> (Huth) Huth
45	Rosaceae	<i>Rubus rosifolius</i> Sm.
46	Rosaceae	<i>Rubus</i> sp. 1
47	Rosaceae	<i>Rubus</i> sp. 2
48	Rubiaceae	<i>Augusta longifolia</i> (Spreng.) Rehder
49	Rubiaceae	<i>Bouvardia ternifolia</i> (Cav.) Schlttdl.
50	Rubiaceae	<i>Cinchona</i> cf. <i>pitayensis</i> (Wedd.) Wedd.
51		
52		
53		
54		
55		
56		
57		
58		
59		
60		

Family	Plant Species
Rubiaceae	<i>Coussarea hydrangeifolia</i> (Benth.) Benth. & Hook.f. ex Müll.Arg.
Rubiaceae	<i>Erithalis fruticosa</i> L.
Rubiaceae	<i>Faramea eurycarpa</i> Donn.Sm.
Rubiaceae	<i>Ferdinandusa sprucei</i> K.Schum.
Rubiaceae	<i>Genipa americana</i> L.
Rubiaceae	<i>Gonzalagunia hirsuta</i> (Jacq.) K.Schum.
Rubiaceae	<i>Hamelia patens</i> Jacq.
Rubiaceae	<i>Hamelia rostrata</i> Bartl. ex DC.
Rubiaceae	<i>Hillia triflora</i> (Oerst.) C.M.Taylor
Rubiaceae	<i>Hoffmannia arborescens</i> Donn.Sm.
Rubiaceae	<i>Isertia hypoleuca</i> Benth.
Rubiaceae	<i>Isertia parviflora</i> Vahl
Rubiaceae	<i>Isertia rosea</i> Spruce ex K.Schum.
Rubiaceae	<i>Manettia aff. sabiceoides</i> Wernham
Rubiaceae	<i>Manettia coccocypseloides</i> Wernham
Rubiaceae	<i>Manettia cordifolia</i> Mart.
Rubiaceae	<i>Manettia dominicensis</i> Wernham
Rubiaceae	<i>Manettia mitis</i> (Vell.) K.Schum.
Rubiaceae	<i>Manettia pubescens</i> Cham. & Schltld.
Rubiaceae	<i>Morinda citrifolia</i> L.
Rubiaceae	<i>Palicourea anderssoniana</i> C.M.Taylor
Rubiaceae	<i>Palicourea angustifolia</i> Kunth
Rubiaceae	<i>Palicourea aschersonianoides</i> (Wernham) Steyerem.
Rubiaceae	<i>Palicourea cf. anacardiifolia</i> (Humb. & Bonpl. ex Schult.) Standl.
Rubiaceae	<i>Palicourea cf. nigricans</i> K.Krause
Rubiaceae	<i>Palicourea cf. vagans</i> Wernham
Rubiaceae	<i>Palicourea coriacea</i> (Cham.) K.Schum.
Rubiaceae	<i>Palicourea crocea</i> (Sw.) Schult.
Rubiaceae	<i>Palicourea demissa</i> Standl.
Rubiaceae	<i>Palicourea fastigiata</i> Kunth
Rubiaceae	<i>Palicourea gomezii</i> C.M.Taylor
Rubiaceae	<i>Palicourea guianensis</i> Aubl.
Rubiaceae	<i>Palicourea lassiorrhachis</i> Oerst.
Rubiaceae	<i>Palicourea marcgravii</i> A.St.-Hil.
Rubiaceae	<i>Palicourea quadrifolia</i> (Rudge) DC.
Rubiaceae	<i>Palicourea rigida</i> Kunth
Rubiaceae	<i>Palicourea rudgeoides</i> (Müll.Arg.) Standl.
Rubiaceae	<i>Palicourea sodiroi</i> Standl.
Rubiaceae	<i>Palicourea sp. 1</i>
Rubiaceae	<i>Palicourea sp. 2</i>
Rubiaceae	<i>Palicourea subspicata</i> Huber
Rubiaceae	<i>Pentagonia monocalis</i> C.M.Taylor
Rubiaceae	<i>Posoqueria sp.</i>
Rubiaceae	<i>Psychotria cf. nemorosa</i> Gardner
Rubiaceae	<i>Psychotria bahiensis</i> DC.
Rubiaceae	<i>Psychotria berteriana</i> DC.
Rubiaceae	<i>Psychotria capitata</i> Ruiz & Pav.



1		
2	<b>Family</b>	<b>Plant Species</b>
3		
4	Rubiaceae	<i>Psychotria elata</i> (Sw.) Hammel
5	Rubiaceae	<i>Psychotria mapourioides</i> DC.
6	Rubiaceae	<i>Psychotria muscosa</i> (Jacq.) Steyerm.
7	Rubiaceae	<i>Psychotria nuda</i> (Cham. & Schldl.) Wawra
8	Rubiaceae	<i>Psychotria platypoda</i> DC.
9	Rubiaceae	<i>Psychotria poeppigiana</i> Müll.Arg.
10	Rubiaceae	<i>Psychotria suterella</i> Müll.Arg.
11	Rubiaceae	<i>Psychotria urbaniana</i> Steyerm.
12	Rubiaceae	<i>Psychotria vellosiana</i> Benth.
13	Rubiaceae	<i>Retiniphyllum speciosum</i> (Spruce ex Benth.) Müll.Arg.
14	Rubiaceae	<i>Rubiaceae</i> sp.
15	Rubiaceae	<i>Rudgea lorentensis</i> Standl.
16	Rubiaceae	<i>Sabicea aspera</i> Aubl.
17	Rubiaceae	<i>Sabicea villosa</i> Willd. ex Schult.
18	Rubiaceae	<i>Schradera exotica</i> (J.F.Gmel.) Standl.
19	Rubiaceae	<i>Staelia galioides</i> DC.
20	Rubiaceae	<i>Warszewiczia coccinea</i> (Vahl) Klotzsch
21	Rutaceae	<i>Decagonocarpus cornutus</i> R.S.Cowan
22	Rutaceae	<i>Erythrochiton fallax</i> Kallunki
23	Rutaceae	<i>Rutaceae</i> sp.
24	Salicaceae	<i>Ryania pyrifera</i> (Rich.) Uittien & Sleumer
25	Sapindaceae	<i>Serjania aff. caracasana</i> (Jacq.) Willd.
26	Sapindaceae	<i>Serjania coradinii</i> Ferrucci & Somner
27	Sapindaceae	<i>Serjania glutinosa</i> Radlk.
28	Schlegeliaceae	<i>Schlegelia brachyantha</i> Griseb.
29	Schlegeliaceae	<i>Schlegelia fastigiata</i> Schery
30	Scrophulariaceae	<i>Buddleja cf. brasiliensis</i> J.Jacq.
31	Solanaceae	<i>Brugmansia arborea</i> (L.) Steud.
32	Solanaceae	<i>Cestrum corymbosum</i> Schldl.
33	Solanaceae	<i>Cestrum macrophyllum</i> Vent.
34	Solanaceae	<i>Cestrum schlechtendahlil</i> G.Don
35	Solanaceae	<i>Cestrum</i> sp. 1
36	Solanaceae	<i>Cestrum</i> sp. 2
37	Solanaceae	<i>Markea coccinea</i> Rich.
38	Styracaceae	<i>Styrax ferrugineus</i> Nees & Mart.
39	Theaceae	<i>Gordonia fruticosa</i> (Schrad.) H.Keng
40	Tropaeolaceae	<i>Tropaeolum moritzianum</i> Klotzsch
41	Tropaeolaceae	<i>Tropaeolum pentaphyllum</i> Lam.
42	Velloziaceae	<i>Barbacenia blanchetii</i> Goethart & Henrard
43	Velloziaceae	<i>Barbacenia flava</i> Mart. ex Schult. & Schult.f.
44	Velloziaceae	<i>Barbacenia gentianoides</i> Goethart & Henrard
45	Velloziaceae	<i>Barbacenia luzulifolia</i> Mart. ex Schult. & Schult.f.
46	Velloziaceae	<i>Barbacenia williamsii</i> L.B.Sm.
47	Velloziaceae	<i>Vellozia epidendroides</i> Mart.
48	Verbenaceae	<i>Citharexylum spinosum</i> L.
49	Verbenaceae	<i>Lantana camara</i> L.
50	Verbenaceae	<i>Lantana nivea</i> Vent.
51		
52		
53		
54		
55		
56		
57		
58		
59		
60		

Family	Plant Species
Verbenaceae	<i>Stachytarpheta crassifolia</i> Schrad.
Verbenaceae	<i>Stachytarpheta gesnerioides</i> Cham.
Verbenaceae	<i>Stachytarpheta glabra</i> Cham.
Verbenaceae	<i>Stachytarpheta jamaicensis</i> (L.) Vahl
Verbenaceae	<i>Stachytarpheta mexiae</i> Moldenke
Verbenaceae	<i>Stachytarpheta quadrangula</i> Nees & Mart.
Verbenaceae	<i>Stachytarpheta urticifolia</i> (Salisb.) Sims
Violaceae	<i>Paypayrola hulkiana</i> Pulle
Vitaceae	<i>Cissus simsiana</i> Roem. & Schult.
Vitaceae	<i>Cissus verticillata</i> (L.) Nicolson & C.E.Jarvis
Vochysiaceae	<i>Qualea grandiflora</i> Mart.
Vochysiaceae	<i>Qualea multiflora</i> Mart.
Vochysiaceae	<i>Qualea parviflora</i> Mart.
Vochysiaceae	<i>Salvertia convallariaeodora</i> A.St.-Hil.
Vochysiaceae	<i>Vochysia cinnamomea</i> Pohl
Vochysiaceae	<i>Vochysia emarginata</i> Vahl
Vochysiaceae	<i>Vochysia pyramidalis</i> Mart.
Vochysiaceae	<i>Vochysia rufa</i> Mart.
Vochysiaceae	<i>Vochysia</i> sp.
Vochysiaceae	<i>Vochysia tucanorum</i> Mart.
Xanthorrhoeaceae	<i>Phormium tenax</i> J.R.Forst. & G.Forst.
Zingiberaceae	<i>Hedychium coronarium</i> J.Koenig
Zingiberaceae	<i>Renealmia alpinia</i> (Rottb.) Maas
Zingiberaceae	<i>Renealmia breviscapa</i> Poepp. & Endl.
Zingiberaceae	<i>Renealmia cernua</i> (Sw. ex Roem. & Schult.) J.F.Macbr.
Zingiberaceae	<i>Renealmia krukovii</i> Maas
Zingiberaceae	<i>Renealmia pyramidalis</i> (Lam.) Maas
Zingiberaceae	<i>Renealmia sessilifolia</i> Gagnep.
Zingiberaceae	<i>Renealmia</i> sp.
Zingiberaceae	<i>Zingiberaceae</i> sp.

## Hummingbird species

### Phylogenetic lineage Hummingbird Species

Bee	<i>Archilochus colubris</i>
Bee	<i>Atthis heloisa</i>
Bee	<i>Calothorax lucifer</i>
Bee	<i>Chaetocercus mulsant</i>
Bee	<i>Mellisuga helenae</i>
Bee	<i>Mellisuga minima</i>
Bee	<i>Selasphorus calliope</i>
Bee	<i>Selasphorus flammula</i>
Bee	<i>Selasphorus platycercus</i>
Bee	<i>Selasphorus rufus</i>

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60**Phylogenetic lineage Hummingbird Species**


---

Bee	<i>Selasphorus sasin</i>
Bee	<i>Selasphorus scintilla</i>
Brilliant	<i>Aglaeactis castelnaudii</i>
Brilliant	<i>Aglaeactis cupripennis</i>
Brilliant	<i>Boissonneaua flavescens</i>
Brilliant	<i>Boissonneaua jardini</i>
Brilliant	<i>Clytolaema rubricauda</i>
Brilliant	<i>Coeligena helianthea</i>
Brilliant	<i>Coeligena lutetiae</i>
Brilliant	<i>Coeligena prunellei</i>
Brilliant	<i>Coeligena torquata</i>
Brilliant	<i>Coeligena wilsoni</i>
Brilliant	<i>Ensifera ensifera</i>
Brilliant	<i>Eriocnemis cupreovertris</i>
Brilliant	<i>Eriocnemis derbyi</i>
Brilliant	<i>Eriocnemis mosquera</i>
Brilliant	<i>Eriocnemis vestita</i>
Brilliant	<i>Heliodoxa aurescens</i>
Brilliant	<i>Heliodoxa imperatrix</i>
Brilliant	<i>Heliodoxa jacula</i>
Brilliant	<i>Heliodoxa rubinoides</i>
Brilliant	<i>Lafresnaya lafresnayi</i>
Brilliant	<i>Ocreatus underwoodii</i>
Brilliant	<i>Pterophanes cyanopterus</i>
Brilliant	<i>Urosticte benjamini</i>
Coquette	<i>Adelomyia melanogenys</i>
Coquette	<i>Aglaiocercus coelestis</i>
Coquette	<i>Aglaiocercus kingii</i>
Coquette	<i>Chalcostigma herrani</i>
Coquette	<i>Heliangelus amethysticollis</i>
Coquette	<i>Lesbia nuna</i>
Coquette	<i>Lophornis chalybeus</i>
Coquette	<i>Lophornis magnificus</i>
Coquette	<i>Metallura tyrianthina</i>
Coquette	<i>Oreonympha nobilis</i>
Coquette	<i>Oreotrochilus estella</i>
Emerald	<i>Amazilia beryllina</i>
Emerald	<i>Amazilia brevirostris</i>
Emerald	<i>Amazilia candida</i>
Emerald	<i>Amazilia cyanifrons</i>
Emerald	<i>Amazilia cyanocephala</i>
Emerald	<i>Amazilia fimbriata</i>
Emerald	<i>Amazilia franciae</i>
Emerald	<i>Amazilia lactea</i>
Emerald	<i>Amazilia rutila</i>
Emerald	<i>Amazilia tobaci</i>
Emerald	<i>Amazilia tzacatl</i>

	<b>Phylogenetic lineage</b>	<b>Hummingbird Species</b>
1		
2		
3		
4	Emerald	<i>Amazilia versicolor</i>
5	Emerald	<i>Amazilia yucatanensis</i>
6	Emerald	<i>Calliphlox amethystina</i>
7	Emerald	<i>Calliphlox mitchellii</i>
8	Emerald	<i>Campylopterus curvipennis</i>
9	Emerald	<i>Campylopterus hemileucurus</i>
10	Emerald	<i>Campylopterus largipennis</i>
11	Emerald	<i>Chlorestes notata</i>
12	Emerald	<i>Chlorostilbon canivetii</i>
13	Emerald	<i>Chlorostilbon gibsoni</i>
14	Emerald	<i>Chlorostilbon lucidus</i>
15	Emerald	<i>Chlorostilbon maugaeus</i>
16	Emerald	<i>Chlorostilbon mellisugus</i>
17	Emerald	<i>Chlorostilbon olivaresi</i>
18	Emerald	<i>Chlorostilbon poortmani</i>
19	Emerald	<i>Chlorostilbon ricordii</i>
20	Emerald	<i>Chrysuronia oenone</i>
21	Emerald	<i>Cyanophaia bicolor</i>
22	Emerald	<i>Cynanthus latirostris</i>
23	Emerald	<i>Eupetomena macroura</i>
24	Emerald	<i>Eupherusa nigriventris</i>
25	Emerald	<i>Hylocharis chrysur</i>
26	Emerald	<i>Hylocharis cyanus</i>
27	Emerald	<i>Hylocharis leucotis</i>
28	Emerald	<i>Klais guimeti</i>
29	Emerald	<i>Leucochloris albicollis</i>
30	Emerald	<i>Orthorhyncus cristatus</i>
31	Emerald	<i>Phaeochroa cuvieri</i>
32	Emerald	<i>Stephanoxis lalandi</i>
33	Emerald	<i>Thalurania colombica</i>
34	Emerald	<i>Thalurania fannyi</i>
35	Emerald	<i>Thalurania furcata</i>
36	Emerald	<i>Thalurania glaucopis</i>
37	Emerald	<i>Trochilus polytmus</i>
38	Gem	<i>Eugenes fulgens</i>
39	Gem	<i>Heliomaster constantii</i>
40	Gem	<i>Heliomaster squamosus</i>
41	Gem	<i>Lampornis amethystinus</i>
42	Gem	<i>Lampornis calolaema</i>
43	Gem	<i>Lampornis castaneoventris</i>
44	Gem	<i>Lampornis clemenciae</i>
45	Gem	<i>Lampornis hemileucus</i>
46	Gem	<i>Lamprolaima rhami</i>
47	Gem	<i>Panterpe insignis</i>
48	Hermit	<i>Anopetia gounellei</i>
49	Hermit	<i>Eutoxeres aquila</i>
50	Hermit	<i>Glaucis hirsutus</i>
51		
52		
53		
54		
55		
56		
57		
58		
59		
60		

## Phylogenetic lineage Hummingbird Species

---

1		
2		
3		
4	Hermit	<i>Phaethornis bourcierii</i>
5	Hermit	<i>Phaethornis eurynome</i>
6	Hermit	<i>Phaethornis guy</i>
7	Hermit	<i>Phaethornis hispidus</i>
8	Hermit	<i>Phaethornis longirostris</i>
9	Hermit	<i>Phaethornis longuemareus</i>
10	Hermit	<i>Phaethornis malaris</i>
11	Hermit	<i>Phaethornis pretrei</i>
12	Hermit	<i>Phaethornis ruber</i>
13	Hermit	<i>Phaethornis squalidus</i>
14	Hermit	<i>Phaethornis striigularis</i>
15	Hermit	<i>Phaethornis stuarti</i>
16	Hermit	<i>Phaethornis subochraceus</i>
17	Hermit	<i>Phaethornis superciliosus</i>
18	Hermit	<i>Phaethornis syrmatophorus</i>
19	Hermit	<i>Ramphodon naevius</i>
20	Hermit	<i>Threnetes niger leucurus</i>
21	Hermit	<i>Threnetes ruckeri</i>
22	Mango	<i>Anthracothorax dominicus</i>
23	Mango	<i>Anthracothorax nigricollis</i>
24	Mango	<i>Anthracothorax prevostii</i>
25	Mango	<i>Anthracothorax viridis</i>
26	Mango	<i>Augastes lumachella</i>
27	Mango	<i>Augastes scutatus</i>
28	Mango	<i>Chrysolampis mosquito</i>
29	Mango	<i>Colibri coruscans</i>
30	Mango	<i>Colibri delphinae</i>
31	Mango	<i>Colibri serrirostris</i>
32	Mango	<i>Colibri thalassinus</i>
33	Mango	<i>Doryfera ludovicica</i>
34	Mango	<i>Eulampis holosericeus</i>
35	Mango	<i>Eulampis jugularis</i>
36	Mango	<i>Heliactin bilophus</i>
37	Mango	<i>Polytmus guainumbi</i>
38	Topaze	<i>Florisuga fusca</i>
39	Topaze	<i>Florisuga mellivora</i>
40	Topaze	<i>Topaza pyra</i>
41		
42		
43		
44		
45		
46		
47		
48		
49		
50		
51		
52		
53		
54		
55		
56		
57		
58		
59		
60		

## Appendix S2 Phylogenetic methods

### *Plant phylogenies*

For each community we extracted the phylogenetic relatedness of plants from Phylomatic's megatree (phylodiversity.net\phylomatic). Branch lengths were thereafter estimated with the *bladj* function from Phylocom (Webb *et al.*, 2008), using calibration dates from Wikström *et al.* (2001).

### *Hummingbird phylogenies*

For each community we extracted the phylogenetic relatedness of hummingbirds from the most complete to date hummingbird phylogeny, published in McGuire *et al.* (2014), which uses a time-calibrated, multi-locus Bayesian phylogenetic estimate constructed from DNA samples representing 284 hummingbird species.

The phylogenetic tree of McGuire *et al.* (2014) did not include eight hummingbird species represented in our networks, for which we had to approximate phylogenetic distances. Approximations were evaluated individually for each species, aiming to find the most accurate and conservative distance value in each case:

- Values for *Schistes geoffroy* were substituted for those of their most closely related species present in the phylogenetic tree: *Augastes lumachella* and *A. scutatus*.
- Values for *Amazilia saucerrottei* were substituted for those of their most closely related species present in the phylogenetic tree: *A. cyanifrons*.
- Values for *Campylopterus excellens* were substituted for those of their most closely related species present in the phylogenetic tree: *C. curvipennis*.
- Values for *Chlorostilbon mellisugus* were substituted for those of their most closely related species present in the phylogenetic tree: *C. gibsoni*.
- We loosely placed *Phaethornis gounellei* in the base of its clade using the minimum

1  
2 value among *Phaethornis* species.

- 3  
4 – We placed *P. stuarti* similarly using the minimum value of the “Pygmornis” clade within  
5  
6 *Phaethornis*.  
7  
8  
9 – Values for *Chlorostilbon olivaresi* and *Mellisuga helenae* were assigned as the average  
10  
11 distance from the con-genera species present in the phylogenetic tree.  
12  
13

### 14 15 16 **Analysis of phylogenetic signal on bipartite interaction networks**

17  
18 Following the methodology described in Ives & Godfray (2006), we examined whether closely  
19  
20 related species interact with a similar array of partners. In other words, we quantified the  
21  
22 effect of phylogeny on species interaction pattern. To do so, first, for each community and  
23  
24 both for plant and hummingbirds, we used the above described phylogenies to calculate  
25  
26 variance-covariance matrices (vcv), which represent the phylogenetic distance among  
27  
28 species. Variance-covariance matrices were constructed using the “vcv” function from the R  
29  
30 package *ape* (Paradis *et al.*, 2004). This approach assumes that the interaction frequency  
31  
32 between plants and hummingbirds is given by the product of the vcv matrices. Therefore, for a  
33  
34 given plant species *i*, the phylogenetic correlation between visits from hummingbirds *a* and *b*  
35  
36 is given by the hummingbird phylogeny, and *vice versa* for hummingbirds. If both plants and  
37  
38 hummingbirds are closely-related, visits between plant *i* and hummingbird *a* and plant *j* and  
39  
40 hummingbird *b* are correlated. The strength of the phylogenetic association between plants  
41  
42 and hummingbirds was then modeled statistically by fitting vcv matrices to each observed  
43  
44 bipartite interaction matrix through a linear model using the “*pbIm*” function from the R  
45  
46 package *picante* (Kembel *et al.*, 2010). This analysis results in a parameter *d* which  
47  
48 determines the strength of the phylogenetic signal on species' interaction pattern. Specifically,  
49  
50 this analysis provides two independent measures of phylogenetic signal (*d*), one for each  
51  
52 trophic group ( $d_{plants}$  and  $d_{hummingbirds}$  in our case), calculated statistically through an estimated  
53  
54  
55  
56  
57  
58  
59  
60

1  
2 generalized least squares model (EGLS), which does not assume that phylogenetic distance  
3  
4 nor the interaction frequencies are normally distributed. Values of  $d=0$  represent a lack of  
5  
6 phylogenetic correlation, whereas  $d=1$  represents a maximum correlation of phylogenetic  
7  
8 signals. Disruptive selection processes show values above 1 (Ives & Godfray, 2006). We  
9  
10 truncated the  $d$ -value to a maximum of  $d=2$ , as  $d_{plants}$  values for network 11 (Cuba) had an  
11  
12 initial value of 6.43, being clearly an outlier (Table Appendix S2). Confidence intervals for  $d$   
13  
14 were calculated by bootstrap replication data from standardized residuals through the same  
15  
16 EGLS used to fit the original data over 2000 replication sets. Thereafter, we calculated the  
17  
18 overall strength of the phylogenetic signal on the interaction matrix by evaluating three  
19  
20 different models, one assuming no phylogenetic signal ( $d_{plants}=d_{hummingbirds}=0$ ; Star model), one  
21  
22 assuming a maximum phylogenetic signal ( $d_{plants}=d_{hummingbirds}=1$ ; Brownian model), and a final  
23  
24 one which incorporated the observed phylogenetic signals combined (estimated  $d_{plants}$  and  
25  
26  $d_{hummingbirds}$ ; Data model). Again, these models were fitted through an EGLS approximation. To  
27  
28 choose the model which best describes the data we quantified the goodness of fit of each  
29  
30 model as mean squared error (MSE), so that the lowest value of MSE indicate the best fit.  
31  
32  
33  
34  
35  
36

37 Notice however that this analysis should not be acknowledged as a test for particular  
38  
39 phylogenetic hypothesis, but rather a measure of how much does the phylogenetic  
40  
41 relatedness among species affect their interaction pattern.  
42  
43  
44  
45

#### 46 REFERENCES

- 47 Ives, A.R. & Godfray, H.C.J. (2006) Phylogenetic analysis of trophic associations. *American Naturalist*, **168**, E1–  
48 E14.  
49 Kembel, S.W., Cowan, P.D., Helmus, M.R., Cornwell, W.K., Morlon, H., Ackerly, D.D., et al. (2010) Picante: R  
50 tools for integrating phylogenies and ecology. *Bioinformatics*, **26**, 1463–1464.  
51 McGuire, J.A., Witt, C.C., Remsen, J.V., Jr., Corl, A., Rabosky, D.L., Altshuler, D.L., et al. (2014) Molecular  
52 phylogenetics and the diversification of hummingbirds. *Current Biology*, **24**, 910–916.  
53 Paradis, E., Claude, J., & Strimmer, K. (2004) APE: analyses of phylogenetics and evolution in R language.  
54 *Bioinformatics*, **20**, 289–290.  
55 Webb, C.O., Ackerly, D.D. & Kembel, S.W. (2008) Phylocom: software for the analysis of phylogenetic  
56 community structure and trait evolution. *Bioinformatics*, **24**, 2098–2100.  
57 Wikström, N., Savolainen, V., & Chase, M.W. (2001). Evolution of the angiosperms: Calibrating the family tree.  
58 *Proceedings of the Royal Society of London B*, **268**, 2211–2220.  
59  
60



Appendix S2 Table A. Results of the phylogenetic analysis for each community

Dataset ID number	MSE <sub>Dat</sub>		MSE <sub>Brownia</sub> n	CI MIN		CI MAX	CI MIN		CI MAX
	a	MSE <sub>Star</sub>		d <sub>plants</sub>	d <sub>plants</sub>		d <sub>hummingbir</sub> ds	d <sub>hummingbir</sub> ds	
1	15.468	19.632	36.420	0.000	0.094	0.452	0.002	0.243	0.467
2	9.887	13.973	16.665	0.000	0.293	0.639	0.000	0.310	2.395
3	25.700	25.277	183.797	0.955	0.883	1.056	0.000	0.000	0.114
4	14.194	15.611	34.123	0.000	0.161	0.403	0.000	0.043	0.257
5	54.965	59.220	101.878	0.000	0.263	0.642	0.000	0.092	0.330
6	2.270	4.220	5.354	0.354	0.700	1.018	0.000	0.000	0.156
7	109.205	116.143	207.956	0.000	0.252	0.521	0.000	0.053	0.284
8	69.545	78.534	122.920	0.000	0.093	0.364	0.000	0.295	0.574
9	16.398	21.877	40.001	0.181	0.587	1.009	0.000	0.000	0.119
10	139.116	139.116	336.877	0.000	0.000	0.000	0.000	0.000	0.000
11	137.292	247.429	242.988	0.931	2.000	17.038	0.000	0.098	0.765
12	6.258	7.458	13.106	0.034	0.654	1.280	0.000	0.000	0.293
13	36.780	37.552	55.636	0.000	0.041	0.637	0.000	0.119	0.787
14	37.913	38.519	60.741	0.000	0.041	1.873	0.000	0.120	1.400
15	16.017	16.433	25.504	0.000	0.151	2.000	0.000	0.000	0.838
16	1.454	1.642	2.379	0.068	0.290	0.499	0.000	0.155	0.319
17	1.017	1.577	1.124	0.089	1.056	2.452	0.000	0.261	1.888
18	4.161	4.471	5.857	0.118	0.706	1.387	0.000	0.065	0.377
19	12.602	21.129	18.718	0.042	1.119	3.076	0.000	0.060	0.514
20	28.627	28.718	43.951	0.000	0.000	0.734	0.000	0.000	0.458
21	1.552	2.167	2.627	0.032	0.383	0.804	0.057	0.310	0.520
22	1.761	2.415	3.948	0.268	1.703	5.247	0.000	0.000	0.291
23	1.490	1.785	1.933	0.000	0.000	0.877	0.000	0.396	0.706
24	63.498	67.344	119.495	0.071	0.420	0.841	0.000	0.006	0.268
25	9.717	9.889	21.005	0.000	0.046	0.232	0.000	0.052	0.213
26	7.456	7.581	15.548	0.000	0.045	0.396	0.000	0.000	0.237
27	2.885	3.426	3.618	0.168	0.397	0.599	0.000	0.292	0.779
28	41.405	42.071	63.699	0.000	0.112	0.423	0.000	0.053	0.306
29	15.717	18.415	29.404	0.025	0.611	1.223	0.000	0.000	0.246
30	52.055	56.779	132.613	0.000	0.009	0.331	0.000	0.084	0.413
31	3.738	4.052	9.265	0.057	0.306	0.546	0.000	0.000	0.095
32	2.873	3.296	4.361	0.056	0.271	0.455	0.056	0.267	0.436
33	0.603	0.723	0.865	0.000	0.066	0.379	0.026	0.405	0.674
34	2.385	2.397	5.426	0.000	0.000	0.186	0.000	0.000	0.130
35	12.454	19.826	23.805	0.257	1.150	2.362	0.000	0.013	0.330
36	6.404	7.762	7.988	0.000	0.240	0.827	0.000	0.499	1.383
37	10.555	10.837	13.897	0.000	0.284	2.633	0.000	0.000	1.158
38	6.022	6.340	11.381	0.007	0.214	0.430	0.000	0.028	0.197
39	23.433	23.874	44.968	0.000	0.000	0.272	0.000	0.136	0.390
40	1.956	2.205	3.601	0.132	0.350	0.551	0.000	0.000	0.142
41	1.616	2.121	2.510	0.108	0.235	0.359	0.266	0.409	0.514
42	1.716	2.047	2.875	0.000	0.009	0.296	0.047	0.388	0.607
43	1.075	1.083	2.293	0.000	0.045	0.178	0.000	0.000	0.085
44	1.783	2.562	2.289	0.157	0.668	1.275	0.001	0.340	0.609
45	1.165	1.168	1.778	0.000	0.000	0.150	0.001	0.143	0.316

Dataset ID number	MSE <sub>Dat</sub>		MSE <sub>Brownia</sub>	CI MIN	d <sub>plant</sub>	CI MAX	CI MIN	d <sub>hummingbir</sub>	CI MAX	d <sub>hummingbir</sub>
	a	MSE <sub>Star</sub>	n	d <sub>plants</sub>	s	d <sub>plants</sub>	ds	ds	ds	ds
46	0.931	0.935	2.388	0.000	0.000	2.800	0.000	0.008	0.412	
47	0.472	0.493	1.565	0.000	0.426	1.899	0.000	0.000	0.470	
48	10.432	11.690	16.730	0.000	0.033	0.961	0.000	0.350	0.971	
49	134.072	149.850	165.457	0.174	0.553	0.932	0.000	0.255	0.519	
50	0.649	0.701	3.976	0.000	0.310	0.871	0.000	0.000	0.159	
51	4.974	6.059	6.535	0.000	0.107	0.476	0.102	0.547	0.798	
52	1.289	1.600	1.505	0.182	0.427	0.765	0.204	0.397	0.614	
53	11.288	11.288	35.158	0.000	0.000	0.228	0.000	0.000	0.186	
54	0.797	0.821	1.811	0.000	0.144	0.340	0.000	0.000	0.132	

For Peer Review

## Appendix S3 Effect of the differences in sampling effort among data sets on network structure

Network data were collected from various studies employing different sampling efforts, which could potentially bias resulting differences in network structure among communities. We measured sampling effort as sampling intensity, i.e. the ratio between number of interactions and the geometric mean of the number of plants and hummingbirds in the community as in Schleuning et al. (2014). Simple correlations presented in Table S2 show a relationship between sampling intensity and corrected complementary specialization and with observed modularity, both in the global and mainland datasets. Therefore, in order to reveal whether these correlations had any repercussions in the conclusions obtained from the macroecological models, we repeated the macroecological models presented in the main manuscript including sampling intensity as a predictor variable (Table Appendix S3).

Overall trends remained constant. For corrected complementary specialization ( $H_2'$ ), the inclusion of sampling intensity had no overall effects on the other predictor variables in the global model, while in the mainland analysis network size lost some of its importance, but the rank of importance remained the same. For observed modularity, the only difference was that  $d_{\text{hummingbirds}}$  lost some of its importance in the global dataset, while insularity became an important predictor, with islands showing a lower modularity than mainland communities (Table Appendix S3), confirming findings from other performed complementary macroecological analysis (Appendix S4) which suggest differences between mainland and island communities.

### REFERENCES

Schleuning, M., Ingmann, L., Strauss, R., Fritz, S.A., Dalsgaard, B., Matthias Dehling, D., et al. (2014) Ecological, historical and evolutionary determinants of modularity in weighted seed-dispersal networks. *Ecology Letters*, 17, 454–463.

**Appendix S3 Table A.** Relationship between species richness, phylogenetic signal and environment, and corrected complementary specialization ( $H_2'$ ) and observed modularity (QuanBiMo) across the Americas (Global dataset) and the mainland, i.e. when excluding insular communities (Only Mainland), when including Sampling effort as a predictor variable, as these network metrics appeared to be correlated with sampling intensity (Table S2). For each network metric, the standardized regression coefficients are reported for ordinary least squares regression for a model including all variables ("Full") and for a model averaged across all possible models using Akaike Weights (AICc  $w_i$  - "Averaged"). For the averaged models, the relative importance of each predictor variable (" $\Sigma w_i$ ") is given by the Akaike Weights (AICc  $w_i$ ). Predictors with high importance are marked in bold. Phylogenetic signal among plants and hummingbirds,  $d_p$  and  $d_h$ , respectively; Mean annual temperature, MAT; Temperature seasonality, TS; Mean annual precipitation, MAP; Precipitation seasonality, PS; Velocity of temperature change since the last Glacial Maximum, MAT velocity; Velocity of precipitation change since the last Glacial Maximum, MAP velocity. The significance of Moran's  $I$  was tested with 10 distance classes and a permutation test with 10,000 iterations. In all models multicollinearity was not an issue, i.e.  $VIF \leq 3.3$ ,  $CN \leq 4.2$ , and there was no positive spatial autocorrelation, i.e. Moran's  $I$  was non-significant.

		Model	AICc	$R^2$	Species richness	Phylogenetic signal		Environmental factors						Sampling effort		
					Network size	$d_p$	$d_h$	Insularity	MAT	TS	MAP	PS	MAT velocity	MAP velocity	Sampling intensity	
Global dataset (54 networks)	$H_2'$ corrected	Full	-60.862	0.620	<b>0.346</b>	-0.027	<b>-0.228</b>	-0.311	0.285	0.138	0.237	-0.165	-0.485	0.140	<b>0.341</b>	
		Averaged			<b>0.415</b>	-0.089	<b>-0.297</b>	-0.245	0.200	0.103	0.305	-0.206	-0.289	0.069	<b>0.342</b>	
		$\Sigma w_i$			<b>0.956</b>	0.264	<b>0.868</b>	0.521	0.385	0.272	0.646	0.516	0.734	0.254	<b>0.944</b>	
	QuanBiMo observed	Full	-69.765	0.658	<b>0.371</b>	-0.170	-0.181	<b>-0.373</b>	0.202	-0.044	0.138	-0.087	-0.052	-0.085	<b>-0.289</b>	
		Averaged			<b>0.380</b>	-0.202	-0.225	<b>-0.345</b>	0.192	-0.071	0.305	-0.098	-0.063	-0.091	<b>-0.304</b>	
		$\Sigma w_i$			<b>0.953</b>	0.653	0.659	<b>0.886</b>	0.447	0.235	0.710	0.279	0.252	0.266	<b>0.872</b>	
	Only Mainland (45 networks)	$H_2'$ corrected	Full	-59.753	0.652	0.212	-0.045	<b>-0.281</b>	na	<b>0.478</b>	0.091	0.114	-0.223	<b>-0.708</b>	0.240	<b>0.344</b>
			Averaged			0.249	-0.057	<b>-0.305</b>	na	<b>0.512</b>	0.044	0.208	-0.266	<b>-0.680</b>	0.277	<b>0.370</b>
			$\Sigma w_i$			0.641	0.195	<b>0.871</b>	na	<b>0.961</b>	0.191	0.340	0.736	<b>0.985</b>	0.614	<b>0.956</b>
QuanBiMo observed		Full	-54.411	0.619	<b>0.262</b>	-0.217	<b>-0.280</b>	na	0.313	-0.118	0.052	-0.103	-0.184	-0.075	<b>-0.311</b>	
		Averaged			<b>0.309</b>	-0.239	<b>-0.292</b>	na	0.287	-0.168	0.327	-0.105	-0.221	-0.086	<b>-0.347</b>	
		$\Sigma w_i$			<b>0.756</b>	0.694	<b>0.754</b>	na	0.583	0.356	0.549	0.262	0.373	0.237	<b>0.809</b>	

## Appendix S4 Results of macroecological models on observed network metrics

As an additional analysis, we performed an equivalent macroecological analysis on observed network metrics instead of on null model corrected metrics as presented in the main manuscript. By analyzing observed network metrics, we are investigating the "raw" level of specialization, regardless of the community context (species and interaction richness, interaction frequency distribution). This complementary analysis allow us to understand more fully the effect of correcting for network metrics and the relationships between the studied variables (Tables S4A and S4B).

The amount of variability explained by the models is similar to that of corrected metrics, but important differences in the role of the different predictor variables arose. First of all, species richness lost its high importance except for modularity in the global dataset. In other words, in general, larger communities did not show higher levels of observed specialization (complementary specialization and modularity). Together with the results from the corrected-metrics analysis, where species richness exhibited a strong and widespread association with higher levels of specialization, we can conclude that *only when compared to communities with the same attributes as empirical matrices* do species richness impact network structure.

Similarly, hummingbird phylogenetic signal, the other variable exhibiting a constant association with corrected network structure, is only related with complementary specialization, and not modularity, when examining observed metrics. As our null model correction does not account for species' phylogenetic relatedness, we may conclude that higher levels of corrected modularity are indeed achieved when closely-related hummingbirds form distinct modules.

As with corrected metrics, mean annual temperature and temperature velocity were related only to higher levels of complementary specialization on mainland communities, both

1 showing the highest regression coefficients of all predictors in the full models.  
2

3  
4 Additionally, insularity and mean annual precipitation became important determinants  
5 of observed specialization, with mainland and humid communities showing higher levels of  
6 complementary specialization and modularity than insular communities or networks from drier  
7 environments. The effect of these predictor variables disappeared when modeling corrected  
8 metrics, suggesting that insularity and MAP may affect either species/interaction richness or  
9 interaction distribution, leading to higher observed levels of specialization (Dalsgaard *et al.*,  
10 2011).  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20

## 21 REFERENCES

22 Dalsgaard, B., Magård, E., Fjeldså, J., Martín González, A.M., Rahbek, C., Olesen, J.M., et al. (2011)  
23 Specialization in plant-hummingbird networks is associated with species richness, contemporary  
24 precipitation and quaternary climate-change velocity. *PLoS ONE*, **6**, e25891.  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Appendix S4 Table A.** Relationship between species richness, phylogenetic signal and environment, and complementary specialization ( $H_2'$ ) and modularity (QuanBiMo) across the Americas (Global dataset) and the mainland, i.e. when excluding insular communities (Only Mainland). Complementary specialization measures the exclusiveness of the interactions in the community, whereas modularity describes a community pattern where species interact more frequently within subsets of species and loosely among subsets. For each network metric, the standardized regression coefficients are reported for ordinary least squares regression for a model including all variables ("Full") and for a model averaged across all possible models using Akaike Weights (AICc  $w_i$  - "Averaged"). For the averaged models, the relative importance of each predictor variable (" $\Sigma w_i$ ") is given by the Akaike Weights (AICc  $w_i$ ). Phylogenetic relatedness among plants and hummingbirds,  $d_p$  and  $d_h$ , respectively; Mean annual temperature, MAT; Temperature seasonality, TS; Mean annual precipitation, MAP; Precipitation seasonality, PS; Velocity of temperature change since the last Glacial Maximum, MAT velocity; Velocity of precipitation change since the last Glacial Maximum, MAP velocity. The significance of Moran's  $I$  was tested with 10 distance classes and a permutation test with 10,000 iterations. In all models multicollinearity was not an issue, i.e. VIF  $\leq$  4.0, CN  $\leq$  3.3, and there was no positive spatial autocorrelation, i.e. Moran's  $I$  was non-significant.

		Species richness			Phylogenetic signal			Environmental factors						
		Model	AICc	R <sup>2</sup>	Network size	$d_p$	$d_h$	Insularity	MAT	TS	MAP	PS	MAT velocity	MAP velocity
Global dataset (54 networks)	$H_2'$	Full	-35.112	0.578	0.210	-0.069	<b>-0.231</b>	<b>-0.453</b>	0.355	0.085	<b>0.344</b>	-0.085	-0.317	0.129
		Averaged			0.249	-0.119	<b>-0.254</b>	<b>-0.401</b>	0.290	0.040	<b>0.423</b>	-0.101	-0.225	0.115
		$\Sigma w_i$			0.659	0.323	<b>0.767</b>	<b>0.916</b>	0.573	0.218	<b>0.965</b>	0.268	0.417	0.295
	QuanBiMo	Full	-65.124	0.602	<b>0.273</b>	-0.119	-0.117	<b>-0.449</b>	0.203	-0.076	<b>0.301</b>	-0.049	-0.037	-0.128
		Averaged			<b>0.259</b>	-0.139	-0.112	<b>-0.446</b>	0.202	-0.098	<b>0.446</b>	-0.028	-0.063	-0.095
		$\Sigma w_i$			<b>0.794</b>	0.411	0.324	<b>0.993</b>	0.483	0.276	<b>0.989</b>	0.216	0.261	0.280
Only Mainland (45 networks)	$H_2'$	Full	-31.253	0.574	0.076	-0.067	<b>-0.300</b>	na	<b>0.518</b>	0.001	0.238	-0.105	<b>-0.491</b>	0.197
		Averaged			0.145	-0.094	<b>-0.382</b>	na	<b>0.525</b>	-0.065	0.327	-0.167	<b>-0.475</b>	0.223
		$\Sigma w_i$			0.325	0.250	<b>0.858</b>	na	<b>0.924</b>	0.223	0.731	0.351	<b>0.832</b>	0.477
	QuanBiMo	Full	-51.864	0.561	0.159	-0.221	-0.234	na	0.314	-0.149	<b>0.232</b>	-0.069	-0.179	-0.136
		Averaged			0.181	-0.232	-0.210	na	0.298	-0.181	<b>0.455</b>	-0.047	-0.230	-0.101
		$\Sigma w_i$			0.412	0.637	0.496	na	0.585	0.383	<b>0.938</b>	0.221	0.384	0.261

**Appendix S4 Table B.** Coefficients of determination ( $R^2$ ) from partial regression local models separating the effect of species richness, species' evolutionary history and environmental factors as predictors of corrected network structure of American communities, when including climate change velocity as an estimate of historical climate stability. We estimated  $R^2$  values in "full" models, i.e. in models including all ten predictor variables (see Appendix S4 Table A). "Total model" reflects the total amount of variation in network structure explained by the full model; "Total species richness", "Total phylogenetic signals" and "Total environmental factors" reflect the variation explained by species richness, phylogenetic signal and environmental factors, respectively; "Only species richness", "Only phylogenetic signal" and "Only environmental factors" reflect the amount of variation in network structure explained by the specified predictor-type and not by any other (i.e. the unique variation); Finally, "shared" reflects the variation attributable to the pair of specified predictor types or to all factors combined. See Table S4A for standardized coefficients of each variable and more details of model fit.

	Global dataset (54 networks)		Only Mainland (45 networks)	
	$H_2'$ observed	QuanBiMo observed	$H_2'$ observed	QuanBiMo observed
Total model	0.58	0.60	0.57	0.56
Total species richness	0.19	0.27	0.11	0.14
Total phylogenetic signal	0.23	0.18	0.25	0.23
Total environmental factors	0.52	0.54	0.51	0.48
Unique species richness	0.03	0.04	0.00	0.02
Unique phylogenetic signal	0.04	0.02	0.06	0.07
Unique environmental factors	0.19	0.21	0.23	0.23
Shared species richness – phylogeny	0.00	0.00	0.00	0.00
Shared species richness – environmental factors	0.14	0.17	0.09	0.08
Shared phylogeny – environmental factors	0.16	0.11	0.17	0.12
Shared all factors	0.04	0.06	0.02	0.05



## Appendix S5 Complementary macroecological analyses

As different climate change metrics may capture different aspects important for biodiversity (Garcia *et al.*, 2014), to evaluate more comprehensively the impact of historical climate-change we calculated another summarizations of historical climatic stability: *the anomaly of mean annual temperature and mean annual precipitation* between the LGM and the present, calculated as pre-industrial minus LGM precipitation/temperature (i.e. positive values indicate that conditions were drier or colder at LGM than at present). Whereas climate change velocities reflect the speed of climate change irrespective of the direction of the change, anomalies reflect the direction of climate change. As with climate change velocities, projections of the global climate during the LGM and pre-industry were generated by Hadley Centre Coupled Model Version 3 (HadCM3) with a resolution of  $3.75 \times 2.5$  arc degrees (Singarayer & Joy 2010), and were downscaled to  $0.1 \times 0.1$  arc degrees.

We performed a set of models equivalent to those presented in the main text (Diniz-Filho *et al.* 2008) but which included climate change anomaly instead of climate change velocity. Temperature anomaly was  $\log_{10}$ -transformed. Models including climatic anomaly gave a poorer fit to our data, explaining less variability and raising multicollinearity concerns. Nevertheless, strong patterns disclosed by the macroecological models with climatic velocity persisted (Tables S4A-B).

The significance and importance of species richness and hummingbird phylogenetic signal was again widespread and showed similar values than the models which included change velocity as measure of historical climate stability. We found a stronger role, although still non-significant, of the phylogenetic signal of plants, associated with lower levels of modularity on both datasets. Finally, and most importantly, neither temperature nor precipitation anomaly had any effect on any of the studied variables.

**REFERENCES**

- Diniz-Filho, J.A.F., Rangel, T.F. L. & Bini, L.M. (2008) Model selection and information theory in geographical ecology. *Global Ecology and Biogeography*, **17**, 479–488.
- Garcia, R.A., Cabeza, M., Rahbek, C., & Araújo, M.B. (2014) Multiple Dimensions of Climate Change and Their Implications for Biodiversity. *Science*, **344**, 6183.
- Singarayer, J.S., & Valdes, P. J. (2010) High-latitude climate sensitivity to ice-sheet forcing over the last 120 kyr. *Quaternary Science Reviews*, **29**, 43–55.

For Peer Review

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Appendix S5 Table A.** Relationship between species richness, phylogenetic signal and environment, and complementary specialization ( $H_2'$ ) and modularity (QuanBiMo) across the Americas (Global dataset) and the mainland, i.e. when excluding insular communities (Only Mainland), when including historical climate anomaly as a predictor of historical climatic stability. Complementary specialization measures the exclusiveness of the interactions in the community, whereas modularity describes a community pattern where species interact more frequently within subsets of species and loosely among subsets. For each network metric, the standardized regression coefficients are reported for ordinary least squares regression for a model including all variables ("Full") and for a model averaged across all possible models using Akaike Weights (AICc  $w_i$  - "Averaged"). For the averaged models, the relative importance of each predictor variable (" $\Sigma w_i$ ") is given by the Akaike Weights (AICc  $w_i$ ). Phylogenetic signal among plants and hummingbirds,  $d_p$  and  $d_h$ , respectively; Mean annual temperature, MAT; Temperature seasonality, TS; Mean annual precipitation, MAP; Precipitation seasonality, PS; Velocity of temperature change since the last Glacial Maximum, MAT velocity; Velocity of precipitation change since the last Glacial Maximum, MAP velocity. The significance of Moran's  $I$  was tested with 10 distance classes and a permutation test with 10,000 iterations. In all models [multicollinearity was not an issue](#), i.e.  $VIF \leq 5.1$ ,  $CN \leq 4.9$ , and [there was no positive spatial autocorrelation](#), i.e. Moran's  $I$  was non-significant.

		Species richness			Phylogenetic signal			Environmental factors						
	Model	AICc	$R^2_{adj}$	Network size	$d_p$	$d_h$	Insularity	MAT	TSN	MAP	PSN	MAT anomaly	MAP anomaly	
Global dataset (54 networks)	$H_2'$	Full	-46.828	0.474	0.512	-0.168	-0.306	-0.046	-0.004	0.111	0.026	-0.251	0.009	-0.046
		Averaged			0.516	-0.165	-0.328	-0.067	-0.024	0.080	0.101	-0.249	0.004	-0.023
		$\Sigma w_i$			0.999	0.458	0.936	0.240	0.218	0.261	0.276	0.738	0.216	0.224
	QuanBiMo	Full	-99.080	0.619	0.630	-0.186	-0.237	-0.210	0.110	0.075	-0.087	-0.242	0.022	-0.217
		Averaged			0.636	-0.194	-0.277	-0.141	0.024	0.054	0.118	-0.214	-0.053	-0.124
		$\Sigma w_i$			1.000	0.667	0.908	0.362	0.227	0.239	0.317	0.696	0.242	0.343
Only Mainland (45 networks)	$H_2'$	Full	-45.189	0.477	0.422	-0.146	-0.337	na	0.135	0.019	-0.388	-0.350	0.216	-0.215
		Averaged			0.424	-0.116	-0.330	na	0.079	0.039	-0.067	-0.275	-0.218	-0.047
		$\Sigma w_i$			0.968	0.282	0.839	na	0.237	0.217	0.246	0.695	0.479	0.229
	QuanBiMo	Full	-76.043	0.558	0.539	-0.217	-0.301	na	0.169	0.028	-0.285	-0.291	-0.081	-0.332
		Averaged			0.534	-0.192	-0.293	na	0.101	0.019	0.079	-0.251	-0.135	-0.183
		$\Sigma w_i$			0.999	0.510	0.794	na	0.265	0.209	0.282	0.676	0.312	0.410

**Appendix S5 Table B.** Coefficients of determination ( $R^2$ ) from partial regression local models separating the effect of species richness, environmental factors and species' evolutionary history as predictors of corrected network structure of American communities when using climatic anomaly as an estimate for historical climatic stability. We estimated  $R^2$  values in “full” models, i.e. in models including all ten predictor variables (see Appendix S5 Table A for standardized coefficients of each variable and more details of model fit). “Total model” reflects the total amount of variation in network structure explained by the model; “Total species richness”, “Total phylogenetic signals” and “Total environmental factors” reflect the total variation explained by species richness, phylogenetic signal and environmental factors, respectively, whereas “Only species richness”, “Only phylogenetic signal” and “Only environmental factors” reflect the unique variation in network structure explained by species richness, phylogenetic signal and environmental factors, respectively. Finally, “shared” reflect the shared variation described by the specified predictor sets, and “shared all factors” reflect variation attributable to species richness, phylogenetic signal and environmental factors.

	Global dataset (54 networks)		Only Mainland (45 networks)	
	$H_2'$	QuanBiMo	$H_2'$	QuanBiMo
Total model	0.47	0.62	0.48	0.56
Total species richness	0.25	0.44	0.20	0.33
Total phylogenetic signal	0.19	0.19	0.18	0.19
Total environmental factors	0.25	0.32	0.28	0.28
Unique species richness	0.16	0.24	0.12	0.20
Unique phylogenetic signal	0.08	0.06	0.08	0.08
Unique environmental factors	0.06	0.06	0.11	0.09
Shared species richness – phylogeny	0.00	0.00	0.00	0.00
Shared species richness – environmental factors	0.07	0.14	0.06	0.09
Shared phylogeny – environmental factors	0.08	0.07	0.08	0.06
Shared all factors	0.04	0.06	0.03	0.05

**Table S1.** Coordinates, description, location and data references for each studied community.

ID number	Latitude	Longitude	Site description and general location	Data Source Reference
1	-20.75	-42.92	Atlantic forest, SE Brazil	Abreu, C.R.M. & Vieira, M.F. (2004) Os beija-flores e seus recursos florais em um fragmento florestal de Viçosa, sudeste brasileiro. <i>Lundiana</i> , <b>5</b> , 129–134.
2	34.22	-116.95	Montane meadow, SW USA	Alarcón, R., Waser, N.M & Ollerton, J. (2008) Year-to-year variation in the topology of a plant-pollinator interaction network. <i>Oikos</i> , <b>117</b> , 1796–1807.
3	38.98	-106.97	Sub-alpine meadow, Central USA	Ollerton, J., Alarcón, R., Waser, N.M, Price, M.V., Watts, S., Cranmer, L., Hingston, A., Peter, C. & Rotenberry, J. (2009) A global test of the pollination syndrome hypothesis. <i>Annals of Botany</i> , <b>103</b> , 1471–1480.
4	-23.35	-44.83	Atlantic forest, SE Brazil	Araujo, A.C. (1996) Beija-flores e seus recursos florais numa área de planície costeira do litoral norte de São Paulo, sudeste do Brasil. <i>MSc. Thesis</i> . Universidade Estadual de Campinas, Brazil.
5	19.50	-105.05	Tropical dry deciduous forest, W Mexico	Arizmendi, M.C. & Ornelas, J.F. (1990) Hummingbirds and their floral resources in a tropical dry forest in Mexico. <i>Biotropica</i> , <b>22</b> , 172–180.
6	-31.80	-52.42	Pampa, S Brazil	Vizentin-Bugoni, J. & Rui, A.M. <i>Unpublished data</i> .
7	-22.50	-44.83	Highland Atlantic forest, SE Brazil	Canela, M.B.F. (2006) Interações entre plantas e beija-flores numa comunidade de floresta atlântica montana em Itatiaia, RJ. <i>Ph.D thesis</i> . Universidade Estadual de Campinas, Brazil.
8	-7.87	-36.40	Caatinga, NE Brazil	Las Casas, F.M.G., Azevedo Júnior, S.M. & Dias Filho, M.M. (2012) The community of hummingbirds (Aves: Trochilidae) and the assemblage of flowers in a Caatinga vegetation. <i>Brazilian Journal of Biology</i> , <b>72</b> , 51–58.
9	-11.48	-41.32	Altitudinal caatinga, NE Brazil	Moura, A.C. & Machado, C.G. Hummingbirds and their flowers at altitudinal dryland vegetation in Chapada Diamantina, northeast Brazil. <i>In preparation</i> .
10	-3.82	-70.27	Amazonian rainforest, SE Colombia	Cotton, P.A. (1998) The hummingbird community of a lowland Amazonian rainforest. <i>Ibis</i> , <b>140</b> , 512–521.
11	22.28	-81.20	Swamp forest, Hurricane disturbed, Cuba	Baquero, A.C. (2014) Evolutionary and ecological insight into hummingbird-plant communities in the Caribbean. <i>MSc Thesis</i> . University of Copenhagen, Denmark.
12	15.35	-61.30	Elfin forest, Dominica	Dalsgaard, B., Martín González, A.M., Olesen, J.M., Ollerton, J., Timmermann, A., Andersen, L.H. & Tossas, A.G. (2009) Plant–hummingbird interactions in the West Indies: floral specialisation gradients associated with environment and hummingbird size. <i>Oecologia</i> , <b>159</b> , 757–766.
13	15.25	-61.37	Coastal dry scrubland, Dominica	Dalsgaard, B., Martín González, A.M., Olesen, J.M., Ollerton, J., Timmermann, A., Andersen, L.H. & Tossas, A.G. (2009) Plant–hummingbird interactions in the West Indies: floral specialisation gradients associated with environment and hummingbird size. <i>Oecologia</i> , <b>159</b> , 757–766.

ID number	Latitude	Longitude	Site description and general location	Data Source Reference
14	12.10	-61.70	Elfin forest, Grenada	Dalsgaard, B., Martín González, A.M., Olesen, J.M., Ollerton, J., Timmermann, A., Andersen, L.H. & Tossas, A.G. (2009) Plant–hummingbird interactions in the West Indies: floral specialisation gradients associated with environment and hummingbird size. <i>Oecologia</i> , <b>159</b> , 757–766.
15	12.10	-61.68	Rainforest, Grenada	Dalsgaard, B., Martín González, A.M., Olesen, J.M., Ollerton, J., Timmermann, A., Andersen, L.H. & Tossas, A.G. (2009) Plant–hummingbird interactions in the West Indies: floral specialisation gradients associated with environment and hummingbird size. <i>Oecologia</i> , <b>159</b> , 757–766.
16	1.25	-77.43	High andean forest, Colombia	Gutierrez Zamora, E.A. & Rojas Nossa, S.V. (2001) Dinámica anual de la interacción colibrí-flor en ecosistemas altoandinos del volcán Galeras, Sur de Colombia. <i>BSc. Thesis</i> . Universidad Nacional de Colombia, Colombia.
17	18.35	-77.65	Evergreen lowland rainforest, Jamaica.	Ingversen, T. (2006) Plant–Pollinator Interactions on Jamaica and Dominica – The Centrality, Asymetry, and Modularity of Networks. <i>MSc. Thesis</i> . University of Aarhus, Denmark.
18	-27.27	-49.01	Atlantic Forest, low elevation	Kohler, G. (2011) Redes de interação planta-beija-flor em um gradiente altitudinal de Floresta Atlântica no Sul do Brasil. <i>MSc. Thesis</i> . Universidade Federal do Paraná, Brazil.
19	-27.26	-49.02	Atlantic Forest, mid elevation	Kohler, G. (2011) Redes de interação planta-beija-flor em um gradiente altitudinal de Floresta Atlântica no Sul do Brasil. <i>MSc. Thesis</i> . Universidade Federal do Paraná, Brazil.
20	-27.26	-49.02	Atlantic Forest, high elevation	Kohler, G. (2011) Redes de interação planta-beija-flor em um gradiente altitudinal de Floresta Atlântica no Sul do Brasil. <i>MSc. Thesis</i> . Universidade Federal do Paraná, Brazil.
21	19.23	-98.97	Highland temperate mosaic forest, Central Mexico	Lara, C. (2006) Temporal dynamics of flower use by hummingbirds in a highland temperate forest in México. <i>Ecoscience</i> , <b>13</b> , 23–29.
22	19.28	-98.23	Suburban forest, Central Mexico	Lara, C. <i>Unpublished data</i> .
23	19.40	-96.80	Protected cloud forest, Central Mexico	Lara, C. <i>Unpublished data</i> .
24	0.07	-72.45	Rainforest, Colombia	Rosero, L. (2003) Interações planta/beija-flor em três comunidades vegetais da parte sul do Parque Nacional Natural Chiribiquete, Amazonas (Colombia). <i>Ph.D. Thesis</i> . Universidade Estadual de Campinas, Brazil. Rosero-Lasprilla, L. & Sazima, M. (2004) Interacciones planta-colibrí en tres comunidades vegetales de la parte suroriental del Parque Nacional Natural Chiribiquete, Colombia. <i>Ornitologia Neotropical</i> , <b>15</b> (Suppl.), 183–190.
25	-13.12	-41.58	Highland caatinga, NE Brazil	Machado, C.G. (2009) Hummingbirds (Aves: Trochilidae) and their floral resources in an area of caatinga vegetation in the Chapada Diamantina, Bahia State, Northeast Brazil. <i>Zoologia</i> , <b>26</b> , 55–65.
26	-13.12	-41.57	Open cerrado, NE	Machado, C.G. (2014) The hummingbird community and the plants which they visit at a savannah in the

ID number	Latitude	Longitude	Site description and general location	Data Source Reference
			Brazil	Chapada Diamantina, Bahia, Brazil. <i>Bioscience Journal</i> , <b>30</b> , 1578–1587.
27	-12.98	-41.33	Campos rupestres, NE Brazil	Machado, C.G., Coelho, A.G., Santana, C.S., Rodrigues, M. (2007) Hummingbirds and their flowers in the 'campos rupestres' of Chapada Diamantina, Bahia, northeastern Brazil. <i>Revista Brasileira de Ornitologia</i> , <b>15</b> , 215–227.
28	10.44	-84.01	Cloud forest, low elevation, Costa Rica	Maglianesi, M.A., Blüthgen, N., Böhning–Gaese, K. & Schleuning, M. (2014) Morphological traits determine specialization and resource use in plant–hummingbird networks in the Neotropics. <i>Ecology</i> , <a href="http://dx.doi.org/10.1890/13-2261.1">http://dx.doi.org/10.1890/13-2261.1</a>
29	10.27	-84.08	Cloud forest, mid elevation, Costa Rica	Maglianesi, M.A., Blüthgen, N., Böhning–Gaese, K. & Schleuning, M. (2014) Morphological traits determine specialization and resource use in plant–hummingbird networks in the Neotropics. <i>Ecology</i> , <a href="http://dx.doi.org/10.1890/13-2261.1">http://dx.doi.org/10.1890/13-2261.1</a>
30	10.18	-84.11	Cloud forest, high elevation, Costa Rica	Maglianesi, M.A., Blüthgen, N., Böhning–Gaese, K. & Schleuning, M. (2014) Morphological traits determine specialization and resource use in plant–hummingbird networks in the Neotropics. <i>Ecology</i> , <a href="http://dx.doi.org/10.1890/13-2261.1">http://dx.doi.org/10.1890/13-2261.1</a>
31	-18.99	-48.30	Cerrado, Central Brazil	Araújo, F.P., Sazima, M. & Oliveira, P.E. (2013) The assembly of plants used as nectar sources by hummingbirds in a Cerrado area of Central Brazil. <i>Plant Systematics and Evolution</i> , <b>299</b> , 1119–1133. Machado, A.O. (2012) Diversidade de recursos florais para beija-flores nos cerrados do Triângulo mineiro e região. <i>Ph.D Thesis</i> . Universidade Federal de Uberlândia, Brazil. Maruyama, P.K., Oliveira, G.M., Ferreira, C., Dalsgaard, B. & Oliveira, P.E. (2013) Pollination syndromes ignored: importance of non-ornithophilous flowers to Neotropical savanna hummingbirds. <i>Naturwissenschaften</i> , <b>100</b> , 1061–1068.
32	-19.16	-48.39	Cerrado, Central Brazil	Araújo, F.P., Sazima, M. & Oliveira, P.E. (2013) The assembly of plants used as nectar sources by hummingbirds in a Cerrado area of Central Brazil. <i>Plant Systematics and Evolution</i> , <b>299</b> , 1119–1133. Machado, A.O. (2012) Diversidade de recursos florais para beija-flores nos cerrados do Triângulo mineiro e região. <i>Ph.D Thesis</i> . Universidade Federal de Uberlândia, Brazil. Maruyama, P.K., Oliveira, G.M., Ferreira, C., Dalsgaard, B. & Oliveira, P.E. (2013) Pollination syndromes ignored: importance of non-ornithophilous flowers to Neotropical savanna hummingbirds. <i>Naturwissenschaften</i> , <b>100</b> , 1061–1068.
33	-17.78	-48.68	Cerrado, Central Brazil	Machado, A.O. (2012) Diversidade de recursos florais para beija-flores nos cerrados do Triângulo mineiro e região. <i>Ph.D Thesis</i> . Universidade Federal de Uberlândia, Brazil.
34	19.50	-96.95	Protected cloud forest, Central Mexico	Ornelas, J.F. <i>Unpublished data</i> .
35	-19.52	-56.98	Pantanal wetland, SW Brazil	Araujo, AC & Sazima, M 2003. The assemblage of flowers visited by hummingbirds in the “capões” of Southern Pantanal, Mato Grosso do Sul, Brazil. <i>Flora</i> , <b>198</b> , 427–435

ID number	Latitude	Longitude	Site description and general location	Data Source Reference
36	18.13	-66.76	Elfin forest, Puerto Rico	Dalsgaard, B., Martín González, A.M., Olesen, J.M., Ollerton, J., Timmermann, A., Andersen, L.H. & Tossas, A.G. (2009) Plant–hummingbird interactions in the West Indies: floral specialisation gradients associated with environment and hummingbird size. <i>Oecologia</i> , <b>159</b> , 757-766.
37	17.95	-66.82	Highly preserved coastal dry scrubland, Puerto Rico	Dalsgaard, B., Martín González, A.M., Olesen, J.M., Ollerton, J., Timmermann, A., Andersen, L.H. & Tossas, A.G. (2009) Plant–hummingbird interactions in the West Indies: floral specialisation gradients associated with environment and hummingbird size. <i>Oecologia</i> , <b>159</b> , 757-766.
38	-19.25	-43.52	Campos rupestres, SE Brazil	Rodrigues, L.C. & Rodrigues, M. (2014) Flowers visited by hummingbirds in the open habitats of the southeastern Brazilian moutaintops: species composition and seasonality. <i>Brazilian Journal of Biology</i> , <b>74</b> , 659–676.
39	-20.44	-54.65	Cerrado, W Brazil	Rodrigues, L.C. & Araujo, A.C. (2011) The hummingbird community and their floral resources in an urban forest remnant in Brazil. <i>Brazilian Journal of Biology</i> , <b>71</b> , 611–622.
40	-22.73	-45.58	Montane Forest, SE Brazil	Sazima, I., Buzato, S. & Sazima, M. (1996) An assemblage of hummingbird-pollinated flowers in a montane forest in southeastern Brazil. <i>Botanica Acta</i> , <b>109</b> , 149–160.
41	10.67	-61.28	Mixed forest, Trinidad	Snow, B.K. & Snow, D.W. (1972) Feeding niches of hummingbirds in a Trinidad Valley. <i>The Journal of Animal Ecology</i> , <b>41</b> , 471–485.
42	4.53	-73.85	Andean forest, Colombia	Snow, D.W. & Snow, B.K. (1980) Relationships between hummingbirds and flowers in the Andes of Colombia. <i>Bulletin of the British Museum of Natural History (Zoology)</i> , <b>38</b> , 105–139.
43	5.90	-73.42	Andean forest, Colombia	Snow, D.W. & Snow, B.K. (1980) Relationships between hummingbirds and flowers in the Andes of Colombia. <i>Bulletin of the British Museum of Natural History (Zoology)</i> , <b>38</b> , 105–139.
44	5.92	-73.53	Andean forest, Colombia	Snow, D.W. & Snow, B.K. (1980) Relationships between hummingbirds and flowers in the Andes of Colombia. <i>Bulletin of the British Museum of Natural History (Zoology)</i> , <b>38</b> , 105–139.
45	-23.63	-45.85	Coastal cloud forest, SE Brazil	Snow D.W. & Snow, B.K. (1986) Feeding ecology of hummingbirds in the Serra do Mar, southeastern Brazil. <i>Hornero</i> , <b>012</b> , 286–296.
46	-17.51	-63.64	Primary forest, Bolivia	Abrahamczyk, S. & Kessler, M. (2010) Hummingbird diversity, food niche characters, and assemblage composition along a latitudinal precipitation gradient in the Bolivian lowlands. <i>Journal of Ornithology</i> , <b>151</b> , 615–625.
47	-16.96	-65.41	Primary forest, Bolivia	Abrahamczyk, S. & Kessler, M. (2010) Hummingbird diversity, food niche characters, and assemblage composition along a latitudinal precipitation gradient in the Bolivian lowlands. <i>Journal of Ornithology</i> , <b>151</b> , 615–625.
48	-19.95	-43.90	Campo rupestre, W Brazil	Vasconcelos, M.F. & Lombardi, J.A. (1999) Padrão sazonal na ocorrência de deis espécie de beija-flores (Apodiformes: Trochilidae) em uma localidade de campo rupestre na Serra do Curral, Minas Gerais. <i>Ararajuba</i> , <b>7</b> , 71–79.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

ID number	Latitude	Longitude	Site description and general location	Data Source Reference
49	-23.28	-45.05	Atlantic forest, SE Brazil	Vizentin–Bugoni, J., Maruyama, P.K. & Sazima, M. (2014) Processes entangling interactions in communities: forbidden links are more important than abundance in a hummingbird–plant network. <i>Proceedings of the Royal Society of London B</i> , <b>281</b> ,1–8.
50	-13.22	-72.12	Subtropical humid montane forest, Perú	Watts, S. (2008) Plant-flower visitor interaction in the Sacred Valley of Perú. <i>PhD. Thesis</i> . University of Northampton, UK.
51	-12.85	-69.37	Lowland primary forest, Perú	Watts, S. (2008) Plant-flower visitor interaction in the Sacred Valley of Perú. <i>PhD. Thesis</i> . University of Northampton, UK.
52	-0.02	-78.77	Andean rainforest, mid-elevation, Ecuador.	Walther, B.A. & Brieschke, H. (2001) Hummingbird-flower relationships in a mid-elevation rainforest near Mindo, northwestern Ecuador. <i>International Journal of Ornithology</i> , <b>4</b> , 115–135.
53	9.57	-83.73	Elfin forest, Costa Rica	Wolf, L.L., Stiles, F.G. & Hainsworth, F.R. (1976) Ecological organization of a tropical, highland hummingbird community. <i>The Journal of Animal Ecology</i> , <b>45</b> , 349–379.
54	9.48	-83.48	Undisturbed highland páramo, Costa Rica	Wolf, L.L., Stiles, F.G. & Hainsworth, F.R. (1976) Ecological organization of a tropical, highland hummingbird community. <i>The Journal of Animal Ecology</i> , <b>45</b> , 349–379.

Peer Review

**Table S2.** Correlations between corrected and observed indices of network structure ( $H_2'$ , complementary specialization; and QuanBiMo, quantitative bipartite modularity) and the predictor variables included in the OLS models: species richness/network size, phylogenetic signal in the interaction pattern of plants ( $d_p$ ) and of hummingbirds ( $d_h$ ), and all included environmental variables (except insularity as it is a dummy variable) calculated within a 10km buffer zone around each locality. MAT, Mean annual temperature; TS, Temperature seasonality; MAP, mean annual precipitation; PS, precipitation seasonality; MAT vel, velocity of temperature change; MAP vel, velocity of precipitation change; MAT anom, temperature anomaly; MAP anom, precipitation anomaly, SE, Sampling effort/intensity. Correlations for the global dataset are below the diagonal (N = 54), and for mainland networks above the diagonal (N = 45). \*\*P<0.01; \*P<0.05 when significance level is based on degrees of freedom corrected for spatial auto-correlation using Dutilleul's (1993) method; †significant when using traditional non-spatial statistics, but non-significant when correcting for spatial auto-correlation; <sup>ns</sup> non-significant. Significant correlations are marked in bold.

	$H_2'$ <i>observed</i>	$H_2'$ <i>corrected</i>	QuanBiMo <i>observed</i>	QuanBiMo <i>corrected</i>	Species richness	$d_p$	$d_h$	MAT	TS	MAP	PS	MAT vel	MAP vel	MAT anom	MAP anom	SE
$H_2'$ <i>observed</i>	-	<b>+0.77**</b>	<b>+0.77**</b>	<b>+0.65**</b>	<b>+0.33*</b>	-0.17 <sup>ns</sup>	<b>-0.43**</b>	<b>+0.42**</b>	-0.29 <sup>ns</sup>	<b>+0.56**</b>	<b>-0.34*</b>	-0.01 <sup>ns</sup>	+0.09 <sup>ns</sup>	<b>-0.47**</b>	+0.31 <sup>†</sup>	-0.28 <sup>ns</sup>
$H_2'$ <i>corrected</i>	<b>+0.82**</b>	-	<b>+0.51**</b>	<b>+0.78**</b>	<b>+0.45*</b>	-0.14 <sup>ns</sup>	<b>-0.37*</b>	+0.23 <sup>ns</sup>	-0.17 <sup>ns</sup>	<b>+0.34*</b>	-0.36 <sup>†</sup>	-0.14 <sup>ns</sup>	+0.19 <sup>ns</sup>	<b>-0.40*</b>	-0.15 <sup>ns</sup>	+0.27 <sup>NS</sup>
QuanBiMo <i>observed</i>	<b>+0.78**</b>	<b>+0.55**</b>	-	<b>+0.76**</b>	<b>+0.38**</b>	<b>+0.32*</b>	<b>-0.30*</b>	<b>+0.34*</b>	-0.43 <sup>†</sup>	<b>+0.61**</b>	-0.27 <sup>ns</sup>	-0.09 <sup>ns</sup>	-0.11 <sup>ns</sup>	<b>-0.43**</b>	+0.40 <sup>†</sup>	<b>-0.41*</b>
QuanBiMo <i>corrected</i>	<b>+0.70**</b>	<b>+0.78**</b>	<b>+0.81**</b>	-	<b>+0.57**</b>	-0.26 <sup>ns</sup>	<b>-0.30*</b>	+0.26 <sup>ns</sup>	-0.27 <sup>ns</sup>	<b>+0.42**</b>	-0.30 <sup>†</sup>	-0.10 <sup>ns</sup>	+0.03 <sup>ns</sup>	<b>-0.38†</b>	+0.25 <sup>ns</sup>	+0.14 <sup>NS</sup>
Species richness	<b>+0.44**</b>	<b>+0.50**</b>	<b>+0.52**</b>	<b>+0.66**</b>	-	-0.23 <sup>ns</sup>	+0.05 <sup>ns</sup>	+0.21 <sup>ns</sup>	<b>-0.41*</b>	<b>+0.31*</b>	+0.01 <sup>ns</sup>	-0.06 <sup>ns</sup>	+0.22 <sup>ns</sup>	-0.34 <sup>†</sup>	+0.13 <sup>ns</sup>	+0.14 <sup>NS</sup>
$d_p$	-0.26 <sup>ns</sup>	-0.25 <sup>ns</sup>	<b>-0.34*</b>	<b>-0.33*</b>	-0.26 <sup>ns</sup>	-	-0.19 <sup>ns</sup>	-0.11 <sup>ns</sup>	+0.21 <sup>ns</sup>	-0.14 <sup>ns</sup>	-0.07 <sup>ns</sup>	+0.11 <sup>ns</sup>	-0.05 <sup>ns</sup>	+0.05 <sup>ns</sup>	-0.07 <sup>ns</sup>	+0.07 <sup>NS</sup>
$d_h$	<b>-0.36**</b>	<b>-0.32*</b>	-0.21 <sup>ns</sup>	-0.24 <sup>ns</sup>	+0.09 <sup>ns</sup>	-0.14 <sup>ns</sup>	-	-0.28 <sup>ns</sup>	-0.05 <sup>ns</sup>	<b>-0.29*</b>	<b>+0.36*</b>	-0.26 <sup>ns</sup>	-0.12 <sup>ns</sup>	+0.12 <sup>ns</sup>	+0.03 <sup>ns</sup>	-0.03 <sup>NS</sup>
MAT	+0.15 <sup>ns</sup>	+0.05 <sup>ns</sup>	+0.06 <sup>ns</sup>	+0.05 <sup>ns</sup>	-0.01 <sup>ns</sup>	+0.01 <sup>ns</sup>	-0.21 <sup>ns</sup>	-	-0.09 <sup>ns</sup>	+0.25 <sup>ns</sup>	+0.02 <sup>ns</sup>	<b>+0.66**</b>	<b>+0.32†</b>	-0.10 <sup>ns</sup>	-0.16 <sup>ns</sup>	+0.00 <sup>NS</sup>
TS	-0.20 <sup>ns</sup>	-0.13 <sup>ns</sup>	-0.30 <sup>†</sup>	-0.20 <sup>ns</sup>	<b>-0.30*</b>	+0.21 <sup>ns</sup>	-0.06 <sup>ns</sup>	-0.17 <sup>ns</sup>	-	-0.47 <sup>†</sup>	+0.14 <sup>ns</sup>	+0.18 <sup>ns</sup>	+0.02 <sup>ns</sup>	+0.17 <sup>ns</sup>	-0.35 <sup>†</sup>	+0.23 <sup>NS</sup>
MAP	<b>+0.51**</b>	<b>+0.34**</b>	<b>+0.50**</b>	<b>+0.36**</b>	+0.25 <sup>ns</sup>	-0.15 <sup>ns</sup>	-0.20 <sup>ns</sup>	+0.25 <sup>ns</sup>	-0.48 <sup>†</sup>	-	<b>-0.47*</b>	-0.14 <sup>ns</sup>	-0.20 <sup>ns</sup>	<b>-0.53**</b>	<b>+0.71**</b>	<b>-0.47*</b>
PS	-0.25 <sup>ns</sup>	-0.28 <sup>†</sup>	-0.15 <sup>ns</sup>	-0.19 <sup>ns</sup>	+0.08 <sup>ns</sup>	-0.03 <sup>ns</sup>	<b>+0.29*</b>	-0.08 <sup>ns</sup>	+0.20 <sup>ns</sup>	<b>-0.50*</b>	-	+0.06 <sup>ns</sup>	-0.03 <sup>ns</sup>	+0.21 <sup>ns</sup>	-0.20 <sup>ns</sup>	+0.12 <sup>NS</sup>
MAT vel	-0.03 <sup>ns</sup>	-0.13 <sup>ns</sup>	-0.05 <sup>ns</sup>	-0.06 <sup>ns</sup>	+0.01 <sup>ns</sup>	+0.19 <sup>ns</sup>	-0.22 <sup>ns</sup>	<b>+0.53**</b>	+0.22 <sup>ns</sup>	-0.20 <sup>ns</sup>	+0.13 <sup>ns</sup>	-	<b>+0.58**</b>	<b>+0.42*</b>	<b>-0.37*</b>	+0.21 <sup>NS</sup>
MAP vel	+0.01 <sup>ns</sup>	+0.07 <sup>ns</sup>	-0.13 <sup>ns</sup>	-0.01 <sup>ns</sup>	+0.17 <sup>ns</sup>	+0.08 <sup>ns</sup>	-0.09 <sup>ns</sup>	<b>+0.29*</b>	+0.05 <sup>ns</sup>	-0.24 <sup>ns</sup>	+0.03 <sup>ns</sup>	<b>+0.61**</b>	-	<b>+0.40**</b>	<b>-0.41*</b>	<b>-0.39*</b>
MAT anom	-0.23 <sup>ns</sup>	-0.13 <sup>ns</sup>	-0.24 <sup>ns</sup>	-0.17 <sup>ns</sup>	-0.16 <sup>ns</sup>	-0.16 <sup>ns</sup>	+0.07 <sup>ns</sup>	-0.14 <sup>ns</sup>	-0.46 <sup>†</sup>	<b>-0.43**</b>	+0.18 <sup>ns</sup>	<b>+0.33*</b>	+0.30 <sup>†</sup>	-	<b>-0.49*</b>	<b>+0.32*</b>
MAP anom	+0.14 <sup>ns</sup>	+0.04 <sup>ns</sup>	-0.15 <sup>ns</sup>	+0.08 <sup>ns</sup>	-0.04 <sup>ns</sup>	-0.03 <sup>ns</sup>	+0.05 <sup>ns</sup>	+0.04 <sup>ns</sup>	-0.37 <sup>†</sup>	<b>+0.67*</b>	+0.27 <sup>†</sup>	<b>+0.38**</b>	<b>+0.38**</b>	<b>-0.44*</b>	-	-0.37 <sup>†</sup>
SE	-0.18 <sup>ns</sup>	<b>+0.29*</b>	<b>-0.33*</b>	+0.16 <sup>NS</sup>	+0.11 <sup>ns</sup>	-0.07 <sup>ns</sup>	-0.07 <sup>ns</sup>	+0.04 <sup>ns</sup>	-0.29 <sup>†</sup>	<b>-0.38*</b>	+0.08 <sup>ns</sup>	+0.15 <sup>ns</sup>	<b>+0.32*</b>	<b>+0.36**</b>	-0.29 <sup>†</sup>	-

**REFERENCES** Dutilleul, P. (1993) Modifying the t test for assessing the correlation between two spatial processes. *Biometrics*, **49**, 305–314.

**Table S3.** Coefficients of determination ( $R^2$ ) from partial regression local models separating the effect of species richness, [phylogenetic signal](#) and environmental factors as predictors of corrected network structure when including climate change velocity as an estimate of historical climate stability. We estimated  $R^2$  values in “full” models, i.e. in models including all ten predictor variables (see Table 1). “Total model” reflects the total amount of variation in network structure explained by the full model; “Total species richness”, “[Total phylogenetic signals](#)” and “Total environmental factors” reflect the variation explained by species richness, phylogenetic signal and environmental factors, respectively; “Only species richness”, “[Only phylogenetic signal](#)”, “Only environmental factors” reflect the amount of variation in network structure explained by the specified predictor-type and not by any other (i.e. the unique variation); Finally, “shared” reflects the variation attributable to the pair of specified predictor types or to all factors combined. See Table 1 for standardized coefficients of each variable and more details of model fit, [and Figure 3 for a graphical representation of these results.](#)

	Global dataset (54 networks)		Only Mainland (45 networks)	
	$H_2'$	QuanBiMo	$H_2'$	QuanBiMo
Total model	0.54	0.62	0.58	0.57
Total species richness	0.25	0.44	0.20	0.33
Total <a href="#">phylogenetic signal</a>	0.19	0.19	0.18	0.19
Total environmental factors	0.38	0.35	0.45	0.32
Unique species richness	0.13	0.24	0.07	0.19
Unique <a href="#">phylogenetic signal</a>	0.06	0.05	0.08	0.07
Unique environmental factors	0.13	0.06	0.22	0.10
Shared species richness – phylogeny	0.00	0.00	0.00	0.00
Shared species richness – environmental factors	0.10	0.14	0.11	0.09
Shared phylogeny – environmental factors	0.10	0.08	0.09	0.07
Shared all factors	0.05	0.08	0.03	0.06

1  
2 **Figure S1.** Scatter plots showing the linear associations between corrected complementary specialization,  $H_2'$ , and corrected quantitative bipartite modularity,  $QuanBiMo$ , and species richness and hummingbird phylogenetic signal, the predictor variables exhibiting the most consistent relationships. Complementary specialization measures the exclusiveness of the interactions in the community, whereas modularity quantifies whether species interact more frequently with subsets of available species within a community. Full and open circles denote mainland and island networks, respectively. Notice the differences in the regression lines fitted to the Global and Only Mainland datasets, in thick and dashed lines, respectively.

