

Genetic Relationship Assessment and Identification of Orchids in the Genus *Eria* Using HAT-RAPD Markers

Jutamas Jiemjuejun, Somchit Damrianant, Theerachai Thanananta
*Department of Biotechnology, Faculty of Science and Technology
Thammasat University, Pathum Thani, Thailand*

Narumol Thanananta*
*Faculty of Science and Technology,
Valaya Alongkorn Rajabhat University under Royal Patronage, Pathum Thani, Thailand*

Received 18 May 2017; Received in revised form 19 September 2017
Accepted 22 September 2017; Available online 15 December 2017

ABSTRACT

High Annealing Temperature-Random Amplified Polymorphic DNA (HAT-RAPD) markers were used to study the genetic relationship among 15 samples of orchids in the genus *Eria*. DNA amplification of 31 random primers out of 72 random primers was successful. In this study, we selected 21 random primers with clear amplified products for further genetic analysis. The result showed significant differences among all samples and also showed specific bands in each sample. Moreover, we found 21 random primers which tended to be DNA markers for further identification. Based on polymorphic bands, a dendrogram was constructed. Genetic similarities among 15 samples of *Eria* species were demonstrated, and the samples could be classified into 3 groups with similarity coefficients ranging from 0.36 to 0.85.

Keywords: *Eria*; Orchids; HAT-RAPD markers; Genetic relationships

Introduction

Orchids are commercially important in Thailand. Export value is expected to increase further because of a wider product range, longer shelf life, and ability to export flowers year round. The orchid family is probably one of the most important plant families from a horticultural point of view. Orchidaceae is one of the largest families among angiosperms. The family includes 800 genera and 25,000 species [1]. Orchids

are distributed in all regions of the world except Antarctica and are found growing in many different habitats and elevation gradients [2]. They are well known for their economic importance and widely cultivated for ornamental purposes and cosmopolitan in distribution [3]. The genus *Eria* is one of the largest and most polymorphous groups of orchids, comprising about 370 taxa [4]. *Eria* plants are among the most prevalent orchids for commercial production of cut

flowers and potted plants and the most popular genus in horticultural industries [4]. The genetic relationships and identification of the *Eria* genus is not well known.

Orchid classification has been poorly understood because of the lack of fossil records, the large number of species, and a historical emphasis on characters related to floral morphology. Circumscription of genera, subtribes, tribes, and subfamilies and the relationships among them were unclear because of homoplasy in morphological and anatomical features [5]. There has been no explicit morphological cladistic analysis of Orchidaceae [6]. Phylogenetic research using DNA sequences is now unraveling the complex picture at every taxonomic level. Assessment of genetic variability is important for the use of genetic resources and for determining the uniqueness of genotypes to exploit heterosis [7]. Genetic relationships can be inferred from genetic similarity or distance estimates determined using phenotypic variation and molecular markers.

Molecular markers are not subject to environmental influence, and because of that, they are considered superior to morphological markers [8]. Several types of molecular markers are available and have been used in assessment of orchid genetic relationships [9], whereas a dendrogram from inter simple sequence repeat (ISSR) was constructed for orchid [10] used the intersimple sequence repeat markers to evaluate genetic relationships and identification among 13 *Dendrobium* species and Tansa-nga *et al.*, [11] determined the genetic relationships among 13 species of *Aerides* species using random amplified polymorphic DNA markers.

The random amplification of polymorphic DNA is relatively simple way of creating genomic DNA 'fingerprints' and can be used successfully to identify orchids. RAPD markers can be useful for the identification and differentiation of species.

Furthermore, high annealing temperature (HAT-RAPD) has been shown to increase the reproducibility of the technique and is used successfully for genetic characterization of numerous plant. Studies on detection of genetic relationship and identification of orchids in the genus *Eria* have not been reported from Thailand. Hence, the present study was undertaken to differentiate *Eria* by using HAT-RAPD markers.

Materials and Methods

Plant Materials

A total of 15 *Eria* species was examined. A list of the *Eria* accessions are provided in Table 1.

Table 1. List of *Eria* species.

No.	Name
1	<i>E. javanica</i> (SW.) Blume
2	<i>E. bractescens</i> Lindl.
3	<i>E. lasiopetala</i> (Willd.) Ormerod
4	<i>E. ornata</i> (Blume) Lindl.
5	<i>E. densa</i> Ridl.
6	<i>E. sarcophylla</i> Schltr.
7	<i>E. biflora</i> (Griff.) Lindl.
8	<i>E. sutepensis</i> Rolfe ex Downie
9	<i>E. cristata</i> Rolfe
10	<i>E. xanthocheila</i> Ridl.
11	<i>E. robusta</i> (Blume) Lindl.
12	<i>E. globulifera</i> Seidenf.
13	<i>E. acervata</i> Lindl.
14	<i>E. discolor</i> Lindl.
15	<i>E. truncata</i> Lindl.

DNA extraction

The genomic DNA of each *Eria* species was extracted from young leaves using the cationic hexadecyl trimethyl ammonium bromide (CTAB) by modified method from Doyle and Doyle [12], and the method from Thanananta *et al.*, [13]. The quality and quantity of extracted DNA samples were assessed by electrophoresis on 0.8 % agarose gel electrophoresis followed

by UV spectrophotometric analysis [14]. The concentration of extracted DNA was adjusted to 100 ng/μl and samples were stored at -20 °C.

HAT-RAPD analysis

Screening of 72 HAT-RAPD primers (Wako Company, Japan) was performed using 15 samples of orchids in the genus *Eria*, only 21 HAT-RAPD primers (A22, A23, A24, A27, A29, A30, A31, A32, B21, B22, B23, B25, B27, B32, C22, C24, C29, C31, D31, E23 and E31) showing strong amplification and good reproducibility were selected. The PCR was performed in a volume of 20 μl containing 100 ng of DNA template, PCR buffer (50 mM KCl, 10 mM Tris-HCl (pH 9.1), 0.1 % Triton™ X-100 and 0.25 mM MgCl₂), 200 μM of dNTPs (dATP, dCTP, dGTP and dTTP), 250 nM of HAT-RAPD primers and *Taq* DNA polymerase (Vivantis technologies Sdn Bhd., Malaysia) 1 U (Unit) [13].

The amplification protocol follows 3 steps; (1) 1 cycle of 94°C for 3 minutes (2) 40 cycles of 94°C for 30 seconds, 46°C for 30 seconds and 72°C for 1 minute and then a final cycle (3) 1 cycle of 72°C for 5 minutes. After the thermal cycling program was complete, stored at 4°C prior to the electrophoresis analysis. The amplification reaction products were at least three PCR amplifications that performed for each sample with HAT-RAPD primers to evaluate the reproducibility of the obtained bands. Amplifications were performed in a thermocycler and PCR products were separated on 1.5% agarose gel electrophoresis and stained with ethidium bromide.

Twenty-one primers were firstly screened with individual of 15 samples of orchids in the genus *Eria*. Primers with reproducible patterns and clear cut polymorphisms were chosen to use for analysis of 15 DNA samples. HAT-RAPD profiles were manually scored as 1 for presence or 0 for absence of a band. The scores were entered into a database program

(NTSYS-pc version 2.0) and a similarity coefficient matrix was constructed. The unweighted pair group method using arithmetic averages (UPGMA) cluster analysis was performed on genetic similarity matrixes, and relationships among species were visualized as a dendrogram using the NTSYS-pc Exeter Software version 2.0 [15].

Results and Discussion

The aim of the present study was to produce HAT-RAPD markers for identification *Eria* species. A total of 72 primers was taken initially and utilized for the amplification from all studied species. Twenty-one HAT-RAPD markers (A22, A23, A24, A27, A29, A30, A31, A32, B21, B22, B23, B25, B27, B32, C22, C24, C29, C31, D31, E23 and E31) were analyzed by PCR. The sequences of these primers are given in Table 2. HAT-RAPD marker was optimized for *Eria* species and produced amplified fragments varying from 100 to 3,000 base pairs (bp) in size for the different primers to be scored across 15 species of orchids in the genus *Eria*. Each primer yielded a wide array of strong and weak bands (Fig. 1). However, only the data of the 21 primers that gave reproducible product formation were included in the statistical analysis.

Table 2. The sequences of primers for HAT-RAPD analysis.

Primers	Sequences
A22	GCCTGCCTCACG
A23	ACTGACCTAGTT
A24	CTCCTGCTGTTG
A27	ATCGCGGAATAT
A29	GGTTCGGGAATG
A30	GACCTGCGATCT
A31	AAGGCGCGAACG
A32	TTGCCGGGACCA

Table 2. The sequences of primers for HAT-RAPD analysis. (Continued)

Primers	Sequences
B21	AAGCCTATACCA
B22	GGTGACTGGTGG
B23	GGTGCCGGAGCA
B25	AGCACTGAATCT
B27	GGCGTTATGAA
B32	ATCGCGGCTTAT
C22	GGTCACCGATCC
C24	CCTTGGCATCGG
C29	GTCGCCTTACCA
C31	TCTGCTGACCGG
D31	GGAGGTCGACCA
E23	AGGTACGCCGCA
E31	GAGGACAGCAA

The genetic similarity coefficients for the 15 species of orchids in the genus *Eria* ranged from 0.36 to 0.65 (Fig. 3). The UPGMA analysis made it possible to cluster all of the species. The dendrogram (Fig. 2) showed a clear distinction into 3 major groups.

The first group was *E. discolor* Lindl. The second group included *E. bractescens* Lindl. and *E. truncata* Lindl. And the third group could be divided into three subgroups. The first subgroup was composed of *E. javanica* (SW.) Blume, *E. densa* Ridl. and *E. sarcophylla* Schltr. The second subgroup contained *E. lasiopetala* (Willd.) Ormerod, *E. ornata* (Blume) Lindl., *E. cristata* Rolfe, *E. sutepensis* Rolfe ex Downie and *E. biflora* (Griff.) Lindl. Within this second subgroup, the *Eria* genus named *E. lasiopetala* (Willd.) Ormerod paired with *E. ornata* (Blume) Lindl. at the closest genetic similarity coefficients of 0.65. And the third subgroup was composed of *E. xanthocheila* Ridl., *E. robusta* (Blume)

Lindl., *E. globulifera* Seidenf. and *E. acervata* Lindl.

From the results, *E. lasiopetala* (Willd.) Ormerod is most closely related to *E. ornata* (Blume) Lindl. (Fig. 2) and appears indistinguishable from it externally. Both taxa share similar features such as slenderly clavate stems topped by a few leaves, biflorous inflorescences with relatively small floral bracts, pubescent flowers with a prominent subglobose mentum, same color, and a more or less subquadrate lip [16]. Not only the HAT-RAPD markers those are effective to identify the *Eria* species but they can also be used to detect other types of orchids such as, *Coelogyne* (Orchidaceae) [17], *Aerides* [18], *Dendrobium* Section *Nigrohirsutae* [19], *Bulbophyllum* Section *Sestochilus* [20] and *Dendrobium* spp. (Ueang Sai group) [21]. Furthermore, The HAT-RAPD approach for genetic characterization is highly sensitive and even scarce amount of DNA template are sufficient for analysis. HAT-RAPD markers have been successfully used for identification of various plant species such as, banana [22], Lychee [23], *Ficus* spp. [24], *Brassica* spp. [25] and *Curcuma* [26].


Conclusion

This paper, study for the characterization of genetic relationship among 15 species of orchids in the genus *Eria* based on HAT-RAPD markers. The genetic similarity coefficients for the 15 species ranged from 0.36 to 0.65. The dendrogram showed a clear distinction within the 3 major groups. This study shows that the molecular markers could be utilized for the comparison of genetic relationship in orchid species, which their morphological

descriptions are most similar. It also indicates that the HAT-RAPD markers may be considered as an effective and convenient method to classify *Eria* species. In conclusion, this research has demonstrated the HAT-RAPD markers as a useful tool for DNA polymorphisms detection, to examine genetic relationship of orchids in the genus *Eria*. The present study highlights the

importance of HAT-RAPD markers. It has been widely used as a genetic screening method [27] because it is rapid, relatively simple to be performed, and required only a small amount of genomic DNA without genome sequence information prior to analysis [28]. Furthermore, it can also be used for the study of genetic diversity of orchid species.

A.



B.




Fig. 1 Examples of HAT-RAPD patterns amplified A32 (A) and B23 (B) primers of the *Eria* 15 species. [Lane M is molecular weight marker 100 bp plus DNA Ladder (Vivantis, Vivantis technologies Sdn. Bhd., Malaysia), Lanes 1-15, (1) *E. javanica* (SW.) Blume (2) *E. bractescens* Lindl. (3) *E. lasiopetala* (Willd.) Ormerod (4) *E. ornata* (Blume) Lindl. (5) *E. densa* Ridl. (6) *E. sarcophylla* Schltr. (7) *E. biflora* (Griff.) Lindl. (8) *E. sutepensis* Rolfe ex Downie (9) *E. cristata* Rolfe (10) *E. xanthocheila* Ridl. (11) *E. robusta* (Blume) Lindl. (12) *E. globulifera* Seidenf. (13) *E. acervata* Lindl. (14) *E. discolor* Lindl. and (15) *E. truncata* Lindl.]

References

- [1] Stewart J, Griffiths M. Manual of orchids. Portland: Timber Press; 1995.
- [2] Pridgeon A. The illustrated encyclopedia of orchids. Portland: Timber Press; 2000.
- [3] Bechtel H, Cribb P, Launert E. The manual of cultivated orchid species. 3rd ed. London: Blandford Press; 1992.
- [4] Mabberley DJ. Mabberley00 Plant Book. 3rd ed. Cambridge: Cambridge University Press; 2008.
- [5] Chase M. Royal Botanic Gardens [Internet]. Richmond, UK: Kew Science & Horticulture; 2006 [cited 2013 Jan 11]. Available from: <http://www.Orchidaceae.html/>.
- [6] Freudenstein JV, Rasmussen FN. What does morphology tell us about orchid relationships—a cladistic analysis. *Amer J Bot* 1999;86:225-48.
- [7] Franco J, Crossa J, Ribaut JM, Betran J, Warburton ML, Khairallah M. A method for combining molecular markers and phenotypic attributes for classifying plant genotypes. *Theor Appl Genet* 2001;103:944-52.
- [8] Ma'ric S, Lari'c S, Artincic J, Peji'c I, Kozumplink V. Genetic diversity of hexaploid wheat cultivars estimated by RAPD markers, morphological traits and coefficients of parentage. *Plant Breed* 2004;123:366-9.
- [9] Chiang CH, Yu TA, Lo SF, Kuo CL, Peng WH, Tsay HS. Molecular authentication of *Dendrobium* species by multiplex polymerase chain reaction and amplification refractory mutation system analysis. *J Amer Soc Hort Sci* 2012;137:438-44.
- [10] Thomsopa T, Thanananta T, Thanananta N. Identification and Genetic Relationship Analysis of *Dendrobium* spp., Ueang Sai group, Using HAT-RAPD and ISSR Techniques. *Thai J Sci Technol* 2014;2:82-91.
- [11] Tansa-nga W, Thanananta T, Thanananta N. Identification and Genetic Relationship Analysis of *Aerides* Varieties Using High Annealing Temperature-Random Amplified Polymorphic DNA (HAT-RAPD) and Inter-Simple Sequence Repeat (ISSR) Techniques. *Thai J Sci Technol* 2014;2:102-12.
- [12] Doyle JJ, Doyle JL. A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochem Bull* 1987;19:11-5.
- [13] Thanananta N, Prasit V, Thanananta T. Identification of Rice Cultivars KDML105 and Its Improved Cultivars by Using HAT-RAPD Technique. *Thai J Sci Technol* 2012;3:169-79.
- [14] Sambrook J, Fritsch EF, Maniatis T. Molecular Cloning: A Laboratory Manual. 2nd ed. New York: Cold Spring Harbor Laboratory; 1989.
- [15] Rohlf FJ. NTSYSpc Numerical Taxonomy and Multivariate Analysis System. New York: Applied Biostatistics; 2002.
- [16] Agrawala DK. Taxonomic studies on the genus *Eria* Lindl. (Orchidaceae) in India [Ph.D. Thesis]. West Bengal: Kalyani University; 2009.
- [17] Thanananta N, Hongtongdee P, Thanananta T. Identification and Assessment of Genetic Relationship Among *Coelogyne* (Orchidaceae) Using HAT-RAPD. *Thai J Sci Technol* 2016;1:88-97.
- [18] Tansa-nga W, Thanananta T, Thanananta N. Identification and Genetic Relationship Analysis of *Aerides* Varieties Using High Annealing Temperature-Random Amplified Polymorphic DNA (HAT-RAPD) and Inter-Simple Sequence Repeat (ISSR) Techniques. *Thai J Sci Technol* 2014;2:102-12.
- [19] Phanroophaw J, Thanananta T, Thanananta N. Genetic Relationships between *Dendrobium* Section *Nigrohirsutae* and their Hybrids base on HAT-RAPD. *Thai J Sci Technol* 2016;1:77-87.
- [20] Saetai K, Thanananta T, Thanananta N. Identification and Genetic Relationship Analysis of *Bulbophyllum* Section *Sestochilus* Using HAT-RAPD and ISSR Markers. *Thai J Sci Technol* 2014;2:92-101.
- [21] Thomsopa T, Thanananta T, Thanananta N. Identification and Genetic Relationship Analysis of *Dendrobium* spp., Ueang Sai group, Using HAT-

- RAPD and ISSR Techniques. Thai J Sci Technol 2014;2:82-91.
- [22] Ruangsuttapha S, Eimert K, Schroder MB, Silayoi B, Denduangboripant J, Kanchanapoom K. Molecular phylogeny of banana cultivars from Thailand based on HAT-RAPD markers. Genet Resour Crop Evol 2007;54:1565-72.
- [23] Chundet R, Cutler RW, Tasanon M, Anuntalabhochai S. Hybrid Detection in Lychee (*Litchee chinensis* Sonn.) Cultivars Using HAT-RAPD Markers. Sci Asia 2007;33:307-11.
- [24] Phromthep W. A new genetic analysis of *Ficus* spp. By HAT-Random amplified Polymorphic DNA Technique, Procedia Engineering. 2012;32:1073-9.
- [25] Sumrith J, Kunpuksri P, Thanananta T. Genetic Relationship Analysis and Identification of *Brassica* spp. Using HAT-RAPD Technique. Thai J Sci Technol 2014;1:23-8.
- [26] Tansa-nga W, Thanananta T, Chongkid B, Thanananta N. Genetic Relationship Analysis and Identification of *Curcuma* Using HAT-RAPD Technique. Thai J Sci Technol 2014;1:29-35.
- [27] Gasser RB. Molecular tools advances opportunities and prospects. Vet Parasitol 2006;136:69-89.
- [28] Williams JG, Kubelik AR, Livak KJ, Rafalski JA, Tingey SV. DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. Nucleic Acids Res 1990;18:6531-5.