


FRESCO™

Brachyscome angustifolia



BRAND
NEW 23



- Large flowers
- Perfect for combinations
- Easy in propagation
- Mounded habit is well suited for hanging baskets and window boxes

FRESCO™ Bright Pink

EU & ROW
ONLY

More varieties in the FRESCO™ series:



FRESCO™ Candy



FRESCO™ Purple

For additional technical growing information:

FRESCO™ [CLICK HERE](#)



FRESCO™

Brachyscome angustifolia



PROPAGATION TIPS

- Stick on priority – Number 2 out of 4 categories



Flowering time

Spring / Summer
Flowering



Width

(8 - 10")
20 - 25 cm



Height

(6 - 8")
15 - 20 cm

Average Time	Temperature	Hormone	Fertilization	Fungicide
4 weeks	Weeks 1 - 2 70°-72° F (21° - 22°C)	Optional	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 5 68° F (18° C)		Weeks 3-5 75 to 100 ppm N	Day of sticking & Week 2



Irrigation specification - Remove from mist as soon as possible



High light and warm temperatures will maintain a good proportion of flowers to foliage. Maintain lower EC levels for best growth and flower size since higher EC will hinder plant quality.

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.6 / 6.2	0.80	72°-74° F (22°-23° C)	50 ppm N in mist			Day of sticking	
Week 2	5.6 to 6.2	.80 to .90	72°-74° F (22°-23° C)	50 ppm N in mist		750 ppm Daminozide	Second fungicide app	
Week 3	5.6 to 6.2	.90 to 1.0	Cool to 68°-70° F (20°-21° C)	Feed 100 ppm to 150 ppm		Daminozide as needed		
Week 4	5.6 to 6.2	.90 to 1.0	65°-68° F (18°-20° C)	feed 100 to 150 ppm		Daminozide as needed		

For additional technical growing information:

FRESCO™ [CLICK HERE](#)



 **DANZIGER**
www.danzigeronline.com | 

FRESCO™

Brachyscome angustifolia



FINISHING TIPS

- Pinch week 3 in the liner or at transplant
- Monitor EC levels so they are not elevated
- Do not over irrigate early in plant finishing

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
6 to 8 wks (12 cm) 5" pot	Average Day 68°-72° F (20° -22° C)	Pinch No lighting required for flowering	100 - 150 ppm N	In liner and finish stages
8 to 10 wks (15 cm) 6 ½"	Weeks 3 – 5 61°-68° F (16°-20° C)		Soil EC 0.8 - 1.0 pH 5.6 to 6.2	

Pests	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS
	Thrips	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
Diseases	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRODIONE, POLYOXIN
	Pythium	PROPAMOCARB
	Phytophthora	PROPAMOCARB



- Avoid covering plant meristem at transplant
- Additional Iron (Fe) or Manganese (Mn) may be required, depending on water quality
- PGR - Spray Daminozide at 750 to 2000 ppm on young & finished plants
- Maintain good airflow and allow plants to dry before nightfall

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 5 Transplant	5.6 to 6.2	0.8 to 1.0	65°-68° F (18°-20° C)	feed 100 to 150 ppm	Day Neutral	Daminozide as needed	Spray fungicide after transplant	
Week 5 to 13	5.6 to 6.2	0.8 to 1.0	65°-68° F (18°-20° C)	feed 100 to 150 ppm	Not Required			

For additional technical growing information:

FRESCO™ [CLICK HERE](#)

